

હવામાન આગાહી આધારીત કૃષિ સલાહ બુલેટીન
દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ
તા. ૦૫-૦૪-૨૦૧૫ થી તા. ૦૮-૦૪-૨૦૧૫

(ગ્રામીણ કૃષિ મૌસમ સેવા, જુનાગઢ કૃષિ યુનિવર્સીટી, જુ.કૃ.યુ.)

હવામાન આગાહી આધારીત બુલેટીન : ૨૦૧૫

પાછલા અઠવાડીયાનું જુનાગઢનું હવામાન

હવામાન પરીબળો	દક્ષિણ સૌરાષ્ટ્ર કૃષિ આબોહવાકીય વિસ્તાર
વરસાદ (મીમી)	૦૦.૦
મહત્તમ તાપમાન (સે.ગે.)	૩૭.૦
લઘુતમ તાપમાન (સે.ગ્રે.)	૧૯.૦
ભેજ(%) (સવાર/બપોરબાદ)	૭૬-૨૮
પવનની ગતિ (કિમી/કલાક)	૫.૪

પત્રક-બ : જિલ્લાવાર હવામાન આગાહી (તા.૦૯-૦૪-૧૫ સુધી)

હવામાન પરીબળો	તારીખ	દક્ષિણ સૌરાષ્ટ્ર	
		જુનાગઢ	પોરબંદર
વરસાદ (મીમી)	૦૫.૦૪.૧૫	૦	૦
	૦૬.૦૪.૧૫	૦	૦
	૦૭.૦૪.૧૫	૦	૦
	૦૮.૦૪.૧૫	૦	૦
	૦૯.૦૪.૧૫	૦	૦
મહત્તમ તાપમાન (સે.ગ્રે.)	૦૫.૦૪.૧૫	૩૨	૩૧
	૦૬.૦૪.૧૫	૩૨	૩૧
	૦૭.૦૪.૧૫	૩૨	૩૦
	૦૮.૦૪.૧૫	૩૩	૩૧
	૦૯.૦૪.૧૫	૩૩	૩૧
લઘુત્તમ તાપમાન (સે.ગ્રે.)	૦૫.૦૪.૧૫	૨૬	૨૫
	૦૬.૦૪.૧૫	૨૬	૨૫
	૦૭.૦૪.૧૫	૨૬	૨૫
	૦૮.૦૪.૧૫	૨૬	૨૬
	૦૯.૦૪.૧૫	૨૬	૨૬
વાદળની સ્થિતિ (ઓકટા)	૦૫.૦૪.૧૫	મોટેભાગે ચોખ્ખુ	મોટેભાગે ચોખ્ખુ
	૦૬.૦૪.૧૫	મોટેભાગે ચોખ્ખુ	ચોખ્ખુ
	૦૭.૦૪.૧૫	મોટેભાગે ચોખ્ખુ	મોટેભાગે ચોખ્ખુ
	૦૮.૦૪.૧૫	મોટેભાગે ચોખ્ખુ	મોટેભાગે ચોખ્ખુ
	૦૯.૦૪.૧૫	ચોખ્ખુ	ચોખ્ખુ
મહત્તમ ભેજ (%)	૦૫.૦૪.૧૫	૮૩	૮૩
	૦૬.૦૪.૧૫	૮૧	૮૧
	૦૭.૦૪.૧૫	૮૦	૭૯
	૦૮.૦૪.૧૫	૭૮	૭૮
	૦૯.૦૪.૧૫	૮૦	૮૦
લઘુત્તમ ભેજ (%)	૦૫.૦૪.૧૫	૭૨	૭૨
	૦૬.૦૪.૧૫	૬૯	૬૯
	૦૭.૦૪.૧૫	૬૧	૬૧
	૦૮.૦૪.૧૫	૬૫	૬૪
	૦૯.૦૪.૧૫	૬૩	૬૨
પવનની ગતિ (કિમી/કલાક)	૦૫.૦૪.૧૫	૦૧૭	૦૧૭
	૦૬.૦૪.૧૫	૦૧૫	૦૧૫
	૦૭.૦૪.૧૫	૦૨૧	૦૨૧
	૦૮.૦૪.૧૫	૦૧૪	૦૧૪
	૦૯.૦૪.૧૫	૦૧૮	૦૧૯
પવનની દિશા (ડીગ્રી)	૦૫.૦૪.૧૫	૨૮૦	૨૭૦
	૦૬.૦૪.૧૫	૨૯૦	૨૯૦
	૦૭.૦૪.૧૫	૨૯૦	૨૯૦
	૦૮.૦૪.૧૫	૨૮૦	૨૯૦
	૦૯.૦૪.૧૫	૩૦૦	૩૦૦

દક્ષિણ ખેત આબોહવાકિય વિસ્તાર (જિલ્લાવાર કૃષિ સલાહ)

પાકનું નામ	કૃષિ સલાહ
ઉનાળુ મગફળી	<ul style="list-style-type: none"> ❖ પાકમાં થ્રીપ્સના નિયંત્રણ માટે પ્રોફેનોફોસ ૨૦ મીલી દવાનો ૧૦ લીટર પાણી ઓગાળીને છંટકાવ કરવો. ❖ વહેલા લાગતા પાનના ટપકાના રોગના નિયંત્રણ માટે ૩૦૦ ગ્રામ કાર્બેન્ડાઝીમ ૧૨ % + મેંકોઝેબ ૬૩ % wp /એકર/ ૧૫૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો. ❖ વધુ પડતા નાઈટ્રોજનયુક્ત ખાતર આપવાથી વાનસ્પતિક વૃદ્ધિ થાય છે અને ઉત્પાદન ઘટે છે. ખોટો ખર્ચ નિવારવા પુર્તિ ખાતરો ના આપવા ❖ પાકને જરૂરીયાત મુજબ પિયત આપવું. ❖ પાકમાં સમયાંતરે આંતરખેડ અને નિંદામણ કરવું.
ઉનાળુ તલ	<ul style="list-style-type: none"> ❖ બીજને વાવતા પહેલા થાઈરમ અથવા મેંકોઝેબ અથવા કાર્બેનડેન્ઝીમ દવાનો ૫૮ પ્રતિ કીલોગ્રામ બીજ દીઠ ૨ થી ૩ ગ્રામ મુજબ આપવો. ❖ પાકને જરૂરીયાત મુજબ પિયત આપવું.
ઉનાળુ મગ	<ul style="list-style-type: none"> ❖ પાયાના ખાતર તરીકે ૧૪.૦ કીલો ડી.એ.પી અને ૧.૬ કીલો યુરીયા પ્રતી વીધા મુજબ આપવું. ❖ પાકને જરૂરીયાત મુજબ પિયત આપવું. ❖ પાકમાં સમયાંતરે આંતરખેડ અને નિંદામણ કરવું.
લસણ	<ul style="list-style-type: none"> ❖ પાન ની ટોચના સુકારા રોગના નિયંત્રણ માટે મેંકોઝેબ દવા ૨૫ ગ્રામ ને ૧૦ લીટર પાણી ઓગાળીને છંટકાવ કરવો. ❖ પાકમાં થ્રીપ્સના નિયંત્રણ માટે પ્રોફેનોફોસ ૨૦ મીલી દવાનો ૧૦ લીટર પાણી ઓગાળીને છંટકાવ કરવો.
દિવેલા	<ul style="list-style-type: none"> ❖ ઘોડીયા ઈયળના નિયંત્રણ માટે ક્વીનાલફોસ ૨૦ મીલી ૧૦ લીટર પાણીમા મીશ્ર કરી છંટકાવ કરવો
શાકભાજી	<ul style="list-style-type: none"> ❖ સ્પોટેડ વીલ્ટ વાયરસના નિયંત્રણ માટે પાકની રોપણી પછી ૧૦ થી ૧૨ દિવસના અંતરે ૫ ગ્રામ કાર્બેન્ડાઝીમ (૫૦ % વે.પા.) ૧૦ લિટર પાણીમા ઓગાળી ને છાંટવું જેથી રોગનું પ્રમાણ ઓછું રહે છે અને ઉત્પાદન વધારે મળે છે. ❖ પાક એક થી દોઢ માસનો થાય ત્યારે ૫૦ % વે.પા. કાર્બેન્ડાઝીમ ૧૦ ગ્રામ અથવા ૭૫ % વે.પા. મેન્કોઝેબ ૨૫ ગ્રામ અથવા ૫૦ % તામ્બાચુક્ત દવા ૫૦ ગ્રામ ૧૦ લીટર પાણીમા ઓગાળી ત્રણ છંટકાવ કરવાથી પાનના ટપકા તથા ફળના સડાનો રોગનું નિયંત્રણ કરી શકાય છે.

આબો	<ul style="list-style-type: none">❖ મધ્યાના નિયંત્રણ માટે ઈમીડાક્લોપ્રીડ ૫ મીલી ૧૦ લીટર પાણીમાં નાખીને છાટવી ૧૦ દીવસના અંતરે બે છંટકાવ કરવા.❖ પાકમાં ભુકીછારાના નિયંત્રણ માટે હેક્ઝાકોનાઝોલ ૧૫ મીલી દવાનો ૧૦ લીટર પાણી ભેળવીને છંટકાવ કરવો.
પશુ પાલન	<ul style="list-style-type: none">❖ પશુઓને યોગ્ય આહાર આપવો.❖ પશુઓમાં ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટાઝ ૨ મીલી/ ૧ લીટર પાણીમાં નાખીને છાટવી.❖ પશુમાં કુટ અને માઉથ રોગના આવે તે માટે જરૂરી રશીકરણ કરાવવું.

JUNAGADH

Phone: 0285-2670139

Date:- 04-04-2015

(Period from 05 to 09 April-2015)

Significant weather for the preceding week

Rain fall (mm)	000.0
Maximum temperature °C	37.0-37.2
Minimum temperature °C	16.5-20.5
RH % (Morning/Afternoon)	76-28
Wind speed (Kmph)	4.2-7.2
Cloud cover	0-4

Weather forecast for Junagadh Dist. until 08:30 hrs of 09-04-2015

Parameters	Day-1 05-04-15	Day-2 06-04-15	Day-3 07-04-15	Day-4 08-04-15	Day-5 09-04-15
Rain fall (mm)	0	0	0	0	0
Maximum temperature °C	32	32	32	33	33
Minimum temperature °C	26	26	26	26	26
Cloud cover (Okta)	1	1	1	1	0
RH % (Morning/After noon)	83-72	81-69	80-61	78-65	80-63
Wind speed (Kmph)	017	015	021	014	018
Wind direction(Deg)	280	290	290	280	300

Agro meteorological Advisories

Crop	Stage	Agromet Advisories
Summer Groundnut	Growth	<ul style="list-style-type: none"> For the control of thrips, spray Profenophos @ 20 ml in 10 liter of water. For the control of early leaf spot, spray 300gm Carbendazim 12 + Mencozeb 63 % WP / acre/ 150 liter of water. Apply irrigation as and when required. Interculturing and hand weeding should be applied.
Summer Sesame	Sowing/ Germination	<ul style="list-style-type: none"> Seed should be treated with thiram or mancozeb or Carbendazim @ 2 to 3g per 1 kg of seed before sowing. Apply irrigation as and when required.
Summer Mung	Sowing/ Germination	<ul style="list-style-type: none"> The basal dose fertilizer should be applied @ 14.0kg DAP /Vigha and 1.6 kg Urea/vigha. Interculturing and hand weeding should be applied.
Garlic	Bulb development	<ul style="list-style-type: none"> For the control of white tip, spray Mencozeb 25 ml/10 liter of water. For the control of thrips, spray Profenophos 20ml in 10 liter of water.
Castor	Capsule formation	<ul style="list-style-type: none"> For the control of semi lopper, spray quinalphos 20 ml in 10 liter of water.

Vegetables	Flowering/Fruiting	<ul style="list-style-type: none"> • For the control of spotted wilt virus, spray Carbendazim (50 % WP) 5gm in 10 liter of water at 10 to 12 days interval. • For the control of fruit rot, three spray of 50 WP carbendazim 10 gm or 75 WP Mencozeb 25 gm in 10 liter of water at 10 to 12 days interval.
Mango	Flowering/Fruiting	<ul style="list-style-type: none"> • For the control of leaf hopper, spray Imidaclopride 5 ml in 10 liter of water. • For the control of powdery mildew, spray Hexaconazol@50ml in 10 liter of water.
Livestock	-	<ul style="list-style-type: none"> • Do vaccination in animal for the prevention of FMD disease. • The beaks of bird in poultry under the age of 15 weeks should be cut.

PORBANDAR

Phone : 0285-2670139

Date :04-04-2015

(Period from 05 to 09 April-2015)

Weather forecast for Junagadh Dist. until 08:30 hrs of 09-04-2015

Parameters	Day-1 05-04-15	Day-2 06-04-15	Day-3 07-04-15	Day-4 08-04-15	Day-5 09-04-15
Rain fall (mm)	0	0	0	0	0
Maximum temperature °C	31	31	30	31	31
Minimum temperature °C	25	25	25	26	26
Cloud cover (Okta)	1	0	1	1	0
RH % (Morning/After noon)	83-72	81-69	79-61	78-64	80-62
Wind speed (Kmph)	017	015	021	014	019
Wind direction(Deg)	270	290	290	290	300

Agro meteorological Advisories

Crop	Stage	Agromet Advisories
Summer Groundnut	Growth	<ul style="list-style-type: none"> For the control of thrips, spray Profenophos @ 20 ml in 10 liter of water. For the control of early leaf spot, spray 300gm Carbendazim 12 + Mencozeb 63 % WP / acre/ 150 liter of water. Apply irrigation as and when required. Interculturing and hand weeding should be applied.
Summer Sesame	Sowing/ Germination	<ul style="list-style-type: none"> Seed should be treated with thiram or mancozeb or Carbendazim @ 2 to 3g per 1 kg of seed before sowing. Apply irrigation as and when required.
Summer Mung	Sowing/ Germination	<ul style="list-style-type: none"> The basal dose fertilizer should be applied @ 14.0kg DAP /Vigha and 1.6 kg Urea/vigha. Interculturing and hand weeding should be applied.
Garlic	Bulb development	<ul style="list-style-type: none"> For the control of white tip, spray Mencozeb 25 ml/10 liter of water. For the control of thrips, spray Profenophos 20ml in 10 liter of water.
Castor	Capsule formation	<ul style="list-style-type: none"> For the control of semi lopper, spray quinalphos 20 ml in 10 liter of water.
Vegetables	Flowering/Fruiting	<ul style="list-style-type: none"> For the control of spotted wilt virus, spray Carbendazim (50 % WP) 5gm in 10 liter of water at 10 to 12 days interval. For the control of fruit rot, three spray of 50 WP carbendazim 10 gm or 75 WP Mencozeb 25 gm in 10 liter of water at 10 to 12 days interval.
Mango	Flowering/Fruiting	<ul style="list-style-type: none"> For the control of leaf hopper, spray Imidaclopride 5 ml in 10 liter of water.

		<ul style="list-style-type: none">• For the control of powdery mildew, spray Hexaconazol@50ml in 10 liter of water.
Livestock	-	<ul style="list-style-type: none">• Do vaccination in animal for the prevention of FMD disease.• The beaks of bird in poultry under the age of 15 weeks should be cut.