



ગ્રામીણ કૃષિ મૌસમ સેવા
કૃષિ હવામાન સેલ, કૃષિ વિજ્ઞાન વિભાગ
જુનાગઢ કૃષિ યુનિવર્સિટી
જુનાગઢ - ૩૬૨૦૦૧

ઈમેઇલ:- amfujind@yahoo.com ફોન નં. ૦૨૮૫-૨૬૭૦૧૩૮



હવામાન આગાહી આધારીત દક્ષિણ સૌરાષ્ટ્ર પેટ આબોહવાકીય વિસ્તારનાં જુનાગઢ જીલ્લા માટે કૃષિ સલાહ બુલેટીન
(કૃષિ હવામાન સેલ, જુનાગઢ કૃષિ યુનિવર્સિટી અને ભારત હવામાન વિભાગ દ્વારા સંપત્તા)

(#આગાહી નો અમૃત મહોત્સવ#)

વાર:- મંગળવાર

તારીખ:- ૧૧/૦૮/૨૦૨૧

હવામાન આગાહી આધારીત બુલેટીન: ૦૩૮ (૨૦૨૧-૨૨)

પાછલા પાંચ દિવસનું હવામાન (તારીખ:- ૦૫ થી ૦૯ ઓગસ્ટ, ૨૦૨૧)					હવામાન પરિબળો	હવામાન આગાહી આગામી પાંચ દિવસ માટે (તારીખ:- ૧૧ થી ૧૫ ઓગસ્ટ, ૨૦૨૧) માન્ય ૧૫ ઓગસ્ટ, ૨૦૨૧. સવારના ૮:૩૦ વાગ્યાં સંધી				
૦૫/૦૮	૦૬/૦૮	૦૭/૦૮	૦૮/૦૮	૦૯/૦૮	તારીખ	૧૧/૦૮	૧૨/૦૮	૧૩/૦૮	૧૪/૦૮	૧૫/૦૮
૦	૦	૦	૦	૦	વરસાદ (મિમી)	૨	૩	૩	૩	૩
૩૨.૫	૩૩.૮	૩૨.૬	૩૨.૭	૩૪.૫	મહત્વમાન તાપમાન (°સે.ગ્ર)	૩૪	૩૪	૩૩	૩૩	૩૪
૨૪.૭	૨૫.૨	૨૪.૬	૨૪.૨	૨૪.૦	લઘુત્વમાન તાપમાન (°સે.ગ્ર)	૨૫	૨૫	૨૫	૨૬	૨૬
મોટેભાગેવ વાદળથાયું	વાદળથાયું	મોટેભાગેવ વાદળથાયું	મોટેભાગેવ વાદળથાયું	મોટેભાગેવ વાદળથાયું	વાદળની સ્થિતિ	મોટેભાગે વાદળથાયું	વાદળથાયું	વાદળથાયું	વાદળથાયું	વાદળથાયું
૮૩	૮૮	૮૧	૮૧	૮૩	સવારના લેજના ટકા	૭૮	૮૨	૮૨	૮૩	૮૩
૫૪	૬૨	૫૧	૫૧	૭૮	બપોરબાદ ના લેજના ટકા	૫૮	૬૪	૬૬	૬૮	૬૮
૧૧.૩	૧૦.૫	૬.૦	૭.૦	૬.૭	પવનની ગતિ (કિમી/કલાક)	૨૩.૨	૨૪.૩	૨૩.૧	૨૪.૫	૨૫.૭
૨૫૦	૨૫૦	૨૫૦	૨૩૦	૨૭૦	પવનની દિશા	૨૪૮	૨૪૮	૨૪૮	૨૪૮	૨૪૮


સામાન્ય સલાહ

ભારત સરકારશ્રીના હવામાન વિભાગ તરફથી મળેલ હવામાન આગાહીની માહીતી અનુસાર અત્રેના ગ્રામીણ કૃષિ મૌસમ સેવા યોજના, જુનાગઢ કૃષિ યુનિવર્સિટી, જુનાગઢ તરફથી જણાવવામા આવે છે કે દક્ષિણ સૌરાષ્ટ્ર પેટ આબોહવાકીય વિસ્તારના જુનાગઢ જીલ્લામાં આગામી તારીખ ૧૧/૦૮/૨૦૨૧ થી ૧૫/૦૮/૨૦૨૧ દરમયાન આકાશ મુખ્યત્વે આગામી પાંચ દિવસોમાં મોટેભાગે વાદળથાયું થી વાદળથાયું રહેશે, જીલ્લામાં આગામી પાંચ દિવસોમાં છુટા છવાયા ભાગોમાં હળવા / આપટાં વરસાદ ની શક્યતાઓ છે. પવનની ઝડપ ૨૩.૧ થી ૨૫.૭ કિમી/કલાક રહેવાની શક્યતા રહેશે. આ સમયગાળામાં મહત્વમાન ૩૩ થી ૩૪ ડિગ્રી સેન્ટીગ્રેડ રહેશે તેમજ લઘુત્વમાન ૨૫ થી ૨૬ ડિગ્રી સેન્ટીગ્રેડ રહેવાની સંભાવના છે.

- માન્ય વરસાદની સૂચી મુજબ જુનાગઢ જીવલામાં ૦૮ જુલાઈ થી ૦૪ ઓગસ્ટ, ૨૦૨૧ ના સમયગાળા દરમિયાન હળવી ભેજ સ્થિતિ જોવા મળેલ છે.
- સાપ્તાહિક વરસાદ આગાહીની સૂચી મુજબ દક્ષિણ સૌરાષ્ટ્રનાં વિસ્તારમાં તા. ૧૫/૦૮/૨૦૨૧ થી તા. ૨૧/૦૮/૨૦૨૧ સામાન્ય વરસાદની શક્યતાઓ છે.
- મેધ્દૂત એપ્લીકેશન:-** કૃષિ હવામાન સલાહ સેવા અને હવામાન આગાહી અંગેની જાણકારી મેળવવા માટે **ડાઉનલોડ** કરો મેધ્દૂત એપ્લીકેશન કે જે ખેડૂતોને તેમની સ્થાનિક ભાષામાં સ્થાન, પાક અને પશુધન વિશે હવામાન આધારિત કૃષિ સલાહ પ્રદાન કરશે.
- મેધ્દૂત એપ્લીકેશનને **ડાઉનલોડ** કરવા માટે નીચેની લિંક પર ક્લિક કરો.
એન્ડ્રોઇડ(મોબાઇલ)ઉપયોગકર્તાઃ
<https://play.google.com/store/apps/details?id=com.aas.meghdoot>
આઈ.ઓ.એસ. ઉપયોગકર્તાઃ-
<https://apps.apple.com/in/app/meghdoot/id1474048155>
- દામની એપ્લીકેશન:-** દામની એપ્લીકેશન દ્વારા આકાશીય વિજળીની અગાઉથી ચેતવણી આપવામાં આવે છે તેમજ આકાશીય વિજળીથી બચવા શું કરવું અને શું નહિ કરવું તેની વિગતવાર માહિતી પૂરી પાડવામાં આવે છે. આથી દામની એપ્લીકેશન ડાઉનલોડ કરવા માટે નીચેની લિંક પર ક્લિક કરો.
એન્ડ્રોઇડ(મોબાઇલ)ઉપયોગકર્તાઃ-
<https://play.google.com/store/apps/details?id=com.lightening.live.damini>
આઈ.ઓ.એસ. ઉપયોગકર્તાઃ-
<https://apps.apple.com/app/id1502385645>
- કોરોના (કોવિડ-૧૯) રોગચાળાની મહામારી ના લીધે, ખેડૂતમિત્રો એ ખેતર પર ખેતીની કામગીરી દરમિયાન ખેત સાધનો / ઓજારો અને ખેત મજુરો સાથે એકબીજાથી સામાજિક અંતર રાખવું અને તે માટે ની સાવધાની અને સલામતી રાખવી તેમજ અગમચેતી જાળવવી.
- ખેતરના સેઢા,પાળા અને વરસાદના પાણી નિકાલની ગટર ને ચોખ્માં કરી ને આગામી ચોમાસા માટે ખેતર ને તૈયાર કરવુ.
- ચોમાસું પાકોમાં આંતરખેડ કરવી અને હાથથી નિંદામણ કરવુ.
- વરાપ સમયે ખાતર, કીટનાશક અને ફૂગનાશક જેવી દવાનો છંટકાવ કરવો.

**NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 30
(23.07.2021 to 29.07.2021) over Agricultural region**

Gujarat



Agricultural vigour is moderate over isolated places of the state.

Source: <https://www.star.nesdis.noaa.gov>

પાકનું નામ	પાક અવસ્થા	રોગ / જીવાત / જાત	કૃષિ સલાહ
ચોમાસું મગાફળી	વૃક્ષ / ફૂલ / સુયા બેસવા	મુંડા, લશકરી ઈયળ, લીલી ઈયળ, ચુસીયા જીવાતો, વહેલા & મોડા પાનનાં ટપકાં	<ul style="list-style-type: none"> ❖ લશકરી ઈયળ અને લીલી ઈયળના નિયંત્રણ માટે કવીનાલફોસ ૨૫ ઇસી અથવા કલોરપાયરીફોસ ૨૦ ઇસી અથવા ફલુબેન્નીયામાઈડ ૨૦ ડબલ્યુ.જી. અથવા નોવાલ્યુરોન ૫.૨૫+ ઇન્ડોક્રાકાર્બ ૪.૫૦ એસ.સી. મુજબ છંટકાવ કરવો. ❖ ચુસીયા જીવાતો માટે લેમડા-સાયહેલોથ્રીન ૫ ઇ.સી. અથવા ઇમીડાકલોપ્રીડ ૧૭.૮ એસ.એલ. અથવા થાયામીથોકામ ૧૨.૬૦ + લેમડા-સાયહેલોથ્રીન ૮.૫૦ ઝેડ.સી. નો છંટકાવ કરવો. ❖ મગાફળીમાં ગેરૂ અને ટીક્કાના નિયંત્રણ માટે કુગનાશક દવા મેન્કોઓબ ૩૦ ગ્રામ દવા ૧૦ લીટર પાણીમાં ભેણવી વાવેતર બાદ ૩૫ દિવસે છંટકાવ કરવો. ❖ મગાફળી પાકને લોહ તત્ત્વની ઉણપના કારણે પાન પીળા દેખાય તો ૧૦૦ ગ્રામ હીરાકશી તથા ૧૦ ગ્રામ લીંબુના ફૂલને ૧૦ લીટર પાણીમાં ઓગાળીને છંટકાવ કરવો. ❖ મગાફળીનાં પાકમાં વરાપ થયે નિંદામણ અને આંતરખેડ કરવી. ❖ ઘેતરમાં પ્રકાશ પિંજર મુકવા.

			<ul style="list-style-type: none"> ❖ ટેબ્યુકોનાડોલ ૨૫.૮ ઇસી અથવા પ્રોપીકોનાડોલ અથવા ડાયફેનકોનાડોલ ૨૫ ટકા ઇસી ૧૦ મી.લી./૧૦ લી. પ્રમાણે ત્રણ છંટકાવ ૩૫, ૫૦ અને ૭૫ દિવસે કરવા અથવા સુડોમોનાસ ફલુરોસન્સ ૨૦ મી.લી./૧૦ લી. મુજબ છંટકાવ કરવો.
કપાસ	વૃદ્ધિ / ડાળી	પાનનો ટપકાંનો રોગ, ચુસીયા પ્રકારની જીવાતો, ખુણીયાં ટપકાં	<ul style="list-style-type: none"> ❖ પાનનો ટપકાંનો રોગ માટે કાર્બોનાઇમ ૫૦% વે. પા. ૫ ગ્રામ/૧૦ લી. મુજબ છંટકાવ કરવો. ❖ પાકની શરૂઆતની અવસ્થામાં કોઈ પણ પ્રકારની રાસાયણિક જંતુનાશક દવાનો છંટકાવ કરવો નહિ કારણકે બીજ માવજત દરમિયાન આપેલી જંતુનાશક દવા પાકને ૩૦-૪૫ દિવસ સુધી રક્ષણ આપે છે. ❖ પાક નાં શરૂઆતની અવસ્થામાં પાકની વૃદ્ધિ દરમિયાન જીવાતનાં કુદરતી દુશ્મન પૂરતા પ્રમાણમાં હોય છે. જે જીવાતને નિયંત્રણ માં રાખે છે. ❖ નુકશાનકારક રસાયણનો છંટકાવ કુદરતી દુશ્મનો નો નાશ કરે છે. ❖ નિયો નિકોટીનોઈડ વર્ગની જંતુનાશકનો ચુસીયા પ્રકારની જીવાત સામેનો વારેવાર ઉપયોગ ટાળવો જોઈએ. ❖ જો ૨૫% જીટલા છોડમાં ચુસીયા પ્રકારની જીવાતનું નુકશાન જોવા મળે તો ફ્લોનીકામીડ ૫૦ ડબ્લ્યુ. જી. ૪ ગ્રામ/ ૧૦ લીટર પાણીમાં લેળવી પાક પર છંટકાવ કરવો. ❖ વાવેલ પાકમાં ખાલા પુરવા અને છોડની પારવણીની કામગીરી હાથ ધરવી. ❖ કપાસનાં પાકમાં વરાપ થયે નિંદામણ અને આંતરખેડ કરવી. ❖ કપાસમાં ખુણીયાં ટપકાં નાં રોગ માટે સ્ટ્રેપ્ટોસાઈકલીન ૧ ગ્રામ/૧૦ લી.પાણીમાં ઓગાળી છંટકાવ કરવો.
સોયાબીન	વૃદ્ધિ / ફૂલ	લશકરી ઈયળ, ઘોડિયા ઈયળ	<ul style="list-style-type: none"> ❖ ગર્ડલ બીટલ, કાતરા અને લશકરી ઈયળથી નુકશાનીત છોડવાઓ, ઈંડાઓનો સમૂહ અને ઈયળનો નાશ કરવો. ❖ સોયાબીનનું વાવેતર કર્યા બાદ ૩૦ દિવસે રીલે આંતરપાક તરીકે બે હાર પછી એક હાર ઓરંડાનું વાવેતર કરવુ જેથી વધુ ઉત્પાદન મેળવી શકાય છે. ❖ સોયાબીનનાં પાકમાં વરાપ થયે નિંદામણ અને આંતરખેડ કરવી.

			<ul style="list-style-type: none"> ❖ સોયાબીનમાં લશકરી અને ધોડિયા ઈયળ માટે પ્રોફેનોફોસ ૫૦ ઇ.સી. અથવા કલોરાનદ્રાનીલીપ્રોલ ૧૮.૫ એસ.સી. અથવા એમામેકટીન બેન્જોએટ ૧.૬૦ ઇ.સી. અથવા ઇથીયોન ૫૦ ઇ.સી. અથવા ફલુબેન્ડીયામાઈડ ૩૮.૩૫ એસ.સી. અથવા ઇન્ડોકાકાર્બ ૧૫.૮૦ ઇ.સી. અથવા લેમડા-સાયહેલોથ્રીન ૪.૬૦ સી.એસ. અથવા સ્પીનેટોરમ ૧૧.૭૦ એસ.સી. અથવા નોવાલ્યુરોન ૫.૨૫+ ઇન્ડોકાકાર્બ ૪.૫૦ એસ.સી. અથવા થાયામીથોકામ ૧૨.૯૦ + લેમડા-સાયહેલોથ્રીન ૬.૫૦ ઎ડ.સી. નો છંટકાવ કરવો.
મગા	વૃદ્ધિ	પાનનાં ટપકાં	<ul style="list-style-type: none"> ❖ કંસિયા, સફેદ માખી અને તડતડીયા વગેરેના નિયંત્રણ માટે તુવેર, ગલગોટા અને દિવેલા જેવા પાકોનો આંતરપાક તરીકે ઉપયોગ કરવો. ❖ રક્ષક પાક તરીકે જીવાર અથવા મકાઈની ૪ હાર મગાની આસપાસ કરવી. ❖ નાઈટ્રોજનયુક્ત ખાતરનો અપ્રમાણસર ઉપયોગ કરવો નહિં. ❖ મગાનાં પાકમાં વરાપ થયે નિંદામણ અને આંતરખેડ કરવી. ❖ રોગની શરૂઆત થયે મેટીરામ પ૫% + પાયરેકલોસ્ટ્રોબીન પ૫% ડબલ્યુજી ૩૦ ગ્રામ/૧૦ લી. અથવા ટેબ્યુંકોનાઓલ ૨૫.૬ ટકા અથવા હેકાકોનાઓલ ૫ ટકા ૧૦ મિલી/૧૦ લિટર મુજબ છંટકાવ કરવો.
અડન	વૃદ્ધિ	પાનનાં ટપકાં	<ul style="list-style-type: none"> ❖ કંસિયા, સફેદ માખી અને તડતડીયા વગેરેના નિયંત્રણ માટે તુવેર, ગલગોટા અને દિવેલા જેવા પાકોનો આંતરપાક તરીકે ઉપયોગ કરવો. ❖ રક્ષક પાક તરીકે જીવાર અથવા મકાઈની ૪ હાર મગાની આસપાસ કરવી. ❖ નાઈટ્રોજનયુક્ત ખાતરનો અપ્રમાણસર ઉપયોગ કરવો નહિં. ❖ અડનનાં પાકમાં વરાપ થયે નિંદામણ અને આંતરખેડ કરવી. ❖ રોગની શરૂઆત થયે મેટીરામ પ૫% + પાયરેકલોસ્ટ્રોબીન પ૫% ડબલ્યુજી ૩૦ ગ્રામ/૧૦ લી. અથવા ટેબ્યુંકોનાઓલ ૨૫.૬ ટકા અથવા હેકાકોનાઓલ ૫ ટકા ૧૦ મિલી/૧૦ લિટર મુજબ છંટકાવ કરવો.
તલ	વૃદ્ધિ	તડતડીયા, સફેદ માખી	<ul style="list-style-type: none"> ❖ તલનાં પાકમાં વરાપ થયે નિંદામણ અને આંતરખેડ કરવી.

			<ul style="list-style-type: none"> ❖ તલમાં તડતડીયા અને સફેદ માખી માટે ઓકાઈડીમેટોન-મિથાઈલ રપ ઇ.સી.નો છંટકાવ કરવો.
તુવેર	ઉગાવો	-	<ul style="list-style-type: none"> ❖ લીલી ઈયળના નિયંત્રણ માટે ગલગોટાનો પિંજર પાક તરીકે ઉપયોગમાં કરી શકાય. ❖ ઈડાનો અને નાની ઈયળોને એકત્રિત કરી નાશ કરવો.
એરંડા	ઉગાવો	-	<ul style="list-style-type: none"> ❖ વાવણી લાયક વરસાદ થયે એરંડાનું વાવેતર કરવું.
શાકભાજી	વૃધ્ઘી/કુલ/ફળ	-	<ul style="list-style-type: none"> ❖ વેલાવાળા શાકભાજીના પાકમાં પાન કોરીયાનો ઉપદ્રવ ટાળવા માટે મુખ્ય પાકની ફરતે દિવેલા તથા ગલગોટાનું વાવેતર કરવાથી તેનું નિયંત્રણ કરી શકાય છે જો ઉપદ્રવ વધુ જોવા મળે તો ૧૦ મીલી મીથાઈલ ઓ ડેમેટોન (૨૫ ઇસી) અથવા ૧૦ મીલી ડાયમીથોએટ (૩૦ ઇસી) ૧૦ લીટર પાણીમાં લેળવી છંટકાવ કરવો ત્યાર બાદ પછીથી બીજો છંટકાવ ૧૫ દિવસ પછી કરવો. ❖ શાકભાજીના પાકમાં ચુસીયા પ્રકારની જીવાતના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦ મીલી અથવા લીબોળીનું મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦મીલી ૧૦ લી. પાણીમાં લેળવી છંટકાવ કરવો. જો વધુ ઉપદ્રવ હોય તો એસીટામ્રાપ્રીડ ૨૦ ટકા ૫ ગ્રામ/૧૦લી. અથવા દ્રાયઓફ્સેસ ૪૦ ટકા ૨૦ મીલી/૧૦ લી.પાણીમાં મીશ્ર કરી છંટકાવ કરવો. ❖ મરચીના પાકમાં થીઝસના નિયંત્રણ માટે ૨૦મીલી પ્રોફેનોઝોસ અને લિમ્બોળીનું તેલ ૫૦ મીલી ૧૦ લીટર પાણી મિશ્રણ કરી છાંટવી. ❖ શાકભાજીના પાકની રોપણી બાદ ચુસીયા પ્રકારની જીવાતના નિયંત્રણ માટે દાણાદાર ફોરેટ ૧૦ જી છોડ દીઠ ૨ ગ્રામ મુજબ આપીને પીયત આપવું. ❖ મરચીના પાકમાં વાઈરસના કારણે પીળા થઈ કુકડાઈ ગયેલ છોડ ઉપાડી જમીનમાં દાટી નાશ કરવો અને ૫ મીલી ઇમીડાક્લોપ્રાઇડ ૧૦ લીટર પાણીમાં લેળવી છંટકાવ કરવો. ❖ રીંગણ અને ટમેટાના ફળ અને અને ડોકાની ઈયળોનો ઉપદ્રવ જણાય તો નુકશાન પામેલ ફળ અને ડોકાને તોડીને જમીનમાં દાટી દેવા અને રીનાક્ષીપાયર દવા ઉમીલીદવા ૧૦ લીટર પાણીમાં લેળવી છંટકાવ કરવો.

			<ul style="list-style-type: none"> ❖ રીંગણ અને ટમેટાના પાકમા ફળમાખીના નિયંત્રણ માટે એસીટેમાપ્રોડ ૨૦ ટકા પગ્રામ દવા ૧૦ લીટર પાણીમાં લેળવી છંટકાવ કરવો. ❖ શાકભાજુ ને સમયસર ઉતારી ગ્રેડીંગ કરી બજારમાં મુકવા.
પશુપાલન	-	-	<ul style="list-style-type: none"> ❖ પશુઓને યોગ્ય આહાર આપવો & પીવા માટે ચોખ્યું પાણી આપવું અને પશુઓના રહેઠાણ ચોખ્યા રાખવા. ❖ માખી મચ્છરથી રક્ષણ માટે ફીનાઈલ છાટવું. ❖ પશુઓને યોગ્ય આહારમાં ઘનિજયુક્ત મિશ્રણ સાથે સંગ્રહિત ચારા આપો. ચારા ને સલામત સ્થળે સંગ્રહિત કરવો જોઈએ. ❖ ચારા માટે ચાફ્ કટર વાપરો. ❖ પશુઓમાં ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટાજ ૨ મીલી/ ૧ લીટર પાણીમાં નાખીને છાટવી. ❖ દુધાળા પશુઓના આઉને ઝીંક ઓક્સાઇડ અથલા બોરીક પાવડરથી બરાબર સાફ્ કરવા.



ગ્રામીણ કૃષિ મૌસમ સેવા
કૃષિ હવામાન સેલ, કૃષિ વિજ્ઞાન વિભાગ
જુનાગઢ કૃષિ યુનિવર્સિટી
જુનાગઢ - ૩૬૨૦૦૧

ઈમેઇલ:- amfujnd@yahoo.com ફોન નં. ૦૨૮૫-૨૬૭૦૧૩૮



હવામાન આગાહી આધારીત દક્ષિણ સૌરાષ્ટ્ર એત આબોહવાકીય વિસ્તારનાં પોરબંદર જીલ્લા માટે કૃષિ સલાહ બુલેટીન
(કૃષિ હવામાન સેલ, જુનાગઢ કૃષિ યુનિવર્સિટી અને ભારત હવામાન વિભાગ દ્વારા સંપદ્ધ)

(#આજાદી નો અમૃત મહોત્સવ#)

વાર:- મંગળવાર

તારીખ:- ૧૦/૦૮/૨૦૨૧

હવામાન આગાહી આધારીત બુલેટીન: ૦૩૮ (૨૦૨૧-૨૨)

પાછલા પાંચ દિવસનું હવામાન (તારીખ:- ૦૫ થી ૦૯ ઓગસ્ટ, ૨૦૨૧)					હવામાન પરિબળો		હવામાન આગાહી આગામી પાંચ દિવસ માટે (તારીખ:- ૧૧ થી ૧૫ ઓગસ્ટ, ૨૦૨૧) માન્ય ૧૫ ઓગસ્ટ, ૨૦૨૧. સવારના ૮:૩૦ વાગ્યાં સુધી				
૦૫/૦૮	૦૬/૦૮	૦૭/૦૮	૦૮/૦૮	૦૯/૦૮	તારીખ	૧૧/૦૮	૧૨/૦૮	૧૩/૦૮	૧૪/૦૮	૧૫/૦૮	
૦	૦	૦	૨.૦	૦	વરસાદ (મિમી)	૨	૨	૩	૩	૩	
૩૧.૦	૩૧.૦	૩૦.૮	૩૨.૦	૩૧.૬	મહત્વમાન તાપમાન (°સે.ગ્રે)	૩૨	૩૧	૩૧	૩૧	૩૦	
૨૬.૫	૨૬.૮	૨૬.૫	૨૬.૨	૨૬.૫	લઘુત્વમાન તાપમાન (°સે.ગ્રે)	૨૭	૨૭	૨૭	૨૬	૨૬	

મોટેલાગે વાદળાયું	વાદળની સ્થિતિ	વાદળાયું	મોટેલાગે વાદળાયું	વાદળાયું	વાદળાયું	વાદળાયું				
૮૩	૮૩	૮૦	૮૭	૭૬	સવારના લેજના ટકા	૭૮	૮૦	૮૦	૮૦	૮૧
૮૨	૮૨	૭૭	૭૬	૭૪	અપોરબાદ ના લેજના ટકા	૬૬	૬૬	૭૩	૭૩	૭૨
૧૮.૫	૨૨.૨	૧૪.૮	૧૪.૮	૧૧.૧	પવનની ગતિ (કિમી/કલાક)	૨૭.૧	૨૮.૦	૨૭.૦	૨૮.૩	૩૦.૨
૨૫૦	૨૫૦	૨૫૦	૨૩૦	૨૫૦	પવનની દિશા	૨૪૮	૨૪૮	૨૪૮	૨૪૮	૨૪૮

સામાન્ય સલાહ

ભારત સરકારની હવામાન વિભાગ તરફથી મળેલ હવામાન આગાહીની માહિતી અનુસાર અતેના ગ્રામીણ કૃષિ મૌસમ સેવા યોજના, જૂનાગઢ કૃષિ યુનિવર્સિટી, જૂનાગઢ તરફથી જણાવવામા આવે છે કે દક્ષિણ સૌરાષ્ટ્ર પેઠ આબોહવાકીય વિસ્તારના પોરબંદર જીલ્લામાં આગામી તારીખ ૧૧/૦૮/૨૦૨૧ થી ૧૫/૦૮/૨૦૨૧ દરમયાન આકાશ મુખ્યત્વે આગામી પાંચ દિવસોમાં મોટેલાગે વાદળાયું થી વાદળાયું રહેશે, જીલ્લામાં આગામી પાંચ દિવસોમાં છુટા છવાયાં ભાગોમાં હળવા / અપાં વરસાદ ની શક્યતાઓ છે. પવનની ઝડપ ૨૭.૦ થી ૩૦.૨ કિમી/કલાક રહેવાની શક્યતા રહેશે. આ સમયગાળામાં મહત્વમાન તાપમાન ૩૦ થી ૩૨ ડિગ્રી સેન્ટીગ્રેડ રહેશે તેમજ લઘુત્તમ તાપમાન ૨૬ થી ૨૭ ડિગ્રી સેન્ટીગ્રેડ રહેવાની સંભાવના છે.

+ માન્ય વરસાદની સૂચી મુજબ પોરબંદર જીલ્લામાં ૦૮ જુલાઈ થી ૦૪ ઓગસ્ટ, ૨૦૨૧ ના સમયગાળા દરમયાન હળવી લેજ ની સ્થિતિ જોવા મળેલ છે.

+ સાપ્તાહિક વરસાદ આગાહીની સૂચી મુજબ દક્ષિણ સૌરાષ્ટ્રનાં વિસ્તારમાં તા. ૧૫/૦૮/૨૦૨૧ થી તા. ૨૧/૦૮/૨૦૨૧ સામાન્ય વરસાદની શક્યતાઓ છે.

+ મેધ્દૂત એપ્લીકેશન:- કૃષિ હવામાન સલાહ સેવા અને હવામાન આગાહી અંગેની જાણકારી મેળવવા માટે ડાઉનલોડ કરો મેધ્દૂત એપ્લીકેશન કે જે ખેડૂતોને તેમની સ્થાનિક ભાષામાં સ્થાન, પાક અને પશુધન વિશે હવામાન આધારિત કૃષિ સલાહ પ્રદાન કરશે.

+ મેધ્દૂત એપ્લીકેશનને ડાઉનલોડ કરવા માટે નીચેની લિંક પર ક્લિક કરો.
એન્ડ્રોઇડ(મોબાઇલ)ઉપયોગકર્તા:

<https://play.google.com/store/apps/details?id=com.aas.meghdoot>

આઈ.ઓ.એસ. ઉપયોગકર્તા:-

<https://apps.apple.com/in/app/meghdoot/id1474048155>

+ દામિની એપ્લીકેશન:- દામિની એપ્લીકેશન દ્વારા આકાશીય વિજળીની અગાઉથી ચેતવણી આપવામાં આવે છે તેમજ આકાશીય વિજળીથી બચવા શું કરવું અને શું નહિ કરવું તેની વિગતવાર માહિતી પૂરી પાડવામાં આવે છે. આથી દામિની એપ્લીકેશન ડાઉનલોડ કરવા માટે નીચેની લિંક પર ક્લિક કરો.

એન્ડ્રોઇડ(મોબાઇલ)ઉપયોગકર્તા:-


<https://play.google.com/store/apps/details?id=com.lightening.live.damini>

આઈ.ଓ.એસ. ઉપયોગકર્તા:-

<https://apps.apple.com/app/id1502385645>

- ❖ કોરોના (કોવીડ-૧૯) રોગચાળાની મહામારી ના લીધે, જેઠૂતમિત્રો એ ખેતર પર ખેતીની કામગીરી દરમિયાન ખેત સાધનો / ઓજારો અને ખેત મજૂરો સાથે એકબીજાથી સામાજુક અંતર રાખવું અને તે માટે ની સાવધાની અને સલામતી રાખવી તેમજ અગમચેતી જાળવવી.
- ❖ ખેતરના સેઢા, પાળા અને વરસાદનાં પાણી નિકાલની ગટર ને ચોમાસું ઋતુમાં ચોખ્ખાં કરવા.
- ❖ ચોમાસું પાકોમાં આંતરખેડ કરવી અને હાથથી નિંદામણ કરવું.
- ❖ વરાપ સમયે ખાતર, કીટનાશક અને ફૂગનાશક જીવી દવાનો છંટકાવ કરવો.

NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 30
(23.07.2021 to 29.07.2021) over Agricultural region



Agricultural vigour is moderate over isolated places of the state.

Source: <https://www.star.nesdis.noaa.gov>

પાકનું નામ	પાક અવસ્થા	રોગ / જીવાત / જાત	ફૂંઝિ સલાહ
ચોમાસું મગફળી	વૃદ્ધિ / ફૂલ / સુયા બેસવા	મુંડા, લશકરી ઈયળ, લીલી ઈયળ, ચુસીયા જીવાતો, વહેલા & મોડા પાનનાં ટપકાં	<ul style="list-style-type: none"> ❖ લશકરી ઈયળ અને લીલી ઈયળના નિયંત્રણ માટે કવીનાલફેસ ૨૫ ઇસી અથવા કલોરપાયરીફેસ ૨૦ ઇસી અથવા ફલુબેન્ડીયામાઈડ ૨૦ ડબલ્યુ.જી. અથવા નોવાલ્યુરોન ૫.૨૫+ ઇન્ડોક્રાકાર્બ ૪.૫૦ એસ.સી. મુજબ છંટકાવ કરવો. ❖ ચુસીયા જીવાતો માટે લેમડા-સાયહેલોથ્રીન ૫ ઇ.સી. અથવા ઇમીડાકલોપ્રીડ ૧૭.૮ એસ.એલ. અથવા થાયામીથોકાઝામ

			<p>૧૨.૬૦ + લેમડા-સાચહેલોથીન ૮.૫૦ ડેડ.સી. નો છંટકાવ કરવો.</p> <ul style="list-style-type: none"> ❖ મગફળીમાં ગેરુ અને ટીક્કાના નિયંત્રણ માટે કુગનાશક દવા મેન્જોરેબ ૩૦ ગ્રામ દવા ૧૦ લીટર પાણીમાં ભેળવી વાવેતર બાદ ૩૫ દિવસે છંટકાવ કરવો. ❖ મગફળી પાકને લોહ તત્વની ઉણપના કારણે પાન પીળા દેખાય તો ૧૦૦ ગ્રામ હીરાકશી તથા ૧૦ ગ્રામ લીંબુના ફૂલને ૧૦ લીટર પાણીમાં ઓગાળીને છંટકાવ કરવો. ❖ મગફળીનાં પાકમાં વરાપ થયે નિંદામણ અને આંતરખેડ કરવી. ❖ ઘેતરમાં પ્રકાશ પિંજર મુકવા. ❖ ટેબ્યુકોનાઓલ ૨૫.૬ ઇસી અથવા પ્રોપીકોનાઓલ અથવા ડાયફેનકોનાઓલ ૨૫ ટકા ઇસી ૧૦ મી.લી./૧૦ લી. પ્રમાણે ત્રણ છંટકાવ ૩૫, ૫૦ અને ૭૫ દિવસે કરવા અથવા સુડોમોનાસ ફલુરોસન્સ ૨૦ મી.લી./૧૦ લી. મુજબ છંટકાવ કરવો.
કપાસ	વૃદ્ધિ / ડાળી	પાનનો ટપકાનો રોગ, ચુસીયા પ્રકારની જીવાતો, ખુણીયાં ટપકાનો	<ul style="list-style-type: none"> ❖ પાનનો ટપકાનો રોગ માટે કાર્બેન્ડાઈમ ૫૦% વે. પા. ૫ ગ્રામ/૧૦ લી. મુજબ છંટકાવ કરવો. ❖ પાકની શરૂઆતની અવસ્થામાં કોઈ પણ પ્રકારની રાસાયણિક જંતુનાશક દવાનો છંટકાવ કરવો નહિ કારણકે બીજ માવજત દરમિયાન આપેલી જંતુનાશક દવા પાકને ૩૦-૪૫ દિવસ સુધી રક્ષણ આપે છે. ❖ પાક નાં શરૂઆતની અવસ્થામાં પાકની વૃદ્ધિ દરમિયાન જીવાતનાં કુદરતી દુશ્મન પૂરતા પ્રમાણમાં હોય છે. જે જીવાતને નિયંત્રણ માં રાખે છે. ❖ નુકશાનકારક રસાયણનો છંટકાવ કુદરતી દુશ્મનો નો નાશ કરે છે. ❖ નિયો નિકોટીનોઈડ વર્ગની જંતુનાશકનો ચુસીયા પ્રકારની જીવાત સામેનો વારંવાર ઉપયોગ ટાળવો જોઈએ. ❖ જો ૨૫% જેટલા છોડમાં ચુસીયા પ્રકારની જીવાતનું નુકશાન જોવા મળે તો ફલોનીકામીડ ૫૦ ડબ્લ્યુ. જી. ૪ ગ્રામ/ ૧૦ લીટર પાણીમાં ભેળવી પાક પર છંટકાવ કરવો. ❖ વાવેલ પાકમાં ખાલા પુરવા અને છોડની પારવણીની કામગીરી હાથ ધરવી.

			<ul style="list-style-type: none"> ❖ કપાસનાં પાકમાં વરાપ થયે નિંદામણ અને આંતરખેડ કરવી. ❖ કપાસમાં ખુણીયાં ટપકાં નાં રોગ માટે સ્ટ્રેપ્ટોસાઇકલીન ૧ ગ્રામ/૧૦ લી.પાણીમાં ઓગાળી છંટકાવ કરવો.
સોયાબીન	વૃદ્ધિ / ફૂલ	લશકરી ઈયળ, ઘોડિયા ઈયળ	<ul style="list-style-type: none"> ❖ ગર્ડલ બીટલ, કાતરા અને લશકરી ઈયળથી નુકાશાનીત છોડવાઓ, ઈંડાઓનો સમૂહ અને ઈયળનો નાશ કરવો. ❖ સોયાબીનનું વાવેતર કર્યા બાદ ૩૦ દિવસે રીલે આંતરપાક તરીકે બે હાર પછી એક હાર એંરંડાનું વાવેતર કરવું જેથી વધુ ઉત્પાદન મેળવી શકાય છે. ❖ સોયાબીનનાં પાકમાં વરાપ થયે નિંદામણ અને આંતરખેડ કરવી. ❖ સોયાબીનમાં લશકરી અને ઘોડિયા ઈયળ માટે પ્રોફેનોફોસ ૫૦ ઇ.સી. અથવા કલોરાનદ્રાનીલીપ્રોલ ૧૮.૫ એસ.સી. અથવા એમામેકટીન બેન્જોએટ ૧.૬૦ ઇ.સી. અથવા ઇથીયોન ૫૦ ઇ.સી. અથવા ફલુબેન્ઝીયામાઈડ ૩૮.૩૫ એસ.સી. અથવા ઇન્ડોકાકાર્બ ૧૫.૮૦ ઇ.સી. અથવા લેમડા-સાયહેલોથ્રીન ૪.૬૦ સી.એસ. અથવા સ્પીનેટોરમ ૧૧.૭૦ એસ.સી. અથવા નોવાલ્યુરોન ૫.૨૫+ ઇન્ડોકાકાર્બ ૪.૫૦ એસ.સી. અથવા થાયામીથોકાકામ ૧૨.૬૦ + લેમડા-સાયહેલોથ્રીન ૮.૫૦ એડ.સી. નો છંટકાવ કરવો.
મગ	વૃદ્ધિ	પાનનાં ટપકાં	<ul style="list-style-type: none"> ❖ કાંસિયા, સફેદ માખી અને તડતડીયા વગેરેના નિયંત્રણ માટે તુવેર, ગલગોટા અને દિવેલા જીવા પાકોનો આંતરપાક તરીકે ઉપયોગ કરવો. ❖ રક્ષક પાક તરીકે જુવાર અથવા મકાઈની ૪ હાર મગની આસપાસ કરવી. ❖ નાઈટ્રોજનયુક્ત ખાતરનો અપ્રમાણસર ઉપયોગ કરવો નહિં. ❖ મગનાં પાકમાં વરાપ થયે નિંદામણ અને આંતરખેડ કરવી. ❖ રોગની શરૂઆત થયે મેટીરામ ૫૫% + પાયરેકલોસ્ટ્રોબીન ૫% ડબલ્યુજી ૩૦ ગ્રામ/૧૦ લી. અથવા ટેબ્યુનોનાઝોલ ૨૫.૬ ટકા અથવા હેકાકોનાઝોલ ૫ ટકા ૧૦ મિલી/૧૦ લિટર મુજબ છંટકાવ કરવો.

અડિ	વૃદ્ધિ	પાનનાં ટપકાં	<ul style="list-style-type: none"> ❖ કંસિયા, સફેદ માખી અને તડતડીયા વગેરેના નિયંત્રણ માટે તુવેર, ગલગોટા અને દિવેલા જેવા પાકોનો આંતરપાક તરીકે ઉપયોગ કરવો. ❖ રક્ષક પાક તરીકે જીવાર અથવા મકાઈની ૪ હાર મગની આસપાસ કરવી. ❖ નાઈટ્રોજનયુક્ત ખાતરનો અપ્રમાણસર ઉપયોગ કરવો નહિ. ❖ અડદનાં પાકમાં વરાપ થયે નિંદામણ અને આંતરખેડ કરવી. ❖ રોગની શરૂઆત થયે મેટીરામ પ્રપ% + પાયરેકલોસ્ટ્રોબીન પ% ડબલ્યુજી ૩૦ ગ્રામ/૧૦ લી. અથવા ટેબ્યુંકોનાઝોલ ૨૫.૬ ટકા અથવા હેકઝોનાઝોલ ૫ ટકા ૧૦ મિલી/૧૦ લિટર મુજબ છંટકાવ કરવો.
તલ	વૃદ્ધિ	તડતડીયા, સફેદ માખી	<ul style="list-style-type: none"> ❖ તલનાં પાકમાં વરાપ થયે નિંદામણ અને આંતરખેડ કરવી. ❖ તલમાં તલમાં તડતડીયા અને સફેદ માખી માટે ઓક્ઝિડિમેટોન-મિથાઈલ ૨૫ છ.સી.નો છંટકાવ કરવો. માટે ઓક્ઝિડિમેટોન-મિથાઈલ ૨૫ છ.સી.નો છંટકાવ કરવો.
તુવેર	ઉગાવો	-	<ul style="list-style-type: none"> ❖ લીલી ઈયળના નિયંત્રણ માટે ગલગોટાનો પિંજર પાક તરીકે ઉપયોગમાં કરી શકાય. ❖ ઈંડાનો અને નાની ઈયળોને એકત્રિત કરી નાશ કરવો.
એરંડા	ઉગાવો	-	<ul style="list-style-type: none"> ❖ વાવણી લાયક વરસાદ થયે એરંડાનું વાવેતર કરવુ.
શાકભાજી	વૃદ્ધી/કુલ/ ફળ	-	<ul style="list-style-type: none"> ❖ વેલાવાળા શાકભાજીના પાકમાં પાન કોરીયાનો ઉપદ્રવ ટાળવા માટે મુખ્ય પાકની ફરતે દિવેલા તથા ગલગોટાનું વાવેતર કરવાથી તેનું નિયંત્રણ કરી શકાય છે જો ઉપદ્રવ વધુ જોવા મળે તો ૧૦ મીલી મીથાઈલ ઓ ડેમેટોન (૨૫ ઈસી) અથવા ૧૦ મીલી ડાયમીથોએટ (૩૦ ઈસી) ૧૦ લિટર પાણીમાં લેળવી છંટકાવ કરવો ત્યાર બાદ પછીથી બીજો છંટકાવ ૧૫ દિવસ પછી કરવો. ❖ શાકભાજીના પાકમાં ચુસીયા પ્રકારની જીવાતના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦ મીલી અથવા લીબોળીનું મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦મીલી ૧૦ લી. પાણીમાં લેળવી છંટકાવ કરવો. જો વધુ ઉપદ્રવ હોય તો એસીટામ્રાગ્રીડ ૨૦ ટકા ૫ ગ્રામ/૧૦લી. અથવા દ્રાવણઝોસ ૪૦ ટકા ૨૦ મીલી/૧૦ લી.પાણીમાં મીશ્ર કરી છંટકાવ કરવો.

		<ul style="list-style-type: none"> ❖ મરચીના પાકમાં થીપ્સના નિયંત્રણ માટે ૨૦મીલી પ્રોફેનોફોસ અને લિપ્બોળીનું તેલ ૫૦ મીલી ૧૦ લીટર પાણી મિશ્રણ કરી છાંટવી. ❖ શાકભાજુના પાકની રોપણી બાદ ચુસીયા પ્રકારની જીવાતના નિયંત્રણ માટે દાણાદાર ફોરેટ ૧૦ જી છોડ દીઠ ૨ ગ્રામ મુજબ આપીને પીયત આપવું. ❖ મરચીના પાકમાં વાઈરસના કારણે પીળા થઈ ફુકડાઈ ગયેલ છોડ ઉપાડી જમીનમાં દાટી નાશ કરવો અને ૫ મીલી ઈમીડાક્લોપ્રાઇડ ૧૦ લીટર પાણીમાં બેળવી છંટકાવ કરવો. ❖ રીંગણ અને ટમેટાના ફળ અને અને ડોકાની ઈયણોનો ઉપદ્રવ જણાય તો નુકશાન પામેલ ફળ અને ડોકાને તોડીને જમીનમાં દાટી દેવા અને રીનાક્ષીપાયર દવા ૩મીલીદવા ૧૦ લીટર પાણીમાં બેળવી છંટકાવ કરવો. ❖ રીંગણ અને ટમેટાના પાકમાં ફળમાખીના નિયંત્રણ માટે એસીટેમાપ્રીડ ૨૦ ટકા પગ્રામ દવા ૧૦ લીટર પાણીમાં બેળવી છંટકાવ કરવો. ❖ શાકભાજુ ને સમયસર ઉતારી ગ્રેડિંગ કરી બજારમાં મુકવા.
પશુપાલન	-	<ul style="list-style-type: none"> ❖ પશુઓને યોગ્ય આહાર આપવો & પીવા માટે યોગ્યનું પાણી આપવું અને પશુઓના રહેઠાણ ચોખ્ખા રાખવા. ❖ માખી મચ્છરથી રક્ષણ માટે ફીનાઈલ છાટવું. ❖ પશુઓને યોગ્ય આહારમાં ખનિજ્યુક્ત મિશ્રણ સાથે સંગ્રહિત ચારા આપો. ચારા ને સલામત સ્થળે સંગ્રહિત કરવો જોઈએ. ❖ ચારા માટે ચાફ કટર વાપરો. ❖ પશુઓમાં ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટાજ ૨ મીલી/ ૧ લીટર પાણીમાં નાખીને છાટવી. ❖ દુધાળા પશુઓના આઉને ઝીંક ઓક્સાઈડ અથલા બોરીક પાવડરથી બરાબર સાફ કરવા.



ગ્રામીણ કૃષિ મૌસમ સેવા
કૃષિ હવામાન સેલ, કૃષિ વિજ્ઞાન વિભાગ
જુનાગઢ કૃષિ યુનિવર્સિટી
જુનાગઢ - ૩૬૨૦૦૧

ઈમેઇલ:- amfujind@yahoo.com ફોન નં. ૦૨૮૫-૨૬૭૦૧૩૬



હવામાન આગાહી આધારીત દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારનાં ગીર સોમનાથ જીલ્લા માટે કૃષિ સલાહ બુલેટીન

(કૃષિ હવામાન સેલ, જુનાગઢ કૃષિ યુનિવર્સિટી અને ભારત હવામાન વિભાગ દ્વારા સંપદ)

(#આગાહી નો અમૃત મહોત્સવ#)

વાર:- મંગળવાર

તારીખ:- ૧૦/૦૮/૨૦૨૧

હવામાન આગાહી આધારીત બુલેટીન: ૦૩૮ (૨૦૨૧-૨૨)

પાછલા પાંચ દિવસનું હવામાન (તારીખ:- ૦૫ થી ૦૯ જુલાઈ, ૨૦૨૧)					હવામાન પરિબળો		હવામાન આગાહી આગામી પાંચ દિવસ માટે (તારીખ:- ૧૧ થી ૧૫ ઓગસ્ટ, ૨૦૨૧)				
૦૫/૦૮	૦૬/૦૮	૦૭/૦૮	૦૮/૦૮	૦૯/૦૮	તારીખ	૧૧/૦૮	૧૨/૦૮	૧૩/૦૮	૧૪/૦૮	૧૫/૦૮	
૦	૦	૦	૦	૦	વરસાદ (મિમી)	૨	૩	૩	૩	૩	
૩૦.૧	૩૦.૮	૩૦.૯	૩૧.૫	૩૨.૬	મહત્વમાન તાપમાન (°સે.ગ્રે)	૩૪	૩૪	૩૩	૩૩	૩૪	
૨૬.૬	૨૬.૮	૨૬.૬	૨૬.૬	૨૬.૮	લઘુત્વમાન (°સે.ગ્રે)	૨૫	૨૫	૨૫	૨૬	૨૬	
મોટેલાગે વાદળાયું	મોટેલાગે વાદળાયું	મોટેલાગે વાદળાયું	મોટેલાગે વાદળાયું	મોટેલાગે વાદળાયું	વાદળાની સ્થિતિ	મોટેલાગે વાદળાયું	વાદળાયું	વાદળાયું	વાદળાયું	વાદળાયું	
૮૬	૮૬	૮૬	૮૬	૮૨	સવારના ભેજના ટકા	૮૨	૮૪	૮૪	૮૪	૮૪	
૮૩	૮૨	૮૩	૮૨	૭૪	બપોરબાદ ના ભેજના ટકા	૬૫	૬૮	૭૧	૭૦	૭૧	
૧૧.૧	૬.૩	૫.૬	-	૩.૭	પવનની ગતિ (કિમી/કલાક)	૨૨.૫	૨૪.૧	૨૩.૫	૨૪.૬	૨૬.૩	
૨૫૦	૨૫૦	૨૫૦	-	૮૮	પવનની દિશા	૨૪૮	૨૪૮	૨૪૮	૨૪૮	૨૪૮	

સામાન્ય સલાહ

ભારત સરકારશીના હવામાન વિભાગ તરફથી મળેલ હવામાન આગાહીની માહીતી અનુસાર અત્રેના ગ્રામીણ કૃષિ મૌસમ સેવા યોજના, જુનાગઢ કૃષિ યુનિવર્સિટી, જુનાગઢ તરફથી જણાવવામા આવે છે કે દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારનાં ગીર સોમનાથ જીલ્લામાં આગામી તારીખ ૧૧/૦૮/૨૦૨૧ થી ૧૫/૦૮/૨૦૨૧ દરમ્યાન આકાશ મુખ્યત્વે આગામી પાંચ દિવસોમાં મોટેલાગે વાદળાયું થી વાદળાયું રહેશે, જીલ્લામાં આગામી પાંચ દિવસોમાં છુટા છવાયા ભાગોમાં હળવા / અપટાં વરસાદ ની શક્યતાઓ છે. પવનની ઝડપ ૨૨.૫ થી ૨૬.૩ કિમી/કલાક રહેવાની શક્યતા રહેશે. આ સમયગાળામાં મહત્વમાન તાપમાન ૩૩ થી ૩૪ ડિગ્રી સેન્ટીગ્રેડ રહેશે તેમજ લઘુત્વમાન રૂપ થી ૨૬ ડિગ્રી સેન્ટીગ્રેડ રહેવાની સંભાવના છે.

- માન્ય વરસાદની સૂચી મુજબ ગીર સોમનાથ જુલામાં ૦૮ જુલાઈ થી ૦૪ ઓગસ્ટ, ૨૦૨૧ ના સમયગાળા દરમિયાન હળવી ભેજ સ્થિતિ જોવા મળેલ છે.
- સાપ્તાહિક વરસાદ આગાહીની સૂચી મુજબ દક્ષિણ સૌરાષ્ટ્રનાં વિસ્તારમાં તા. ૧૫/૦૮/૨૦૨૧ થી તા. ૨૧/૦૮/૨૦૨૧ સામાન્ય વરસાદની શક્યતાઓ છે.

મેઘદૂત એપ્લીકેશન:- કૃષિ હવામાન સલાહ સેવા અને હવામાન આગાહી અંગેની જાણકારી મેળવવા માટે ડાઉનલોડ કરો મેઘદૂત એપ્લીકેશન કે જે ખેડૂતોને તેમની સ્થાનિક ભાષામાં સ્થાન, પાક અને પશુધન વિશે હવામાન આધારિત કૃષિ સલાહ પ્રદાન કરશે.

- મેઘદૂત એપ્લીકેશનને ડાઉનલોડ કરવા માટે નીચેની લિંક પર ક્લિક કરો.
એન્ડ્રોઇડ(મોબાઇલ)ઉપયોગકર્તાઃ

<https://play.google.com/store/apps/details?id=com.aas.meghdoot>

આઈ.ઓ.એસ. ઉપયોગકર્તાઃ-

<https://apps.apple.com/in/app/meghdoot/id1474048155>

- દામિની એપ્લીકેશનઃ- દામિની એપ્લીકેશન દ્વારા આકાશીય વિજળીની અગાઉથી ચેતવણી આપવામાં આવે છે તેમજ આકાશીય વિજળીથી બચવા શું કરવું અને શું નહિ કરવું તેની વિગતવાર માહિતી પૂરી પાડવામાં આવે છે. આથી દામિની એપ્લીકેશન ડાઉનલોડ કરવા માટે નીચેની લિંક પર ક્લિક કરો.

એન્ડ્રોઇડ(મોબાઇલ)ઉપયોગકર્તાઃ-

<https://play.google.com/store/apps/details?id=com.lightening.live.damini>


આઈ.ઓ.એસ. ઉપયોગકર્તાઃ-

<https://apps.apple.com/app/id1502385645>

- કોરોના (કોવીડ-૧૯) રોગચાળાની મહામારી ના લીધે, ખેડૂતમિત્રો એ ખેતર પર ખેતીની કામગીરી દરમિયાન ખેત સાધનો / ઓજારો અને ખેત મજૂરો સાથે એકબીજાથી સામાજીક અંતર રાખવું અને તે માટે ની સાવધાની અને સલામતી રાખવી તેમજ અગમચેતી જળવવી.
- ખેતરના સેઢા, પાળા અને વરસાદનાં પાણી નિકાલની ગટર ને ચોમાસું ઋતુમાં ચોખ્માં કરવા.
- ચોમાસું પાકોમાં આંતરખેડ કરવી અને હાથથી નિંદામણ કરવુ.
- વરાપ સમયે ખાતર, કીટનાશક અને કૂગનાશક જેવી દવાનો છંટકાવ કરવો.

**NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 30
(23.07.2021 to 29.07.2021) over Agricultural region**

Gujarat



Agricultural vigour is moderate over isolated places of the state.

Source: <https://www.star.nesdis.noaa.gov>

પાકનું નામ	પાક અવસ્થા	રોગ / જીવાત / જાત	કૃષિ સલાહ
ચોમાસું મગફળી	વૃદ્ધિ / ફૂલ / સુયા બેસવા	મુંડા, લશકરી ઈયળ, લીલી ઈયળ, ચુસીયા જીવાતો, વહેલા & મોડા પાનનાં ટપકાં	<ul style="list-style-type: none"> ❖ લશકરી ઈયળ અને લીલી ઈયળના નિયંત્રણ માટે કવીનાલફોસ ૨૫ ઇસી અથવા કલોરપાયરીફોસ ૨૦ ઇસી અથવા ફલુબેનીયામાઇડ ૨૦ ડબલ્યુ.જી. અથવા નોવાલ્યુરોન ૫.૨૫+ ઇન્ડોકાકાર્બ ૪.૫૦ એસ.સી. મુજબ છંટકાવ કરવો. ❖ ચુસીયા જીવાતો માટે લેમડા-સાચહેલોથીન ૫ એસ.સી. અથવા ઇમીડાકલોપ્રીડ ૧૭.૮ એસ.એલ. અથવા થાયામીથોકાઝમ ૧૨.૬૦ + લેમડા-સાચહેલોથીન ૮.૫૦ એડ.સી. નો છંટકાવ કરવો. ❖ મગફળીમાં ગેરુ અને ટીક્કાના નિયંત્રણ માટે કુગનાશક દવા મેન્કોઝેબ ૩૦ ગ્રામ દવા ૧૦ લીટર પાણીમાં ભેળવી વાવેતર બાદ ૩૫ દિવસે છંટકાવ કરવો. ❖ મગફળી પાકને લોહ તત્વની ઉણપના કારણે પાન પીળા દેખાય તો ૧૦૦ ગ્રામ હીરાકશી તથા ૧૦ ગ્રામ લીંબુના ફૂલને ૧૦ લીટર પાણીમાં ઓગાળીને છંટકાવ કરવો. ❖ મગફળીનાં પાકમાં વરાપ થયે નિંદામણ અને આંતરમેડ કરવી. ❖ ઘેતરમાં પ્રકાશ પિંજર મુકવા.

			<ul style="list-style-type: none"> ટેબ્યુકોનાડોલ ૨૫.૮ ઇસી અથવા પ્રોપીકોનાડોલ અથવા ડાયફેનકોનાડોલ ૨૫ ટકા ઇસી ૧૦ મી.લી./૧૦ લી. પ્રમાણે ત્રણ છંટકાવ ઉપલ ૫૦ અને ૭૫ દિવસે કરવા અથવા સુડોમોનાસ ફલુરોસન્સ ૨૦ મી.લી./૧૦ લી. મુજબ છંટકાવ કરવો.
કપાસ	વૃદ્ધિ / ડાળી	પાનનો ટપકાંનો રોગ, ચુસીયા પ્રકારની જીવાતો, ખુણીયાં ટપકાં	<ul style="list-style-type: none"> પાનનો ટપકાંનો રોગ માટે કાર્બેન્ડાઈમ ૫૦% વે. પા. ૫ ગ્રામ/૧૦ લી. મુજબ છંટકાવ કરવો. પાકની શરૂઆતની અવસ્થામાં છોઈ પણ પ્રકારની રાસાયણિક જંતુનાશક દવાનો છંટકાવ કરવો નહિ કારણકે બીજ માવજત દરમિયાન આપેલી જંતુનાશક દવા પાકને ૩૦-૪૫ દિવસ સુધી રક્ષણ આપે છે. પાક નાં શરૂઆતની અવસ્થામાં પાકની વૃદ્ધિ દરમિયાન જીવાતનાં કુદરતી દુશ્મન પૂરતા પ્રમાણમાં હોય છે. જે જીવાતને નિયંત્રણ માં રાખે છે. નુકશાનકારક રસાયણનો છંટકાવ કુદરતી દુશ્મનો નો નાશ કરે છે. નિયો નિકોટીનોઈડ વર્ગની જંતુનાશકનો ચુસીયા પ્રકારની જીવાત સામેનો વારેવાર ઉપયોગ ટાળવો જોઈએ. જો ૨૫% જીટલા છોડમાં ચુસીયા પ્રકારની જીવાતનું નુકશાન જોવા મળે તો ફ્લોનીકામીડ ૫૦ ડબ્લ્યુ. જી. ૪ ગ્રામ/ ૧૦ લીટર પાણીમાં લેળવી પાક પર છંટકાવ કરવો. વાવેલ પાકમાં ખાલા પુરવા અને છોડની પારવણીની કામગીરી હાથ ધરવી. કપાસનાં પાકમાં વરાપ થયે નિંદામણ અને આંતરઘેડ કરવી. કપાસમાં ખુણીયાં ટપકાં નાં રોગ માટે સ્ટ્રેપ્ટોસાઈકલીન ૧ ગ્રામ/૧૦ લી.પાણીમાં ઓગાળી છંટકાવ કરવો.
સોયાબીન	વૃદ્ધિ / ફૂલ	લશકરી ઈયળ, ઘોડિયા ઈયળ	<ul style="list-style-type: none"> ગર્ડલ બીટલ, કાતરા અને લશકરી ઈયળથી નુકશાનીત છોડવાઓ, ઈંડાઓનો સમૂહ અને ઈયળનો નાશ કરવો. વાવેલ પાકમાં ખાલા પુરવા અને છોડની પારવણીની કામગીરી હાથ ધરવી. સોયાબીનનું વાવેતર કર્યા બાદ ૩૦ દિવસે રીતે આંતરપાક તરીકે બે હાર પછી એક હાર એંરંડાનું વાવેતર કરવુ જેથી વધુ ઉત્પાદન મેળવી શકાય છે.

			<ul style="list-style-type: none"> ❖ સોયાબીનનાં પાકમાં વરાપ થયે નિંદામણ અને આંતરખેડ કરવી. ❖ સોયાબીનમાં લશકરી અને ધોડિયા ઈચ્છા માટે પ્રોફેનોફોસ ૫૦ ઇ.સી. અથવા કલોરાનદ્રાનીલીપ્રોલ ૧૮.૫ એસ.સી. અથવા એમામેકટીન બેન્જોએટ ૧.૬૦ ઇ.સી. અથવા ઇથીયોન ૫૦ ઇ.સી. અથવા ફલુબેન્ડીયામાઈડ ૩૬.૩૫ એસ.સી. અથવા ઇન્ડોકાકાર્બ ૧૫.૮૦ ઇ.સી. અથવા લેમડા-સાયહેલોથ્રીન ૪.૬૦ સી.એસ. અથવા સ્પીનેટોરમ ૧૧.૭૦ એસ.સી. અથવા નોવાલ્યુરોન ૫.૨૫+ ઇન્ડોકાકાર્બ ૪.૫૦ એસ.સી. અથવા થાયામીથોકામ ૧૨.૬૦ + લેમડા-સાયહેલોથ્રીન ૮.૫૦ એડ.સી. નો છંટકાવ કરવો.
મગા	વૃદ્ધિ	પાનનાં ટપકાનું	<ul style="list-style-type: none"> ❖ કંસિયા, સફેદ માખી અને તડતડીયા વગેરેના નિયંત્રણ માટે તુવેર, ગલગોટા અને દિવેલા જેવા પાકોનો આંતરપાક તરીકે ઉપયોગ કરવો. ❖ રક્ષક પાક તરીકે જીવાર અથવા મકાઈની ૪ હાર મગની આસપાસ કરવી. ❖ નાઈટ્રોજનયુક્ત ખાતરનો અપ્રમાણસર ઉપયોગ કરવો નહિં. ❖ મગનાં પાકમાં વરાપ થયે નિંદામણ અને આંતરખેડ કરવી. ❖ રોગની શરૂઆત થયે મેટીરામ પ૫% + પાયરેકલોસ્ટ્રોબીન પ૫% ડબલ્યુજી ૩૦ ગ્રામ/૧૦ લી. અથવા ટેબ્યુન્કોનાઓલ ૨૫.૮ ટકા અથવા હેકાકોનાઓલ ૫ ટકા ૧૦ મિલી/૧૦ લિટર મુજબ છંટકાવ કરવો.
અડન	વૃદ્ધિ	પાનનાં ટપકાનું	<ul style="list-style-type: none"> ❖ કંસિયા, સફેદ માખી અને તડતડીયા વગેરેના નિયંત્રણ માટે તુવેર, ગલગોટા અને દિવેલા જેવા પાકોનો આંતરપાક તરીકે ઉપયોગ કરવો. ❖ રક્ષક પાક તરીકે જીવાર અથવા મકાઈની ૪ હાર મગની આસપાસ કરવી. ❖ નાઈટ્રોજનયુક્ત ખાતરનો અપ્રમાણસર ઉપયોગ કરવો નહિં. ❖ અડનનાં પાકમાં વરાપ થયે નિંદામણ અને આંતરખેડ કરવી. ❖ રોગની શરૂઆત થયે મેટીરામ પ૫% + પાયરેકલોસ્ટ્રોબીન પ૫% ડબલ્યુજી ૩૦ ગ્રામ/૧૦ લી. અથવા ટેબ્યુન્કોનાઓલ ૨૫.૮ ટકા અથવા હેકાકોનાઓલ ૫ ટકા ૧૦ મિલી/૧૦ લિટર મુજબ છંટકાવ કરવો.

તલ	વૃક્ષ	તડતડીયા, સફેદ માખી	<ul style="list-style-type: none"> ❖ તલનાં પાકમાં વરાપ થયે નિંદામણ અને આંતરખેડ કરવી. ❖ તલમાં તડતડીયા અને સફેદ માખી માટે ઓકાઈડીમેટોન-મિથાઈલ ૨૫ ઈ.સી.નો છંટકાવ કરવો.
તુવેર	ઉગાવો	-	<ul style="list-style-type: none"> ❖ લીલી ઈયળના નિયંત્રણ માટે ગલગોટાનો પિંજર પાક તરીકે ઉપયોગમાં કરી શકાય.
એરંડા	ઉગાવો	-	<ul style="list-style-type: none"> ❖ વાવણી લાયક વરસાદ થયે એરંડાનું વાવેતર કરવું.
શાકભાજુ	વૃધ્ઘી/કુલા/ફળ	-	<ul style="list-style-type: none"> ❖ વેલાવાળા શાકભાજુના પાકમાં પાન કોરીયાનો ઉપદ્રવ ટાળવા માટે મુખ્ય પાકની ફરતે દિવેલા તથા ગલગોટાનું વાવેતર કરવાથી તેનું નિયંત્રણ કરી શકાય છે જો ઉપદ્રવ વધુ જોવા મળે તો ૧૦ મીલી મીથાઈલ ઓ ડેમેટોન (૨૫ ઈસી) અથવા ૧૦ મીલી ડાયમીથોએટ (૩૦ ઈસી) ૧૦ લીટર પાણીમાં લેળવી છંટકાવ કરવો ત્યાર બાદ પછીથી બીજો છંટકાવ ૧૫ દિવસ પછી કરવો. ❖ શાકભાજુના પાકમાં ચુસીયા પ્રકારની જીવાતના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦ મીલી અથવા લીબોળીનું મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦મીલી ૧૦ લી. પાણીમાં લેળવી છંટકાવ કરવો. જો વધુ ઉપદ્રવ હોય તો એસીટામ્રાપ્રીડ ૨૦ ટકા ૫ ગ્રામ/૧૦લી. અથવા દ્રાવણોફ્લોસ ૪૦ ટકા ૨૦ મીલી/૧૦ લી.પાણીમાં મીશ્ર કરી છંટકાવ કરવો. ❖ મરચીના પાકમાં થીપ્સના નિયંત્રણ માટે ૨૦મીલી પ્રોફેનોફ્લોસ અને લિમ્પોળીનું તેલ ૫૦ મીલી ૧૦ લીટર પાણી મિશ્રણ કરી છાંટવી. ❖ શાકભાજુના પાકની રોપણી બાદ ચુસીયા પ્રકારની જીવાતના નિયંત્રણ માટે દાણાદાર ફોરેટ ૧૦ જી છોડ દીઠ ૨ ગ્રામ મુજબ આપીને પીયત આપવું. ❖ મરચીના પાકમાં વાઈરસના કારણે પીળા થઈ કુકડાઈ ગયેલ છોડ ઉપાડી જમીનમાં દાટી નાશ કરવો અને ૫ મીલી ઈમીડાકલોપ્રાઈડ ૧૦ લીટર પાણીમાં લેળવી છંટકાવ કરવો. ❖ રીંગાણ અને ટમેટાના ફળ અને અને ડોકાની ઈયળોનો ઉપદ્રવ જણાય તો નુકશાન પામેલ ફળ અને ડોકાને તોડીને જમીનમાં દાટી દેવા અને રીનાક્ષીપાયર દવા ૩મીલીદવા ૧૦ લીટર પાણીમાં લેળવી છંટકાવ કરવો.

		<ul style="list-style-type: none"> ❖ રીંગાણ અને ટમેટાના પાકમા ફળમાખીના નિયંત્રણ માટે એસીટેમાપ્રીડ ૨૦ ટકા પગ્રામ દવા ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો. ❖ શાકભાજુ ને સમયસર ઉતારી ગ્રેડિંગ કરી બજારમાં મુકવા.
પશુપાલન	-	<ul style="list-style-type: none"> ❖ પશુઓને યોગ્ય આહાર આપવો & પીવા માટે ચોખ્યું પાણી આપવું અને પશુઓના રહેઠાણ ચોખ્યા રાખવા. ❖ માખી મચ્છરથી રક્ષણ માટે ફિનાઈલ છાટવુ. ❖ પશુઓને યોગ્ય આહારમાં ખનિજયુક્ત મિશ્રણ સાથે સંગ્રહિત ચારા આપો. ચારા ને સલામત સ્થળે સંગ્રહિત કરવો જોઈએ. ❖ ચારા માટે ચાફ કટર વાપરો. ❖ પશુઓમાં ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટાજ ૨ મીલી/ ૧ લીટર પાણીમાં નાખીને છાટવી. ❖ દુધાળા પશુઓના આઉને ઝીંક ઓક્સાઇડ અથલા બોરીક પાવડરથી બરાબર સાફ કરવા.

યાંત્રિક અધિકારી
ગ્રામીણ કૃષિ મૌસમ સેવા
જ.કૃ.યુ.



Gramin Krushi Mausam Seva (GKMS)
Agro meteorological Cell
Department of Agronomy
Junagadh Agricultural University
Junagadh-362001

E-mail: amfujnd@yahoo.com Phone: 0285-2670139



Agromet Advisory Service Bulletin for Junagadh District of South Saurashtra Agro-Climatic Zone

(Issued jointly by Junagadh Agricultural University and India Meteorological Department)

(#AZADI KA AMRIT MAHOTSAV#)

Day:- Tuesday

Date: 10-08-2021

Agromet Advisory Service bulletin. No:- 038 (2021-22)

Observed Weather during last Five days (Dated 05 th to 09 th August, 2021)					Weather Parameters	Weather Forecast upcoming Five days (Valid for 11 th to 15 th August, 2021) until 08:30 hrs of 15-08-2021				
05/08	06/08	07/08	08/08	09/08	Date	11/08	12/08	13/08	14/08	15/08
0	0	0	0	0	Rainfall(mm)	2	3	3	3	3
32.5	33.8	32.6	32.7	34.5	T _{max} (°C)	34	34	33	33	34
24.7	25.2	24.9	24.2	24.0	T _{min} (°C)	25	25	25	26	26
5	7	5	5	5	Cloud Cover	7	8	8	8	8
83	88	91	91	83	RH-I (%)	79	82	82	83	83
54	62	61	51	78	RH-II (%)	58	64	69	68	68
11.3	10.5	9.0	7.0	6.7	Wind speed (km/hr)	23.2	24.3	23.1	24.5	25.7
250	250	250	230	270	Wind direction	248	248	248	248	248

Agro meteorological Advisories

General Advice	Rain fall	As per IMD Forecast, There is forecasting of light rainfall occurs at isolated places during next five days. Sky will be remain mainly cloudy to cloudy.
	Temperature	Maximum and minimum temperature likely to be in range of 33 - 34°C and 25 - 26°C, respectively.
	RH (%)	The Relative Humidity is likely to be in the range of 58 – 83 %.
	Wind Direction and Wind Speed	Direction and speed of wind is mostly W and 23.1 – 25.7 km/h, respectively.

- Standardized Precipitation Index:** - For the period of **08th July to 04th August, 2021** for Junagadh district considered under the mildly wet condition.
- AS per ERFS products** **Normal rainfall may occur in South Saurashtra region during 15th to 21st August, 2021.**
- MEGHDOOT Mobile Application**:- **Download** the MEGHDOOT mobile application for Agromet Advisory Services and information on weather forecasting, which will provide weather based agricultural advice to farmers for their location, crops and livestock in their local language.

Click on the following link to download the MEGHDOOT application:

Android users: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>

IOS users: <https://apps.apple.com/in/app/meghdoot/id1474048155>

- DAMINI Mobile application**:- **Download** the DAMINI Mobile application to ensure the safety through monitoring sky ghtning activity. It will provide the alert and detail description of instruction, precaution, Do and Don't.


Click on the following link to download the DAMINI application.

Android users: <https://play.google.com/store/apps/details?id=com.lightening.live.damini>

IOS users: <https://apps.apple.com/app/id1502385645>

- Due to pandemic of CORONA (COVID-19), Farmers should maintain social distancing and safety precaution while handling farm machines and labour in the field.**
- Cleaning ridge, fencing and drainage system of the field during monsoon season.**
- Interculturing and hand weeding should be carried out in kharif crops.**
- To apply fertilizer, insecticide, pesticide and fungicide during vaspa condition.**

**NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 30
(23.07.2021 to 29.07.2021) over Agricultural region**



Source: <https://www.star.nesdis.noaa.gov>

Crop	Variety/ Pest/Insect	Stage	Agro met Advisories
Kharif Groundnut	White Grubs, Spodoptera, Helicoverpa	Vegetative / Flowering / Pegging	<ul style="list-style-type: none"> ➤ To Spray Quinolphos 25 EC 0.05% or Chlorpyrifos 20 EC or Flubendiamide 20 WG or Novaluron 5.25 + Indoxacarb 4.50 SC for managing defoliators like Spodoptera and Helicoverpa. ➤ Neem based application of Lambda-cyhalothrin 5 EC or or Imidacloprid 17.8 SL or Thiamethoxam 12.60 + Lambda-cyhalothrin 9.50 ZC for managing sucking pests. ➤ For the control of rust and tikka, spray fungicides mencozeb 26 gm mix in 10 litre of water at after sowing of 35 days. ➤ In Ground nut, spray 0.1% FeSo₄ @ 100 gram with 10 gram citric acid solution to correct chlorosis. ➤ Carry out inter culturing and weeding at vapsa condition. ➤ Installation of light trap in the field. ➤ Spray with Tebuconazole 25.9 EC or Propiconazole or Difenconazole 25 % EC 10 ml/10 lit at 30,50 and 75 days after sowing or Spray Pseudomonas fluorescens 10 ml/10 l.
Cotton	Leaf Spot, Sucking pest, Angular Leaf Spot	Vegetative / Branching	<ul style="list-style-type: none"> ➤ To control leaf spot, Spray Carbendazim 50 % WP 5 g/10 l. ➤ Avoid spraying of any chemical insecticides during initial crop growth stage if seeds treated with insecticide used for sowing

			<p>which gives protection up to initial 30-45 DAS.</p> <ul style="list-style-type: none"> ➤ During the initial phase of crop growth, natural enemies are abundant and keep the pest population under check ➤ Sprays of harmful chemicals destroy the natural enemies. ➤ Avoid repeated sprays of neonicotinic group of insecticides against sucking pests. ➤ If 25% plants are infested by sucking pest, spray Flonicamid 50 WG 4g/ 10 liter water. ➤ Gap filling and Thinning operation should be carried out in sown crop. ➤ Carry out inter culturing and weeding at vapsa condition. ➤ To Control Angular Leaf Spot, Spray Streptocycline 1 gm/10 l water.
Soybean	Tobacco Caterpillar, Green semiloopers	Vegetative / Flowering	<ul style="list-style-type: none"> ➤ Collect and destroy girdle beetle infested plant parts, egg masses and gregariously feeding larvae of hairy caterpillar and tobacco caterpillar. ➤ 30 days after sowing of soybean relay intercropping with castor planting can be obtained more production. ➤ Carry out inter culturing and weeding at vapsa condition. ➤ To Control Tobacco Caterpillar & Green semiloopers, Apply Profenofos 50 EC or Chlorantraniliprole 18.5 SC or Emamectin benzoate 1.90 EC or Ethion 50 EC or Flubendiamide 39.35 SC or Indoxacarb 15.80

			EC or Lambda-cyhalothrin 4.90 CS or Spinetoram 11.70 SC or Novaluron 5.25 + Indoxacarb 4.50 SC or Thiamethoxam 12.60 +Lambda-cyhalothrin 9.50 ZC.
Black gram	Leaf spot	Vegetative	<ul style="list-style-type: none"> ➤ Growing intercrops such as Pigeon pea, marigold and castor for the control of blister beetle, whitefly and leaf hoppers. ➤ Guard crop sorghum or maize in 4 rows all around green gram crop as guard crop. ➤ Avoid excessive use of nitrogenous fertilizers. ➤ Carry out inter culturing and weeding at vapsa condition. ➤ To Spray Metiram 55% + Pyraclostrobin 5% WG 30 g/10 l or Tebuconazole 25.9 EC or Hexaconazole 5 EC 10 ml/10 l control of leaf spot.
Green Gram	Leaf spot	Vegetative	<ul style="list-style-type: none"> ➤ Growing intercrops such as Pigeon pea, marigold and castor for the control of blister beetle, whitefly and leaf hoppers. ➤ Guard crop sorghum or maize in 4 rows all around green gram crop as guard crop. ➤ Avoid excessive use of nitrogenous fertilizers. ➤ Carry out inter culturing and weeding at vapsa condition. ➤ To Spray Metiram 55% + Pyraclostrobin 5% WG 30 g/10 l or Tebuconazole 25.9 EC or

			Hexaconazole 5 EC 10 ml/10 l control of leaf spot.
Sesame	Leaf hopper, White fly	Vegetative	<ul style="list-style-type: none"> ➤ Carry out inter culturing and weeding at vapsa condition. ➤ To Spray Oxydemeton-methyl 25 EC for the control of leaf hopper & white fly.
Pigeon pea	-	Germination	<ul style="list-style-type: none"> ➤ Marigold plantation should be adopted as trap crop for pod borer. ➤ Collect and destroy eggs and early stage larvae during early stages of the crop.
Castor	-	Germination	<ul style="list-style-type: none"> ➤ Timely sowing should be done at the time of sufficient rainfall.
Vegetable crops	-	Sowing/ Flowering/ Fruiting	<ul style="list-style-type: none"> ➤ For the control of leaf webber in fruit leafy vegetable, sowing of trap crop (castor and galgota) around the vegetable crop. For the heavy incidence spray 10 ml methyl O demeton (25 EC) Or 10 ml diamethoate (30 EC) in 10 liter of water. ➤ For the control of thrips and white fly, spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. ➤ For the control of thrips in chilli, spray 20 ml profenophos in 10 litres of water. ➤ For the control of fruit and shoot borer, spray neem seed extract 5 % or neem seed oil 50 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.

			<ul style="list-style-type: none"> ➤ Rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of 5 ml Imidacloprid per 10 litre of water is advised for control the vector, when sky remains clear. ➤ Apply regular irrigation at 6 to 8 days to the crop at the flowering and fruiting stage. ➤ In brinjal and tomato crops constant monitoring of borers by using pheromone traps 4-6 per acre is advised, for the control of fruit borer, spray reenaxipayar @ 3.0ml in 10 lit of water. ➤ In brinjal and tomato crops, for the control of fruit fly, spray acetemaphrid 5.0 ml in 10 lit of water. ➤ Keep vegetable crop weed free, apply irrigation regularly and send vegetable fruit to the market often grading.
Livestock	-	-	<ul style="list-style-type: none"> ➤ To provide adequate food & clean water for drinking and keep the animals clean shed. ➤ Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Give stored fodder with mineral mixture. Fodder should be stored at safe place. ➤ Feed silage & hay material along with concentrate feed. ➤ Use chaff cutter for fodder.

			<ul style="list-style-type: none"> ➤ For the control of tick and other parasites, spray deltemethrin or aemitaz 2 ml in 1 litre of water. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.
--	--	--	--

 Gramin Krushi Mausam Seva (GKMS) Agro meteorological Cell Department of Agronomy Junagadh Agricultural University Junagadh-362001 E-mail: amfujnd@yahoo.com Phone: 0285-2670139	
---	---

Agromet Advisory Service Bulletin for Porbandar District of South Saurashtra Agro-Climatic Zone
(Issued jointly by Junagadh Agricultural University and India Meteorological Department)

(#AZADI KA AMRIT MAHOTSAV #)

Day:- Tuesday

Date: 10-08-2021

Agromet Advisory Service bulletin. No:- 038 (2021-22)

Observed Weather during last Five days (Dated 05 th to 09 th August, 2021)					Weather Parameters	Weather Forecast upcoming Five days (Valid for 11 th to 15 th August, 2021) until 08:30 hrs of 15-08-2021				
05/08	06/08	07/08	08/08	09/08	Date	11/08	12/08	13/08	14/08	15/08
0	0	0	2.0	0	Rainfall(mm)	2	2	3	3	3
31.0	31.0	30.8	32.0	31.6	Tmax (°C)	32	31	31	31	30
26.5	26.8	26.5	26.2	26.5	Tmin (°C)	27	27	27	26	26
7	7	7	6	5	Cloud Cover	8	7	8	8	8
83	83	80	87	79	RH-I (%)	78	80	80	80	81
82	82	77	76	74	RH-II (%)	66	69	73	73	72
18.5	22.2	14.8	14.8	11.1	Wind speed (km/hr)	27.1	28.0	27.0	28.3	30.2
250	250	250	230	250	Wind direction	248	248	248	248	248

Agro meteorological Advisories

General Advice	Rain fall	As per IMD Forecast, There is forecasting light rainfall occurs at isolated places during next five days. Sky will be remain mainly cloudy to cloudy.
	Temperature	

	RH (%)	Maximum and minimum temperature likely to be in range of 30 - 32°C and 26 - 27°C, respectively.
	Wind Direction and Wind Speed	The Relative Humidity is likely to be in the range of 66 – 81 %. Direction and speed of wind is to be mostly W and 27.0 – 30.2 km/h, respectively.

- ❖ **Standardized Precipitation Index:** - For the period **08th July to 04th August, 2021** for Porbandar district considered under the mildly wet condition.
- ❖ **AS per ERFS products** **Normal rainfall may occur in South Saurashtra region during 15th to 21st August, 2021.**
- ❖ **MEGHDOOT Mobile Application**:- **Download** the MEGHDOOT mobile application for Agromet Advisory Services and information on weather forecasting, which will provide weather based agricultural advice to farmers for their location, crops and livestock in their local language.

Click on the following link to download the MEGHDOOT application:

Android users: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>

IOS users: <https://apps.apple.com/in/app/meghdoot/id1474048155>

- ❖ **DAMINI Mobile application**:- **Download** the DAMINI Mobile application to ensure the safety through monitoring sky ghtning activity. It will provide the alert and detail description of instruction, precaution, Do and Don't.

Click on the following link to download the DAMINI application.


Android users: <https://play.google.com/store/apps/details?id=com.lightening.live.damini>

IOS users: <https://apps.apple.com/app/id1502385645>

- ❖ **Due to pandemic of CORONA (COVID-19), Farmers should maintain social distancing and safety precaution while handling farm machines and labour in the field.**
- ❖ **Cleaning ridge, fencing and drainage system of the field during monsoon season.**
- ❖ **Interculturing and hand weeding should be carried out in kharif crops.**
- ❖ **To apply fertilizer, insecticide, pesticide and fungicide during vaspa condition.**

**NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 30
(23.07.2021 to 29.07.2021) over Agricultural region**

Gujarat



Agricultural vigour is moderate over isolated places of the state.

Source: <https://www.star.nesdis.noaa.gov>

Crop	Variety/ Pest / Insect	Stage	Agro met Advisories
Kharif Groundnut	White Grubs, Spodoptera, Helicoverpa	Vegetative / Flowering / Pegging	<ul style="list-style-type: none"> ➤ To Spray Quinolphos 25 EC 0.05% or Chlorpyrifos 20 EC or Flubendiamide 20 WG or Novaluron 5.25 + Indoxacarb 4.50 SC for managing defoliators like Spodoptera and Helicoverpa. ➤ Neem based application of Lambda-cyhalothrin 5 EC or Imidacloprid 17.8 SL or Thiamethoxam 12.60 + Lambda-cyhalothrin 9.50 ZC for managing sucking pests. ➤ For the control of rust and tikka, spray fungicides mencozeb 26 gm mix in 10 litre of water at after sowing of 35 days. ➤ In Ground nut, spray 0.1% FeSo₄ @ 100 gram with 10 gram citric acid solution to correct chlorosis. ➤ Carry out inter culturing and weeding at vapsa condition. ➤ Installation of light trap in the field. ➤ Spray with Tebuconazole 25.9 EC or Propiconazole or Difenconazole 25 % EC 10 ml/10 lit at 30,50 and 75 days after sowing

			or Spray Pseudomonas fluorescens 10 ml/10 l.
Cotton	Leaf Spot, Sucking pest, Angular Leaf Spot	Vegetative / Branching	<ul style="list-style-type: none"> ➤ To control leaf spot, Spray Carbendazim 50 % WP 5 g/10 l. ➤ Avoid spraying of any chemical insecticides during initial crop growth stage if seeds treated with insecticide used for sowing which gives protection up to initial 30-45 DAS. ➤ During the initial phase of crop growth, natural enemies are abundant and keep the pest population under check ➤ Sprays of harmful chemicals destroy the natural enemies. ➤ Avoid repeated sprays of neonicotinic group of insecticides against sucking pests. ➤ If 25% plants are infested by sucking pest, spray Flonicamid 50 WG 4g/ 10 liter water. ➤ Gap filling and Thinning operation should be carried out in sown crop. ➤ Carry out inter culturing and weeding at vapsa condition. ➤ To Control Angular Leaf Spot, Spray Streptocycline 1 gm/10 l water.
Soybean	Tobacco Caterpillar, Green semiloopers	Vegetative / Flowering	<ul style="list-style-type: none"> ➤ Collect and destroy girdle beetle infested plant parts, egg masses and gregariously feeding larvae of hairy caterpillar and tobacco caterpillar.

			<ul style="list-style-type: none"> ➤ 30 days after sowing of soybean relay intercropping with castor planting can be obtained more production. ➤ Carry out inter culturing and weeding at vapsa condition. ➤ To Control Tobacco Caterpillar & Green semiloopers, Apply Profenofos 50 EC or Chlorantraniliprole 18.5 SC or Emamectin benzoate 1.90 EC or Ethion 50 EC or Flubendiamide 39.35 SC or Indoxacarb 15.80 EC or Lambda-cyhalothrin 4.90 CS or Spinetoram 11.70 SC or Novaluron 5.25 + Indoxacarb 4.50 SC or Thiamethoxam 12.60 +Lambda-cyhalothrin 9.50 ZC.
Black gram	Leaf spot	Vegetative	<ul style="list-style-type: none"> ➤ Growing intercrops such as Pigeon pea, marigold and castor for the control of blister beetle, whitefly and leaf hoppers. ➤ Guard crop sorghum or maize in 4 rows all around green gram crop as guard crop. ➤ Avoid excessive use of nitrogenous fertilizers. ➤ Carry out inter culturing and weeding at vapsa condition. ➤ To Spray Metiram 55% + Pyraclostrobin 5% WG 30 g/10 l or Tebuconazole 25.9 EC or Hexaconazole 5 EC 10 ml/10 l control of leaf spot.
Green Gram	Leaf spot	Vegetative	<ul style="list-style-type: none"> ➤ Growing intercrops such as Pigeon pea, marigold and castor for the control of blister beetle, whitefly and leaf hoppers. ➤ Guard crop sorghum or maize in 4 rows all around green gram crop as guard crop.

			<ul style="list-style-type: none"> ➤ Avoid excessive use of nitrogenous fertilizers. ➤ Carry out inter culturing and weeding at vapsa condition. ➤ To Spray Metiram 55% + Pyraclostrobin 5% WG 30 g/10 l or Tebuconazole 25.9 EC or Hexaconazole 5 EC 10 ml/10 l control of leaf spot.
Sesame	Leaf hopper, White fly	Vegetative	<ul style="list-style-type: none"> ➤ Carry out inter culturing and weeding at vapsa condition. ➤ To Spray Oxydemeton-methyl 25 EC for the control of leaf hopper & white fly.
Pigeon pea	-	Germination	<ul style="list-style-type: none"> ➤ Marigold plantation should be adopted as trap crop for pod borer. ➤ Collect and destroy eggs and early stage larvae during early stages of the crop.
Castor	-	Germination	<ul style="list-style-type: none"> ➤ Timely sowing should be done at the time of sufficient rainfall.
Vegetable crops	-	Sowing/ Flowering/ Fruiting	<ul style="list-style-type: none"> ➤ For the control of leaf webber in fruit leafy vegetable, sowing of trap crop (castor and galgota) around the vegetable crop. For the heavy incidence spray 10 ml methyl O demeton (25 EC) Or 10 ml diamethoate (30 EC) in 10 liter of water. ➤ For the control of thrips and white fly, spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. ➤ For the control of thrips in chilli, spray 20 ml profenophosin 10 litres of water.

			<ul style="list-style-type: none"> ➤ For the control of fruit and shoot borer, spray neem seed extract 5 % or neem seed oil 50 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. ➤ Rouging of viralinfested crop should be done in chilli. Infectedplants should be uprooted and buried in the soil.After rouging, spraying of 5 mi Imidacloprid per 10litre of water is advisedfor control thevector, when sky remains clear. ➤ Apply regular irrigation at 6 to 8 days to the crop at the flowering and fruiting stage. ➤ In brinjal and tomato crops constant monitoring of borers by using pheromone traps 4-6 per acre is advised, for the control of fruit borer, spray reenaxipayar @ 3.0ml in 10 lit of water. ➤ In brinjal and tomato crops, for the control of fruit fly, spray acetemaprid 5.0 ml in 10 lit of water. ➤ Keep vegetable crop weed free, apply irrigation regularly and send vegetable fruit to the market often grading.
Livestock	-	-	<ul style="list-style-type: none"> ➤ To provide adequate food & clean water for drinking and keep the animals clean shed. ➤ Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Give stored fodder with mineral mixture. Fodder should be stored at safe place.

			<ul style="list-style-type: none"> ➤ Feed silage & hay material along with concentrate feed. ➤ Use chaff cutter for fodder. ➤ For the control of tick and other parasites, spray deltemethrin or aemitaz 2 ml in 1 litre of water. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.
--	--	--	---

	Gramin Krushi Mausam Seva (GKMS) Agrometeorological Cell Department of Agronomy Junagadh Agricultural University Junagadh-362001 E-mail: amfujnd@yahoo.com Phone: 0285-2670139	
---	---	--

Agromet Advisory Service Bulletin for Gir Somnath District of South Saurashtra Agro-Climatic Zone

(Issued jointly by Junagadh Agricultural University and India Meteorological Department)

(#AZADI KA AMRIT MAHOTSAV#)

Day:- Tuesday

Date: 10-08-2021

Agromet Advisory Service bulletin. No:- 038 (2021-22)

Observed Weather during last Five days (Dated 05 th to 09 th August, 2021)					Weather Parameters	Weather Forecast upcoming Five days (Valid for 11 th to 15 th August, 2021) until 08:30 hrs of 15-08-2021				
05/08	06/08	07/08	08/08	09/08	Date	11/08	12/08	13/08	14/08	15/08
0	0	0	0	0	Rainfall(mm)	2	3	3	3	3
30.1	30.8	30.6	31.5	32.6	Tmax (°C)	34	34	33	33	34
26.9	26.8	26.9	26.9	26.8	Tmin (°C)	25	25	25	26	26
5	7	5	5	5	Cloud Cover	7	8	8	8	8
86	86	89	89	82	RH-I (%)	82	84	84	84	84
83	82	83	82	74	RH-II (%)	65	68	71	70	71
11.1	9.3	5.6	-	3.7	Wind speed (km/hr)	22.5	24.1	23.5	24.9	26.3
250	250	250	-	99	Wind direction	248	248	248	248	248

Agro meteorological Advisories

General Advice	Rain fall	As per IMD Forecast, There is forecasting of light rainfall occurs at isolated places during
-----------------------	------------------	---

	Temperature	next five days. Sky will be remain mainly cloudy to cloudy. Maximum and minimum temperature likely to be in range of 33 - 34°C and 25 - 26°C, respectively.
	RH (%)	The Relative Humidity is likely to be in the range of 65 – 84 %.
	Wind Direction and Wind Speed	Direction and speed of wind is to be mostly W and 22.5 – 26.3 km/h, respectively.

+ **Standardized Precipitation Index**: - For the period of **08th July to 04th August, 2021** for Gir Somnath district considered under the mildly wet condition.

+ **AS per ERFS products Normal rainfall may occur in South Saurashtra region during 15th to 21st August, 2021.**

+ **MEGHDOOT Mobile Application**:- **Download** the MEGHDOOT mobile application for Agromet Advisory Services and information on weather forecasting, which will provide weather based agricultural advice to farmers for their location, crops and livestock in their local language.

Click on the following link to download the MEGHDOOT application:

Android users: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>

IOS users: <https://apps.apple.com/in/app/meghdoot/id1474048155>

+ **DAMINI Mobile application**:- **Download** the DAMINI Mobile application to ensure the safety through monitoring sky lightning activity. It will provide the alert and detail description of instruction, precaution, Do and Don't.


Click on the following link to download the DAMINI application.

Android users: <https://play.google.com/store/apps/details?id=com.lightening.live.damini>

IOS users: <https://apps.apple.com/app/id1502385645>

- ❖ **Due to pandemic of CORONA (COVID-19), Farmers should maintain social distancing and safety precaution while handling farm machines and labour in the field.**
- ❖ **Cleaning ridge, fencing and drainage system of the field during monsoon season.**
- ❖ **Interculturing and hand weeding should be carried out in kharif crops.**
- ❖ **To apply fertilizer, insecticide, pesticide and fungicide during vaspa condition.**

**NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 30
(23.07.2021 to 29.07.2021) over Agricultural region**



Agricultural vigour is moderate over isolated places of the state.

Source: <https://www.star.nesdis.noaa.gov>

Crop	Variety/ Pest / Insect	Stage	Agro met Advisories
Kharif Groundnut	White Grubs, Spodoptera, Helicoverpa, Early & Late leaf spot	Vegetative / Flowering / Pegging	<ul style="list-style-type: none"> ➤ To Spray Quinolphos 25 EC 0.05% or Chlorpyrifos 20 EC or Flubendiamide 20 WG or Novaluron 5.25 + Indoxacarb 4.50 SC for managing defoliators like Spodoptera and Helicoverpa. ➤ Need based application of Lambda-cyhalothrin 5 EC or Imidacloprid 17.8 SL or Thiamethoxam 12.60 + Lambda-cyhalothrin 9.50 ZC for managing sucking pests. ➤ For the control of rust and tikka, spray fungicides mencozeb 26 gm mix in 10 litre of water at after sowing of 35 days. ➤ In Ground nut, spray 0.1% FeSo₄ @ 100 gram with 10 gram citric acid solution to correct chlorosis. ➤ Carry out inter culturing and weeding at vapsa condition. ➤ Installation of light trap in the field. ➤ Spray with Tebuconazole 25.9 EC or Propiconazole or Difenconazole 25 % EC 10 ml/10 lit at 30,50 and 75 days after sowing or Spray Pseudomonas fluorescens 10 ml/10 l.

Cotton	Leaf Spot, Sucking pest, Angular Leaf Spot	Vegetative / Branching	<ul style="list-style-type: none"> ➤ To control leaf spot, Spray Carbendazim 50 % WP 5 g/10 l. ➤ Avoid spraying of any chemical insecticides during initial crop growth stage if seeds treated with insecticide used for sowing which gives protection up to initial 30-45 DAS. ➤ During the initial phase of crop growth, natural enemies are abundant and keep the pest population under check ➤ Sprays of harmful chemicals destroy the natural enemies. ➤ Avoid repeated sprays of neonicotinic group of insecticides against sucking pests. ➤ If 25% plants are infested by sucking pest, spray Flonicamid 50 WG 4g/ 10 liter water. ➤ Gap filling and Thinning operation should be carried out in sown crop. ➤ Carry out inter culturing and weeding at vapsa condition. ➤ To Control Angular Leaf Spot, Spray Streptocycline 1 gm/10 l water.
Soybean	Tobacco Caterpillar, Green semiloopers	Vegetative / Flowering	<ul style="list-style-type: none"> ➤ Collect and destroy girdle beetle infested plant parts, egg masses and gregariously feeding larvae of hairy caterpillar and tobacco caterpillar. ➤ 30 days after sowing of soybean relay intercropping with castor planting can be obtained more production. ➤ Carry out inter culturing and weeding at vapsa condition.

			<ul style="list-style-type: none"> ➤ To Control Tobacco Caterpillar & Green semiloopers, Apply Profenofos 50 EC or Chlorantraniliprole 18.5 SC or Emamectin benzoate 1.90 EC or Ethion 50 EC or Flubendiamide 39.35 SC or Indoxacarb 15.80 EC or Lambda-cyhalothrin 4.90 CS or Spinetoram 11.70 SC or Novaluron 5.25 + Indoxacarb 4.50 SC or Thiamethoxam 12.60 +Lambda-cyhalothrin 9.50 ZC.
Black gram	Leaf spot	Vegetative	<ul style="list-style-type: none"> ➤ Growing intercrops such as Pigeon pea, marigold and castor for the control of blister beetle, whitefly and leaf hoppers. ➤ Guard crop sorghum or maize in 4 rows all around green gram crop as guard crop. ➤ Avoid excessive use of nitrogenous fertilizers. ➤ Carry out inter culturing and weeding at vapsa condition. ➤ To Spray Metiram 55% + Pyraclostrobin 5% WG 30 g/10 l or Tebuconazole 25.9 EC or Hexaconazole 5 EC 10 ml/10 l control of leaf spot.
Green Gram	Leaf spot	Vegetative	<ul style="list-style-type: none"> ➤ Growing intercrops such as Pigeon pea, marigold and castor for the control of blister beetle, whitefly and leaf hoppers. ➤ Guard crop sorghum or maize in 4 rows all around green gram crop as guard crop. ➤ Avoid excessive use of nitrogenous fertilizers. ➤ Carry out inter culturing and weeding at vapsa condition. ➤ To Spray Metiram 55% + Pyraclostrobin 5% WG 30 g/10 l or Tebuconazole 25.9 EC or

			Hexaconazole 5 EC 10 ml/10 l control of leaf spot.
Sesame	Leaf hopper, White fly	Vegetative	<ul style="list-style-type: none"> ➤ Carry out inter culturing and weeding at vapsa condition. ➤ To Spray Oxydemeton-methyl 25 EC for the control of leaf hopper & white fly.
Pigeon pea	-	Germination	<ul style="list-style-type: none"> ➤ Marigold plantation should be adopted as trap crop for pod borer. ➤ Collect and destroy eggs and early stage larvae during early stages of the crop.
Castor	-	Germination	<ul style="list-style-type: none"> ➤ Timely sowing should be done at the time of sufficient rainfall.
Vegetable crops	-	Sowing/ Flowering/ Fruiting	<ul style="list-style-type: none"> ➤ For the control of leaf webber in fruit leafy vegetable, sowing of trap crop (castor and galgota) around the vegetable crop. For the heavy incidence spray 10 ml methyl O demeton (25 EC) Or 10 ml diamethoate (30 EC) in 10 liter of water. ➤ For the control of thrips and white fly, spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. ➤ For the control of thrips in chilli, spray 20 ml profenophosin10 litres of water. ➤ For the control of fruit and shoot borer, spray neem seed extract 5 % or neem seed oil 50 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.

			<ul style="list-style-type: none"> ➤ Rouging of viralinfested crop should be done in chilli. Infectedplants should be uprooted and buried in the soil. After rouging, spraying of 5 mi Imidacloprid per 10litre of water is advisedfor control thevector, when sky remains clear. ➤ Apply regular irrigation at 6 to 8 days to the crop at the flowering and fruiting stage. ➤ In brinjal and tomato crops constant monitoring of borers by using pheromone traps 4-6 per acre is advised, for the control of fruit borer, spray reenaxipayar @ 3.0ml in 10 lit of water. ➤ In brinjal and tomato crops, for the control of fruit fly, spray acetemaphrid 5.0 ml in 10 lit of water. ➤ Keep vegetable crop weed free, apply irrigation regularly and send vegetable fruit to the market often grading.
Livestock	-	-	<ul style="list-style-type: none"> ➤ To provide adequate food & clean water for drinking and keep the animals clean shed. ➤ Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Give stored fodder with mineral mixture. Fodder should be stored at safe place. ➤ Feed silage & hay material along with concentrate feed. ➤ Use chaff cutter for fodder.

			<ul style="list-style-type: none">➤ For the control of tick and other parasites, spray deltemethrin or aemitaz 2 ml in 1 litre of water.➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.
--	--	--	---

**Technical Officer
Gramin Krushi Mausam Seva (GKMS)
J.A.U., Junagadh**