



75
आज़ादी का
अमृत महोत्सव

ग्रामीણ कृषि मौसम सेवा
कृषि हवामान सेल
कृषि विज्ञान विभाग
जुनागढ कृषि युनिवर्सिटी
जुनागढ - 352001

ईमेल: - amfujnd@yhaoo.com फ़ोन नं. 0224-2590936



G20
भारत 2023 INDIA

हवामान आगाही आधारित दक्षिण सौराष्ट्र भेत् आबोहवाकीय विस्तारनां **जूनागढ जुल्ला** माटे कृषि सलाह बुलेटीन
(कृषि हवामान सेल, जूनागढ कृषि युनिवर्सिटी अने भारत हवामान विभाग द्वारा संपन्न)

वार:- शुक्रवार

तारीख:- 21/03/2024

हवामान आगाही आधारित बुलेटीन: **102 (2024-25)**

पाछला पांच दिवसनुं हवामान (तारीख:- 15 थी 20 मार्च, 2024)					हवामान परिबर्णो	हवामान आगाही आगामी पांच दिवस माटे (तारीख:- 22 थी 26 मार्च, 2024) मान्य 26 मार्च, 2024 : सवारना 6:30 वाज्यां सुधी				
15/03	16/03	17/03	18/03	20/03	तारीख	22/03	23/03	24/03	25/03	26/03
0	0	0	0	0	वरसाद(मिमी)	0	0	0	0	0
39.5	35.4	35.0	39.1	39.0	महतम तापमान (°स.ग.)	36	40	40	40	36
21.0	21.3	23.6	21.4	16.1	लघुतम तापमान (°स.ग.)	22	23	21	16	16
योग्य	योग्य	योग्य	योग्य	योग्य	वादननी स्थिति	योग्य	योग्य	योग्य	योग्य	योग्य
49	48	36	44	41	सवारना लज्जा टका	50	50	50	54	40
20	18	19	11	19	अपारवादन लज्जा टका	20	14	14	20	30
4.4	5.9	5.6	4.6	8.6	पवनना गत (किमी/कलाक)	9	5	9	5	6
320	350	260	320	260	पवननी दिशा	325	349	16	349	316

सामान्य सलाह

भारत सरकारश्रीना हवामान विभाग तरङ्गथी मजेल हवामान आगाहीनी माहीती अनुसार अत्रेनां ग्रामीण कृषि मौसम सेवा योजना, जूनागढ कृषि युनिवर्सिटी, जूनागढ तरङ्गथी जणाववामा आवे छेके दक्षिण सौराष्ट्र भेत् आबोहवाकीय विस्तारना जूनागढ जुल्लामां आगामी तारीख **22/03/2024 थी 26/03/2024** दरम्यान आकाश मुभ्यत्वे आगामी पांच दिवसोमां मुभ्यत्वे **योग्य** रहेशे, जुल्लामां आगामी पांच दिवसो दरम्यान **वरसादनी** शक्यताओ **नहिवत** छे. पवननी झडप **05 थी 06** कीमी/कलाक रहेवानी शक्यता रहेशे. आ समयगाणामां महतम तापमान **36 थी 40** डीग्री सेन्टीग्रेड रहेशे तेमज लघुतम तापमान **16 थी 23** डीग्री सेन्टीग्रेड रहेवानी संभावना छे.

✦ માન્ય વરસાદની સૂચી મુજબ જુનાગઢ જિલ્લામાં ૨૬ ડિસેમ્બર, ૨૦૨૪ થી ૨૨ જાન્યુઆરી, ૨૦૨૫ નાં સમયગાળા દરમિયાન હળવી સૂકી સ્થિતિ જોવા મળેલ છે.

✦ સાપ્તાહિક હવામાન આગાહીની સૂચી મુજબ સૌરાષ્ટ્ર અને કચ્છનાં વિસ્તારોમાં તા. ૨૬/૦૩/૨૦૨૫ થી તા. ૦૧/૦૪/૨૦૨૫ દરમિયાન વરસાદ નથી & મહત્તમ તાપમાન સામાન્ય થી નીચે અને લઘુત્તમ તાપમાન સામાન્ય રહેવાની શક્યતાઓ છે.

✦ મેઘદૂત એપ્લિકેશન:-કૃષિ હવામાન સલાહ સેવા અને હવામાન આગાહી અંગેની જાણકારીમેળવવામાટે ડાઉનલોડ કરો મેઘદૂત એપ્લિકેશન કે જે ખેડૂતોને તેમનીસ્થાનિક ભાષામાંસ્થાન, પાક અને પશુધન વિશે હવામાન આધારિત કૃષિ સલાહ પ્રદાન કરશે.

મેઘદૂત એપ્લિકેશનનેડાઉનલોડ કરવા માટે નીચેની લિંક પર ક્લિક કરો.

એન્ડ્રોઈડ(મોબાઈલ)ઉપયોગકર્તા:<https://play.google.com/store/apps/details?id=com.aas.meghdoot>


આઈ.ઓ.એસ.ઉપયોગકર્તા:-<https://apps.apple.com/in/app/meghdoot/id1474048155>

❖ શિયાળું પાકોમાં, દેહધાર્મિક પરિપક્વ અવસ્થાએ કાપણીની કામગીરી કરવી.

❖ શિયાળુ પાકોની કાપણી થઈ ગયા બાદ આખા ખેતરનું અવલોકન કરી જમીનના નમુના લેવા અને પૃથ્થકરણ કરવા માટે લેબોરેટરીમાં મોકલી આપવા.

❖ ઉનાળુ પાકોના વાવેતર માટે ખેત આયોજન કરવું તે મુજબ બીયારણ, ખાતર અને દવાનો પુરવઠો મેળવી લેવો.

❖ ખેડૂતોને ઉનાળુ પાકોનાં વાવેતર માટે ખેતરો તૈયાર કરવા સલાહ આપવામાં આવે છે.



પાકનું નામ	પાક અવસ્થા	રોગ / જીવાત / જાત	કૃષિ સલાહ
ઉનાળું મગફળી	વૃદ્ધિ / ફૂલ	ચૂસીયા જીવાત	❖ પાકને નિયમિત સમયાંતરે પિયત આપવાની કામગીરી કરવી. ❖ મગફળીમાં ચુસીયા, મોલોમશી, થ્રીપ્સ ના નિયંત્રણ માટે

			ફોસ્ફામીડોન ૩ મીલી દવા અથવા ડાયમીથોએટ ૧૫ મીલી દવા ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો. મોલોમશી સાથે દાળીયા જીવાત જોવા મળે તો દાવાનો છંટકાવ મુલતવી રાખવો. મગફળીના પાકમાં આંતરખેડ કરવી અને હાથથી નિદામણ કરવું.
ઉનાળું મગ	ઉગાવો / વૃદ્ધિ	ચૂસીયા જીવાત	❖ પાકને સમયાંતરે પિયત આપવાની કામગીરી કરવી. ❖ ચૂસીયા જીવાતોના નિયંત્રણ માટે લીમડાનું તેલ @૫ મિલી/ લીટર મુજબ છાંટવું. પાકમાં આંતરખેડ કરવી અને હાથથી નિદામણની કામગીરી કરવી.
ઉનાળું અડદ	ઉગાવો / વૃદ્ધિ	ચૂસીયા જીવાત	❖ પાકને સમયાંતરે પિયત આપવાની કામગીરી કરવી. ❖ ચૂસીયા જીવાતોના નિયંત્રણ માટે લીમડાનું તેલ @૫ મિલી/ લીટર મુજબ છાંટવું. પાકમાં આંતરખેડ કરવી અને હાથથી નિદામણની કામગીરી કરવી.
ઉનાળું તલ	ઉગાવો / વૃદ્ધિ	તડતડિયા / સફેદમાખી	❖ પાકને સમયાંતરે પિયત આપવાની કામગીરી કરવી. પાકમાં તડતડિયા અને સફેદમાખીનાં નિયંત્રણ માટે ઓક્ઝીડીમેટોન - મિથાઈલ ૨૫ ઇ.સી.નો છંટકાવ કરવો. પાકમાં આંતરખેડ કરવી અને હાથથી નિદામણની કામગીરી કરવી.
આંબા	ફૂલ / ફળ અવસ્થા	મધીયા & ભુકીછારા	❖ આંબામાં મધીયાના નિયંત્રણમાટે ૫૦ % કાર્બારીલ ૪૦ ગ્રામ પાવડર અથવા ૩૬ % ઇ.સી. મોનોક્રોટોફોસ ૧૦ મીલી દવા ૧૦ લી.પાણીમાં ભેળવી છંટકાવ કરવો. ❖ આંબામાં ભુકીછારા ના નિયંત્રણ માટે ૧૦ લી. પાણીમાં ૩૦ ગ્રામ દ્રાવ્ય ગંધક + કેરેથોન ૪૮ % ઇ.સી. ૧૦ મીલી દવાનો છંટકાવ કરવો. ❖ આંબાને નિયમિત સમયાંતરે પિયત આપવાની કામગીરી કરવી. કેરી સોપારી જેવડા કદની થાય એટલે ૨૦ પી.પી.એમ. એન.એ.એ. અને ૨% ચુરીયા ના દ્રાવણનો છંટકાવ કરવાથી કેરી ખરતી અટકે છે અને ફળના કદમાં વધારો થાય છે.
શાકભાજી	વૃદ્ધી/ફુલ/ ફળ	-	❖ શાકભાજીના પાકોને નીદામણ મુક્ત કરવું. શાકભાજી ને સમયસર ઉતારી ગ્રેડીંગ કરી બજારમાં મુકવા. ❖ પાકને નિયમિત સમયાંતરે પિયત આપવાની કામગીરી કરવી. આગાહી સમયગાળા દરમિયાન ખેતર પરનાં બધાં ખેતકાર્યો કરવા.

પશુપાલન	-	-	❖ પશુઓને યોગ્ય આહાર આપવો & પીવા માટે ચોખ્ખું પાણી આપવું અને પશુઓના રહેઠાણ ચોખ્ખા રાખવા. ❖ માખી મચ્છરથી રક્ષણ માટે ફીનાઈલ છાટવું. ❖ પશુઓને યોગ્ય આહારમાં ખનિજયુક્ત મિશ્રણ સાથે સંગ્રહિત ચારા આપો. ચારા ને સલામત સ્થળે સંગ્રહિત કરવો જોઈએ. ❖ ચારા માટે ચાફ કટર વાપરો. ❖ પશુઓમાં ઇતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટાઝ ૨ મીલી/ ૧ લીટર પાણીમાં નાખીને છાટવી. દુધાળા પશુઓના આઉને ઝીંક ઓક્સાઈડ અથવા બોરીક પાવડરથી બરાબર સાફ કરવા.
---------	---	---	--



75
આઝાદી કા
અમૃત મહોત્સવ

ગ્રામીણ કૃષિ મૌસમ સેવા
કૃષિ હવામાન સેલ
કૃષિ વિજ્ઞાન વિભાગ
જૂનાગઢ કૃષિ યુનિવર્સિટી
જૂનાગઢ - ૩૬૨૦૦૧

ઈમેઈલ : - amfujnd@yahoo.com ફોન નં. ૦૨૮૫-૨૬૭૦૧૩૯



G20
ભારત ૨૦૨૩ INDIA

હવામાન આગાહી આધારીત દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારનાં **પોરબંદર જીલ્લા** માટે કૃષિ સલાહ બુલેટીન
(કૃષિ હવામાન સેલ, જૂનાગઢ કૃષિ યુનિવર્સિટી અને ભારત હવામાન વિભાગ દ્વારા સંપન્ન)

વાર:- શુક્રવાર

તારીખ: ૨૧/૦૩/૨૦૨૫

હવામાન આગાહી આધારીત બુલેટીન: **૧૦૨ (૨૦૨૪-૨૫)**

પાછલા પાંચ દિવસનું હવામાન (તારીખ:- ૧૬ થી ૨૦ માર્ચ, ૨૦૨૫)					હવામાન પરિબળો	હવામાન આગાહી આગામી પાંચ દિવસ માટે (તારીખ:- ૨૨ થી ૨૬ માર્ચ, ૨૦૨૫) માન્ય ૨૬ માર્ચ, ૨૦૨૫ : સવારના ૮:૩૦ વાગ્યાં સુધી				
૧૬/૦૩	૧૭/૦૩	૧૮/૦૩	૧૯/૦૩	૨૦/૦૩	તારીખ	૨૨/૦૩	૨૩/૦૩	૨૪/૦૩	૨૫/૦૩	૨૬/૦૩
૦	૦	૦	૦	૦	વરસાદ (મિમી)	૦	૦	૦	૦	૦
૩૨.૨	૩૫.૦	૩૫.૧	૩૬.૬	૩૫.૨	મહત્તમ તાપમાન (°સે.ગ્રે)	૩૬	૩૭	૩૬	૩૪	૩૨
૧૯.૬	૧૮.૮	૨૦.૯	૨૦.૬	૧૯.૦	લઘુત્તમ તાપમાન (°સે.ગ્રે)	૨૨	૨૨	૨૧	૧૯	૧૯
ચોખ્ખું	ચોખ્ખું	ચોખ્ખું	ચોખ્ખું	ચોખ્ખું	વાદળની સ્થિતિ	ચોખ્ખું	ચોખ્ખું	ચોખ્ખું	ચોખ્ખું	ચોખ્ખું
૯૧	૮૧	૪૨	૭૮	૭૩	સવારના ભેજના ટકા	૭૦	૭૦	૭૦	૭૫	૮૦
૬૩	૩૫	૩૯	૩૭	૫૧	બપોરબાદના ભેજના ટકા	૩૫	૩૫	૪૦	૫૦	૫૫
-	૯.૩	૧૪.૮	૧૧.૧	૧૧.૧	પવનની ગતિ (કિમી/કલાક)	૧૨	૧૧	૧૦	૯	૧૨

-	૩૨૦	૩૪૦	૩૬૦	૩૬૦	પવનની દિશા	૩૨૫	૩૨૯	૯૦	૩૪૨	૩૧૫
---	-----	-----	-----	-----	------------	-----	-----	----	-----	-----

સામાન્ય સલાહ

ભારત સરકારશ્રીના હવામાન વિભાગ તરફથી મળેલ હવામાન આગાહીની માહિતી અનુસાર અત્રેના ગ્રામીણ કૃષિ મૌસમ સેવા યોજના, જૂનાગઢ કૃષિ યુનિવર્સિટી, જૂનાગઢ તરફથી જણાવવામા આવે છેકે દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના પોરબંદર જીલ્લામાં આગામી તારીખ **૨૨/૦૩/૨૦૨૫ થી ૨૬/૦૩/૨૦૨૫** દરમિયાન આકાશ મુખ્યત્વે આગામી પાંચ દિવસોમાં મુખ્યત્વે **ચોખ્ખું** રહેશે, જીલ્લામાં આગામી પાંચ દિવસો દરમિયાન **વરસાદની** શક્યતાઓ **નહિવત છે**. પવનની ઝડપ **૦૯ થી ૧૨** કીમી/કલાક રહેવાની શક્યતા રહેશે. આ સમયગાળામાં મહત્તમ તાપમાન **૩૨ થી ૩૭** ડીગ્રી સેન્ટીગ્રેડ રહેશે તેમજ લઘુત્તમ તાપમાન **૧૯ થી ૨૨** ડીગ્રી સેન્ટીગ્રેડ રહેવાની સંભાવના છે.

- ✦ માન્ય વરસાદની સૂચી મુજબ પોરબંદર જીલ્લામાં ૨૬ ડિસેમ્બર, ૨૦૨૪ થી ૨૨ જાન્યુઆરી, ૨૦૨૫ નાં સમયગાળા દરમિયાન **હળવી સૂકી સ્થિતિ** જોવા મળેલ છે.
- ✦ સાપ્તાહિક હવામાન આગાહીની સૂચી મુજબ સૌરાષ્ટ્ર અને કચ્છનાં વિસ્તારોમાં તા. ૨૬/૦૩/૨૦૨૫ થી તા. ૦૧/૦૪/૨૦૨૫ દરમિયાન વરસાદ **નથી** & મહત્તમ તાપમાન **સામાન્ય થી નીચે** અને લઘુત્તમ તાપમાન **સામાન્ય** રહેવાની શક્યતાઓ છે.
- ✦ મેઘદૂત એપ્લીકેશન:-કૃષિ હવામાન સલાહ સેવા અને હવામાન આગાહી અંગેની જાણકારી મેળવવામાટે **ડાઉનલોડ** કરો મેઘદૂત એપ્લીકેશન કે જે ખેડૂતોને તેમની સ્થાનિક ભાષામાં સ્થાન, પાક અને પશુધન વિશે હવામાન આધારિત કૃષિ સલાહ પ્રદાન કરશે.



મેઘદૂત એપ્લીકેશનને **ડાઉનલોડ** કરવા માટે નીચેની લિંક પર ક્લિક કરો.

એન્ડ્રોઈડ(મોબાઈલ)ઉપયોગકર્તા:-<https://play.google.com/store/apps/details?id=com.aas.meghdoot>

આઈ.ઓ.એસ.ઉપયોગકર્તા:-<https://apps.apple.com/in/app/meghdoot/id1474048155>

- ❖ શિયાળું પાકોમાં, દેહધાર્મિક પરિપક્વ અવસ્થાએ કાપણીની કામગીરી કરવી.
- ❖ શિયાળુ પાકોની કાપણી થઈ ગયા બાદ આખા ખેતરનું અવલોકન કરી જમીનના નમુના લેવા અને પૃથ્થકરણ કરવા માટે લેબોરેટરીમાં મોકલી આપવા.
- ❖ ઉનાળુ પાકોના વાવેતર માટે ખેત આયોજન કરવું તે મુજબ બીયારણ, ખાતર અને દવાનો પુરવઠો મેળવી લેવો.
- ❖ ખેડૂતોને ઉનાળુ પાકોનાં વાવેતર માટે ખેતરો તૈયાર કરવા સલાહ આપવામાં આવે છે.

NOAA/VIIIRS/BLENDED NDVI Composite ending on Week no 9
(26.02.2025 to 04.03.2025) over Agricultural region
Gujarat



Agriculture vigour is good at isolated places of the state.

Source: <https://www.star.nesdis.noaa.gov>

પાકનું નામ	પાક અવસ્થા	રોગ / જીવાત / જાત	કૃષિ સલાહ
ઉનાળું મગફળી	વૃદ્ધિ / ફૂલ	ચૂસીયા જીવાત	❖ પાકને નિયમિત સમયાંતરે પિયત આપવાની કામગીરી કરવી. ❖ મગફળીમાં ચૂસીયા, મોલોમશી, થ્રીપ્સ ના નિયંત્રણ માટે ફોસ્ફામીડોન ૩ મીલી દવા અથવા ડાયમીથોએટ ૧૫ મીલી દવા ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો. મોલોમશી સાથે દાળીયા જીવાત જોવા મળે તો દાવાનો છંટકાવ મુલતવી રાખવો. મગફળીના પાકમાં આંતરખેડ કરવી અને હાથથી નિદામણ કરવું.
ઉનાળું મગ	ઉગાવો / વૃદ્ધિ	ચૂસીયા જીવાત	❖ પાકને સમયાંતરે પિયત આપવાની કામગીરી કરવી. ❖ ચૂસીયા જીવાતોના નિયંત્રણ માટે લીમડાનું તેલ @૫ મિલી/લીટર મુજબ છાંટવું. - પાકમાં આંતરખેડ કરવી અને હાથથી નિદામણની કામગીરી કરવી.
ઉનાળું અડદ	ઉગાવો / વૃદ્ધિ	ચૂસીયા જીવાત	❖ પાકને સમયાંતરે પિયત આપવાની કામગીરી કરવી. ❖ ચૂસીયા જીવાતોના નિયંત્રણ માટે લીમડાનું તેલ @૫ મિલી/લીટર મુજબ છાંટવું. - પાકમાં આંતરખેડ કરવી અને હાથથી નિદામણની કામગીરી કરવી.
ઉનાળું તલ	ઉગાવો / વૃદ્ધિ	તડતડિયા / સફેદમાખી	❖ પાકને સમયાંતરે પિયત આપવાની કામગીરી કરવી. - પાકમાં તડતડિયા અને સફેદમાખીનાં નિયંત્રણ માટે ઓક્ઝીડીમેટોન - મિથાઈલ ૨૫ ઇ.સી.નો છંટકાવ કરવો.

			પાકમાં આંતરખેડ કરવી અને હાથથી નિદામણની કામગીરી કરવી.
આંબા	ફૂલ / ફળ અવસ્થા	મધીયા & ભુકીછારા	❖ આંબામાં મધીયાના નિયંત્રણમાટે ૫૦ % કાર્બારીલ ૪૦ ગ્રામ પાવડર અથવા ૩૬ % ઈ.સી. મોનોક્રોટોફોસ ૧૦ મીલી દવા ૧૦ લી.પાણીમાં ભેળવી છંટકાવ કરવો. ❖ આંબામાં ભુકીછારા ના નિયંત્રણ માટે ૧૦ લી. પાણીમાં ૩૦ ગ્રામ દ્રાવ્ય ગંધક + કેરેથોન ૪૮ % ઈ.સી. ૧૦ મીલી દવાનો છંટકાવ કરવો. ❖ આંબાને નિયમિત સમયાંતરે પિયત આપવાની કામગીરી કરવી. કેરી સોપારી જેવડા કદની થાય એટલે ૨૦ પી.પી.એમ. એન.એ.એ. અને ૨% ચુરીયા ના દ્રાવણનો છંટકાવ કરવાથી કેરી ખરતી અટકે છે અને ફળના કદમાં વધારો થાય છે.
શાકભાજી	વૃદ્ધી/ફુલ/ ફળ	-	❖ શાકભાજીના પાકોને નીદામણ મુક્ત કરવું. શાકભાજી ને સમયસર ઉતારી ગ્રેડીંગ કરી બજારમાં મુકવા. ❖ પાકને નિયમિત સમયાંતરે પિયત આપવાની કામગીરી કરવી. આગાહી સમયગાળા દરમિયાન ખેતર પરનાં બધાં ખેતકાર્યો કરવા.
પશુપાલન	-	-	❖ પશુઓને યોગ્ય આહાર આપવો & પીવા માટે ચોખ્ખું પાણી આપવું અને પશુઓના રહેઠાણ ચોખ્ખા રાખવા. ❖ માખી મચ્છરથી રક્ષણ માટે ફીનાઈલ છાટવું. ❖ પશુઓને યોગ્ય આહારમાંખનિજયુક્ત મિશ્રણ સાથે સંગ્રહિત ચારા આપો. ચારા ને સલામત સ્થળે સંગ્રહિત કરવો જોઈએ. ❖ ચારા માટે ચાફ કટર વાપરો. ❖ પશુઓમાં ઈંતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટાઝ ૨ મીલી/ ૧ લીટર પાણીમાં નાખીને છાટવી. દુધાળા પશુઓના આઉને ઝીંક ઓક્સાઈડ અથવા બોરીક પાવડરથી બરાબર સાફ કરવા.



75
આઝાદી કા
અમૃત મહોત્સવ

ગ્રામીણ કૃષિ મૌસમ સેવા
કૃષિ હવામાન સેલ
કૃષિ વિજ્ઞાન વિભાગ
જૂનાગઢ કૃષિ યુનિવર્સિટી
જૂનાગઢ - ૩૬૨૦૦૧

ઈમેઈલ : - amfujnd@yhaoo.com ફોન નં. ૦૨૮૫-૨૬૭૦૧૩૯



G20
ભારત 2023

હવામાન આગાહી આધારીત દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારનાં ગીર સોમનાથ જીલ્લા માટે કૃષિ સલાહ બુલેટીન
(કૃષિ હવામાન સેલ, જૂનાગઢ કૃષિ યુનિવર્સિટી અને ભારત હવામાન વિભાગ દ્વારા સંપન્ન)

વાર:- શુક્રવાર

તારીખ:-૨૧/૦૩/૨૦૨૫

હવામાન આગાહી આધારીત બુલેટીન: ૧૦૨ (૨૦૨૪-૨૫)

પાછલા પાંચ દિવસનું હવામાન (તારીખ:- ૧૬ થી ૨૦ માર્ચ, ૨૦૨૫)					હવામાન પરિબલો	હવામાન આગાહી આગામી પાંચ દિવસ માટે (તારીખ:- ૨૨ થી ૨૬ માર્ચ, ૨૦૨૫) માન્ય ૨૬ માર્ચ, ૨૦૨૫ સવારના ૮:૩૦ વાગ્યાં સુધી				
૧૬/૦૩	૧૭/૦૩	૧૮/૦૩	૧૯/૦૩	૨૦/૦૩	તારીખ	૨૨/૦૩	૨૩/૦૩	૨૪/૦૩	૨૫/૦૩	૨૬/૦૩
૦	૦	૦	૦	૦	વરસાદ (મિમી)	૦	૦	૦	૦	૦
૩૧.૮	૩૩.૯	૩૩.૯	૩૫.૬	૩૦.૪	મહત્તમ તાપમાન (°સે.ગ્રે)	૩૪	૩૪	૩૩	૩૨	૩૧
૨૨.૨	૨૧.૨	૨૦.૩	૨૧.૩	૨૧.૯	લઘુતમ તાપમાન (°સે.ગ્રે)	૨૪	૨૪	૨૨	૨૧	૨૧
ચોખ્ખું	ચોખ્ખું	ચોખ્ખું	ચોખ્ખું	ચોખ્ખું	વાદળની સ્થિતિ	ચોખ્ખું	ચોખ્ખું	ચોખ્ખું	ચોખ્ખું	ચોખ્ખું
૮૦	૭૮	૭૪	૭૧	૬૯	સવારના ભેજના ટકા	૮૫	૮૫	૮૦	૮૦	૮૦
૭૪	૬૭	૪૯	૪૯	૪૦	બપોરબાદના ભેજના ટકા	૬૦	૬૦	૭૦	૭૦	૭૦
૩.૭	૫.૬	-	-	૫.૬	પવનની ગતિ(કિમી/કલાક)	૧૨	૧૨	૯	૧૦	૧૫
૯૯	૨૦	-	-	૨૦	પવનની દિશા (ડિગ્રી)	૩૨૯	૩૪૨	૭	૩૪૬	૩૨૫

સામાન્ય સલાહ

ભારત સરકારશ્રી ના હવામાન વિભાગ તરફથી મળેલ હવામાન આગાહીની માહિતી અનુસાર અત્રેના ગ્રામીણ કૃષિ મૌસમ સેવા યોજના, જૂનાગઢ કૃષિ યુનિવર્સિટી, જૂનાગઢ તરફથી જણાવવામા આવે છે કે દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના ગીર સોમનાથ જીલ્લામાં આગામી તારીખ ૨૨/૦૩/૨૦૨૫ થી ૨૬/૦૩/૨૦૨૫ દરમિયાન આકાશ મુખ્યત્વે આગામી દિવસોમાં મુખ્યત્વે ચોખ્ખું રહેશે, જીલ્લામાં આગામી પાંચ દિવસો દરમિયાન વરસાદની શક્યતાઓ નહિવત છે. પવનની ઝડપ ૦૯ થી ૧૫ કીમી/કલાક રહેવાની શક્યતાઓ રહેશે. આ સમયગાળામાં મહત્તમ તાપમાન ૩૧ થી ૩૪ ડીગ્રી સેન્ટીગ્રેડ રહેશે તેમજ લઘુતમ તાપમાન ૨૧ થી ૨૪ ડીગ્રી સેન્ટીગ્રેડ રહેવાની સંભાવના છે.

✦ માન્ય વરસાદની સૂચી મુજબ ગીર સોમનાથ જિલ્લામાં ૨૬ ડિસેમ્બર, ૨૦૨૪ થી ૨૨ જાન્યુઆરી, ૨૦૨૫ નાં સમયગાળા દરમિયાન હળવી સૂકી સ્થિતિ જોવા મળેલ છે.

✦ સાપ્તાહિક હવામાન આગાહીની સૂચી મુજબ સૌરાષ્ટ્ર અને કચ્છનાં વિસ્તારોમાં તા. ૨૬/૦૩/૨૦૨૫ થી તા. ૦૧/૦૪/૨૦૨૫ દરમિયાન વરસાદ નથી & મહત્તમ તાપમાન સામાન્ય થી નીચે અને લઘુત્તમ તાપમાન સામાન્ય રહેવાની શક્યતાઓ છે.

✦ મેઘદૂત એપ્લિકેશન:-કૃષિ હવામાન સલાહ સેવા અને હવામાન આગાહી અંગેની જાણકારી મેળવવા માટે ડાઉનલોડ કરો મેઘદૂત એપ્લિકેશન કે જે ખેડૂતોને તેમની સ્થાનિક ભાષામાં સ્થાન, પાક અને પશુધન વિશે હવામાન આધારિત કૃષિ સલાહ પ્રદાન કરશે.

મેઘદૂત એપ્લિકેશનને ડાઉનલોડ કરવા માટે નીચેની લિંક પર ક્લિક કરો.

એન્ડ્રોઈડ(મોબાઈલ)ઉપયોગકર્તા:-<https://play.google.com/store/apps/details?id=com.aas.meghdoot>


આઈ.ઓ.એસ.ઉપયોગકર્તા:-<https://apps.apple.com/in/app/meghdoot/id1474048155>

❖ શિયાળું પાકોમાં, દેહધાર્મિક પરિપક્વ અવસ્થાએ કાપણીની કામગીરી કરવી.

❖ શિયાળું પાકોની કાપણી થઈ ગયા બાદ આખા ખેતરનું અવલોકન કરી જમીનના નમુના લેવા અને પૃથ્થકરણ કરવા માટે લેબોરેટરીમાં મોકલી આપવા.

❖ ઉનાળું પાકોના વાવેતર માટે ખેત આયોજન કરવું તે મુજબ બીયારણ, ખાતર અને દવાનો પુરવઠો મેળવી લેવો.

❖ ખેડૂતોને ઉનાળું પાકોનાં વાવેતર માટે ખેતરો તૈયાર કરવા સલાહ આપવામાં આવે છે.



પાકનું નામ	પાક અવસ્થા	રોગ / જીવાત / જાત	કૃષિ સલાહ
ઉનાળું મગફળી	વૃદ્ધિ / ફૂલ	યૂસીયા જીવાત	❖ પાકને નિયમિત સમયાંતરે પિયત આપવાની કામગીરી કરવી. ❖ મગફળીમાં યુસીયા, મોલોમશી, થ્રીપ્સ ના નિયંત્રણ માટે

			ફોસ્ફોર્મીડોન ૩ મીલી દવા અથવા ડાયમીથોએટ ૧૫ મીલી દવા ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો. મોલોમશી સાથે દાળીયા જીવાત જોવા મળે તો દાવાનો છંટકાવ મુલતવી રાખવો. મગફળીના પાકમાં આંતરખેડ કરવી અને હાથથી નિદામણ કરવું.
ઉનાળું મગ	ઉગાવો / વૃદ્ધિ	ચૂસીયા જીવાત	❖ પાકને સમયાંતરે પિયત આપવાની કામગીરી કરવી. ❖ ચૂસીયા જીવાતોના નિયંત્રણ માટે લીમડાનું તેલ @૫ મિલી/ લીટર મુજબ છાંટવું. પાકમાં આંતરખેડ કરવી અને હાથથી નિદામણની કામગીરી કરવી.
ઉનાળું અડદ	ઉગાવો / વૃદ્ધિ	ચૂસીયા જીવાત	❖ પાકને સમયાંતરે પિયત આપવાની કામગીરી કરવી. ❖ ચૂસીયા જીવાતોના નિયંત્રણ માટે લીમડાનું તેલ @૫ મિલી/ લીટર મુજબ છાંટવું. પાકમાં આંતરખેડ કરવી અને હાથથી નિદામણની કામગીરી કરવી.
ઉનાળું તલ	ઉગાવો / વૃદ્ધિ	તડતડિયા / સફેદમાખી	❖ પાકને સમયાંતરે પિયત આપવાની કામગીરી કરવી. પાકમાં તડતડિયા અને સફેદમાખીનાં નિયંત્રણ માટે ઓક્ઝીડીમેટોન - મિથાઈલ ૨૫ ઇ.સી.નો છંટકાવ કરવો. પાકમાં આંતરખેડ કરવી અને હાથથી નિદામણની કામગીરી કરવી.
આંબા	ફળ અવસ્થા	મધીયા & ભુકીછારા	❖ આંબામાં મધીયાના નિયંત્રણ માટે ૫૦ % કાર્બારીલ ૪૦ ગ્રામ પાવડર અથવા ૩૬ % ઇ.સી. મોનોક્રોટોફોસ ૧૦ મીલી દવા ૧૦ લી.પાણીમાં ભેળવી છંટકાવ કરવો. ❖ આંબામાં ભુકીછારા ના નિયંત્રણ માટે ૧૦ લી. પાણીમાં ૩૦ ગ્રામ દ્રાવ્ય ગંધક + કેરેથોન ૪૮ % ઇ.સી. ૧૦ મીલી દવાનો છંટકાવ કરવો. ❖ આંબાને નિયમિત સમયાંતરે પિયત આપવાની કામગીરી કરવી. કેરી સોપારી જેવડા કદની થાય એટલે ૨૦ પી.પી.એમ. એન.એ.એ. અને ૨% ચુરીયા ના દ્રાવણનો છંટકાવ કરવાથી કેરી ખરતી અટકે છે અને ફળના કદમાં વધારો થાય છે.
શાકભાજી	વૃદ્ધી/ફુલ/ ફળ	-	❖ શાકભાજીના પાકોને નીદામણ મુક્ત કરવું. શાકભાજી ને સમયસર ઉતારી ગ્રેડીંગ કરી બજારમાં મુકવા. ❖ પાકને નિયમિત સમયાંતરે પિયત આપવાની કામગીરી કરવી. આગાહી સમયગાળા દરમિયાન ખેતર પરનાં બધાં ખેતકાર્યો કરવા.

પશુપાલન	-	-	❖ પશુઓને યોગ્ય આહાર આપવો & પીવા માટે ચોખ્ખું પાણી આપવું અને પશુઓના રહેઠાણ ચોખ્ખા રાખવા. ❖ માખી મચ્છરથી રક્ષણ માટે ફીનાઈલ છાટવું. ❖ પશુઓને યોગ્ય આહારમાંખનિજયુક્ત મિશ્રણ સાથે સંગ્રહિત ચારા આપો. ચારા ને સલામત સ્થળે સંગ્રહિત કરવો જોઈએ. ❖ ચારા માટે ચાફ કટર વાપરો. ❖ પશુઓમાં ઇતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટાઝ ૨ મીલી/ ૧ લીટર પાણીમાં નાખીને છાટવી. દુધાળા પશુઓના આઉને ઝીંક ઓક્સાઈડ અથવા બોરીક પાવડરથી બરાબર સાફ કરવા.
---------	---	---	---

યાંત્રિક અધિકારી
ગ્રામીણ કૃષિ મૌસમ સેવા
જી.કૃ.યુ.

AMFU, JAU, Junagadh



Gramin Krushi Mausam Seva (GKMS)
Agrometeorological Cell
Department of Agronomy
Junagadh Agricultural University
Junagadh-362001

E-mail: amfujnd@yahoo.com Phone: 0285-2670139



Agromet Advisory Service Bulletin for Junagadh District of South Saurashtra Agro-Climatic Zone
(Issued jointly by Junagadh Agricultural University and India Meteorological Department)

Day:- Friday

Date: 21-03-2025

Agromet Advisory Service bulletin. No:- 102 (2024-25)

Observed Weather during last Five days (Dated 16 th to 20 th March, 2025)					Weather Parameters	Weather Forecast upcoming Five days (Valid for 22 nd to 26 th March, 2025) until 08:30 hrs of 26-03-2025				
16/03	17/03	18/03	19/03	20/03	Date	22/03	23/03	24/03	25/03	26/03
0	0	0	0	0	Rainfall(mm)	0	0	0	0	0
37.6	36.5	36.0	37.1	37.0	T _{max} (°C)	39	40	40	40	38
21.0	21.3	23.8	21.5	18.1	T _{min} (°C)	22	23	21	19	19
0	0	0	0	0	Cloud Cover	0	0	0	0	0
57	54	39	45	51	RH-I (%)	60	60	60	65	70
20	14	17	11	17	RH-II (%)	20	15	15	20	30
5.5	6.7	6.8	5.9	4.9	Wind speed (km/hr)	7	6	7	6	9
320	360	290	320	290	Wind direction	326	347	18	357	318

Agro meteorological Advisories

General Advice	Rain fall	As per IMD Forecast, There is forecasting of no rainfall occurs during next five days. Sky will be remaining clear.
	Temperature	Maximum and minimum temperature likely to be in range of 38 - 40°C and 19 - 23°C , respectively.
	RH (%)	The Relative Humidity is likely to be in the range of 15 - 70 % .
	Wind Direction and Wind Speed	Direction and speed of wind is to be NW & N and around 06 to 09 km/h , respectively.

✚ **Standardized Precipitation Index:** - For the period of 26th December, 2024 to 22nd January, 2025. Junagadh district considered under the mildly dry condition.

✚ As per ERFs products, no rainfall and T_{max} below normal & T_{min} normal may occur in Saurashtra & Kutch region during 26th March to 01st April, 2025.

✚ **MEGHDOOT Mobile Application:-** Download the MEGHDOOT mobile application for Agromet Advisory Services and information on weather forecasting, which will provide weather based agricultural advice to farmers for their location, crops and livestock in their local language.

Click on the following link to download the MEGHDOOT application:

Android users: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>


IOS users: <https://apps.apple.com/in/app/meghdoot/id1474048155>

✚ In *rabi* crops, harvesting operation should be carry out at physiological maturity stage.

✚ After harvesting of rabi crops, take the soil samples and send to the soil testing laboratory for the soil analysis.

✚ To get the supply of seeds, fertilizers and insecticide, pesticide and fungicide according to planning the farm for summer season.

✚ Farmers are advised to prepare their field for summer season crops sowing.



Crop	Variety/ Pest/Insect	Stage	Agro met Advisories
Summer Groundnut	Sucking pest	Vegetative / Flowering	➤ Apply irrigation to the crop at regular an interval. ➤ For the control of sucking pest (aphid, thrips jassid), spray 3 ml Phosphamidon OR 15 ml

			Dimethoate in 10 litre of water. For ladybird beetle is observed with aphid then postpone the spraying operation. ➤ Inter culturing and hand weeding should be carried out in groundnut.
Summer Green gram	Sucking pest	Germination / Vegetative	➤ Apply irrigation to the crop at regular an interval. To spray neem oil @ 5 ml/l for the control of sucking pest. ➤ Inter culturing and hand weeding should be carried out in crop.
Summer Black gram	Sucking pest	Germination / Vegetative	➤ Apply irrigation to the crop at regular an interval. ➤ To spray neem oil @ 5 ml/l for the control of sucking pest. ➤ Inter culturing and hand weeding should be carried out in crop.
Summer Sesame	Leafhopper / Whitefly	Germination / Vegetative	➤ Apply irrigation to the crop at regular an interval. To control of leafhopper and whitefly, Spray Oxydemeton-methyl 25 EC. ➤ Inter culturing and hand weeding should be carried out in crop.
Mango	Madhiya & Powdery mildew	Fruiting	➤ For the control of madhiya, spray 50 % WP carbaryl 40 gm OR 36 % EC monocrotophos 10 ml in 10 liter of water. ➤ For the control of powdery mildew, spray 30 gm sulphur + 48 % EC carethon 10 ml in 10 liter of water. ➤ Apply irrigation to the mango at regular an

			interval. ➤ To Spray 20 PPM NAA and 2 % urea solution for the control of stops fruit break and increase the fruit size.
Vegetable crops	-	Sowing/ Flowering/ Fruiting	➤ Keep vegetable crop weed free, apply irrigation regularly and send vegetable fruit to the market often grading. ➤ Apply irrigation to the crop at regular an interval. ➤ To done all the activity of farm during forecast period.
Livestock	-	-	➤ To provide adequate food & clean water for drinking and keep the animals clean shed. ➤ Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Give stored fodder with mineral mixture. Fodder should be stored at safe place. ➤ Feed silage & hay material along with concentrate feed. ➤ Use chaff cutter for fodder. ➤ For the control of tick and other parasites, spray deltemethrin or aemitaz 2 ml in 1 litre of water. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.




आज़ादी का
अमृत महोत्सव

Gramin Krushi Mausam Seva (GKMS)
Agrometeorological Cell
Department of Agronomy
Junagadh Agricultural University
Junagadh-362001

E-mail: amfujnd@yahoo.com Phone: 0285-2670139




Agromet Advisory Service Bulletin for Porbandar District of South Saurashtra Agro-Climatic Zone
(Issued jointly by Junagadh Agricultural University and India Meteorological Department)

Day:- Friday

Date: 21-03-2025

Agromet Advisory Service bulletin. No:- 102 (2024-25)

Observed Weather during last Five days (Dated 16 th to 20 th March,2025)					Weather Parameters	Weather Forecast upcoming Five days (Valid for 22 nd to 26 th March,2025) until 08:30 hrs of 26-03-2025				
16/03	17/03	18/03	19/03	20/03	Date	22/03	23/03	24/03	25/03	26/03
0	0	0	0	0	Rainfall (mm)	0	0	0	0	0
32.2	35.0	35.1	36.6	35.2	Tmax (°C)	36	37	36	34	32
19.6	18.8	20.9	20.6	19.0	Tmin (°C)	22	22	21	19	19
0	0	0	0	0	Cloud Cover	0	0	0	0	0
91	81	42	78	73	RH-I (%)	70	70	70	75	80
63	35	39	37	51	RH-II (%)	35	35	40	50	55
-	9.3	14.8	11.1	11.1	Wind speed (km/hr)	12	11	10	9	12
-	320	340	360	360	Wind direction	325	329	90	342	315

Agro meteorological Advisories

General Advice	Rain fall	As per IMD Forecast, There is forecasting of no rainfall occurs during next five days. Sky will be remaining clear.
	Temperature	Maximum and minimum temperature likely to be in range of 32 - 37°C and 19 - 22°C , respectively.
	RH (%)	The Relative Humidity is likely to be in the range of 35 – 80 % .
	Wind Direction and Wind Speed	Direction and speed of wind is to be NW & N and 09 - 12 km/h , respectively.

- ❖ **Standardized Precipitation Index:** - For the period **26th December, 2024 to 22nd January, 2025**. Porbandar district considered under the mildly dry condition.
- ❖ **As per ERFs products**, **no** rainfall and T_{max} **below normal** & T_{min} **normal** may occur in Saurashtra & Kutch region during **26th March to 01st April, 2025**.


- ❖ **MEGHDOOT Mobile Application:-Download** the MEGHDOOT mobile application for Agromet Advisory Services and information on weather forecasting, which will provide weather based agricultural advice to farmers for their location, crops and livestock in their local language.

Click on the following link to download the MEGHDOOT application:

Android users:<https://play.google.com/store/apps/details?id=com.aas.meghdoot>

IOS users:<https://apps.apple.com/in/app/meghdoot/id1474048155>

- ✚ In *rabi* crops, harvesting operation should be carry out at physiological maturity stage.
- ✚ After harvesting of rabi crops, take the soil samples and send to the soil testing laboratory for the soil analysis.
- ✚ To get the supply of seeds, fertilizers and insecticide, pesticide and fungicide according to planning the farm for summer season.
- ✚ Farmers are advised to prepare their field for summer season crops sowing.



Crop	Variety/ Pest / Insect	Stage	Agro met Advisories
Summer Groundnut	Sucking pest	Germination / Vegetative	➤ Apply irrigation to the crop at regular an interval. ➤ For the control of sucking pest (aphid, thrips jassid), spray 3 ml Phosphamidon

			For ladybird beetle is observed with aphid then postpone the spraying operation. ➤ Interculturing and hand weeding should be carried out in groundnut.
Summer Green gram	Sucking pest	Germination / Vegetative	➤ Apply irrigation to the crop at regular an interval. To spray neem oil @ 5 ml/l for the control of sucking pest. ➤ Inter culturing and hand weeding should be carried out in crop.
Summer Black gram	Sucking pest	Germination / Vegetative	➤ Apply irrigation to the crop at regular an interval. ➤ To spray neem oil @ 5 ml/l for the control of sucking pest. ➤ Inter culturing and hand weeding should be carried out in crop.
Summer Sesame	Leafhopper / Whitefly	Germination / Vegetative	➤ Apply irrigation to the crop at regular an interval. To control of leafhopper and whitefly, Spray Oxydemeton-methyl 25 EC. ➤ Inter culturing and hand weeding should be carried out in crop.
Mango	Madhiya & Powdery mildew	Fruiting	➤ For the control of madhiya, spray 50 % WP carbaryl 40 gm OR 36 % EC monocrotophos 10 ml in 10 liter of water. ➤ For the control of powdery mildew, spray 30 gm sulphur + 48 % EC carethon 10 ml in 10 liter of water. ➤ Apply irrigation to the mango at regular an interval.

			➤ To Spray 20 PPM NAA and 2 % urea solution for the control of stops fruit break and increase the fruit size.
Vegetable crops	-	Sowing/ Flowering/ Fruiting	➤ Keep vegetable crop weed free, apply irrigation regularly and send vegetable fruit to the market often grading. ➤ Apply irrigation to the crop at regular an interval. ➤ To done all the activity of farm during forecast period.
Livestock	-	-	➤ To provide adequate food & clean water for drinking and keep the animals clean shed. ➤ Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Give stored fodder with mineral mixture. Fodder should be stored at safe place. ➤ Feed silage & hay material along with concentrate feed. ➤ Use chaff cutter for fodder. ➤ For the control of tick and other parasites, spray deltemethrin or aemitaz 2 ml in 1 litre of water. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.



75
आज़ादी का
अमृत महोत्सव

Gramin Krushi Mausam Seva (GKMS)
Agrometeorological Cell
Department of Agronomy
Junagadh Agricultural University
Junagadh-362001

E-mail: amfujnd@yahoo.com Phone: 0285-2670139



Agromet Advisory Service Bulletin for Gir Somnath District of South Saurashtra Agro-Climatic Zone
(Issued jointly by Junagadh Agricultural University and India Meteorological Department)

Day:- Friday
Date: 21-03-2025

Agromet Advisory Service Bulletin. No:-102 (2024-25)

Observed Weather during last Five days (Dated 16 th to 20 th March,2025)					Weather Parameters	Weather Forecast upcoming Five days (Valid for 22 nd to 26 th March,2025) until 08:30 hrs of 26-03-2025				
16/03	17/03	18/03	19/03	20/03	Date	22/03	23/03	24/03	25/03	26/03
0	0	0	0	0	Rainfall (mm)	0	0	0	0	0
31.8	33.9	33.9	35.6	30.4	Tmax (°C)	34	34	33	32	31
22.2	21.2	20.3	21.3	21.9	Tmin (°C)	24	24	22	21	21
1	4	0	0	0	Cloud cover	0	0	0	0	0
80	78	74	71	69	RH-I (%)	85	85	90	90	90
74	67	49	49	40	RH-II (%)	60	60	70	70	70
3.7	5.6	-	-	5.6	Wind speed (km/hr)	12	12	9	10	15
99	20	-	-	20	Wind direction	329	342	7	346	325

Agro meteorological Advisories

General Advice	Rain fall	As per IMD Forecast, There is forecasting of no rainfall occurs during next five days. Sky will be remaining clear.
	Temperature	Maximum and minimum temperature likely to be in range of 31 - 34°C and 21 - 24°C , respectively.
	RH (%)	The Relative Humidity is likely to be in the range of 60 – 90 % .
	Wind Direction and Wind Speed	Direction and speed of wind is to be mainly NW & N and around 09 – 15 km/h , respectively.

Standardized Precipitation Index: - For the period **26th December, 2024 to 22nd January,**

2025. Gir Somnath district considered under the mildly dry condition.

✚ As per ERFs products, no rainfall and $T_{\max}$ below normal & $T_{\min}$ normal may occur in Saurashtra & Kutch region during 26th March to 01st April, 2025.

✚ **MEGHDOOT Mobile Application:-**Download the MEGHDOOT mobile application for Agromet Advisory Services and information on weather forecasting, which will provide weather based agricultural advice to farmers for their location, crops and livestock in their local language.

Click on the following link to download the MEGHDOOT application:

Android users: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>


IOS users: <https://apps.apple.com/in/app/meghdoot/id1474048155>

✚ In rabi crops, harvesting operation should be carry out at physiological maturity stage.

✚ After harvesting of rabi crops, take the soil samples and send to the soil testing laboratory for the soil analysis.

✚ To get the supply of seeds, fertilizers and insecticide, pesticide and fungicide according to planning the farm for the next summer season.

✚ Farmers are advised to prepare their field for an upcoming summer season crops sowing.



Crop	Variety/ Pest / Insect	Stage	Agro met Advisories
Summer Groundnut	Sucking pest	Vegetative / Flowering	➤ Apply irrigation to the crop at regular an interval. ➤ For the control of sucking pest (aphid, thrips jassid), spray 3 ml Phosphamidon OR 15 ml Dimethoate in 10 litre of water. For ladybird

			the spraying operation. ➤ Interculturing and hand weeding should be carried out in groundnut.
Summer Green gram	Sucking pest	Germination / Vegetative	➤ Apply irrigation to the crop at regular an interval. To spray neem oil @ 5 ml/l for the control of sucking pest. ➤ Inter culturing and hand weeding should be carried out in crop.
Summer Black gram	Sucking pest	Germination / Vegetative	➤ Apply irrigation to the crop at regular an interval. ➤ To spray neem oil @ 5 ml/l for the control of sucking pest. ➤ Inter culturing and hand weeding should be carried out in crop.
Summer Sesame	Leafhopper / Whitefly	Germination / Vegetative	➤ Apply irrigation to the crop at regular an interval. To control of leafhopper and whitefly, Spray Oxydemeton-methyl 25 EC. ➤ Inter culturing and hand weeding should be carried out in crop.
Mango	Madhiya & Powdery mildew	Fruiting	➤ For the control of madhiya, spray 50 % WP carbaryl 40 gm OR 36 % EC monocrotophos 10 ml in 10 liter of water. ➤ For the control of powdery mildew, spray 30 gm sulphur + 48 % EC carethon 10 ml in 10 liter of water. ➤ Apply irrigation to the mango at regular an interval.

			➤ To Spray 20 PPM NAA and 2 % urea solution for the control of stops fruit break and increase the fruit size.
Vegetable crops	-	Sowing/ Flowering/ Fruiting	➤ Keep vegetable crop weed free, apply irrigation regularly and send vegetable fruit to the market often grading. ➤ Apply irrigation to the crop at regular an interval. ➤ To done all the activity of farm during forecast period.
Livestock	-	-	➤ To provide adequate food & clean water for drinking and keep the animals clean shed. ➤ Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Give stored fodder with mineral mixture. Fodder should be stored at safe place. ➤ Feed silage & hay material along with concentrate feed. ➤ Use chaff cutter for fodder. ➤ For the control of tick and other parasites, spray deltemethrin or aemitaz 2 ml in 1 litre of water. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.

Technical Officer
Gramin Krishi Mausam Seva (GKMS)
J.A.U., Junagadh