

હવામાન આગાહી આધારીત કૃષિ સલાહ બુલેટીન દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ

તા. ૨૫-૦૫-૨૦૧૯ થી તા. ૨૯-૦૫-૨૦૧૯

(ગ્રામીણ કૃષિ મૌસમ સેવા, જુનાગઢ કૃષિ યુનિવર્સિટી, જુ.કૃ.યુ.)

હવામાન આગાહી આધારીત બુલેટીન : ૨૦૧૯-૨૦

પાછલા અઠવાડીયાનું જુનાગઢનું હવામાન

હવામાન પરીબળો	દક્ષિણ સૌરાષ્ટ્ર કૃષિ આબોહવાકીય વિસ્તાર
વરસાદ (મીમી)	૦૦૦.૦
મહત્તમ તાપમાન (°સે.)	૪૦.૯
લઘુતમ તાપમાન (°સે.)	૨૬.૦
ભેજ(%) (સવાર/બપોરબાદ)	૮૦ - ૩૨
પવનની ગતિ (કિમી/કલાક)	૭.૫

પત્રક-બ : જુલ્લાવાર હવામાન આગાહી (તા.૨૯-૦૫-૨૦૧૯ સુધી)

ભારત સરકારશ્રીના હવામાન વિભાગ તરફથી મળેલ હવામાન આગાહીની માહિતી અનુસાર અત્રેના ગ્રામીણ કૃષિ મૌસમ સેવા યોજના, જુનાગઢ કૃષિ યુનિવર્સિટી, જુનાગઢ તરફથી જણાવવામા આવે છે કે દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના

૧.જુનાગઢ:- જુલ્લામા આગામી તારીખ ૨૫/૦૫/૨૦૧૯ થી ૨૯/૦૫/૨૦૧૯ દરમ્યાન આકાશ મુખ્યત્વે આગામી પાંચ દિવસોમાં ચોખ્ખું થી મોટેભાગે ચોખ્ખું રહેશે, આવતા પાંચ દિવસોમાં વરસાદ ની કોઈ શક્યતાઓ રહેલી નથી. પવનની ઝડપ ૧૪ થી ૧૬ કીમી/કલાક રહેવાની શક્યતા રહેશે. આ સમયગાળામાં મહત્તમ તાપમાન ૩૯ થી ૪૦ ડીગ્રી સેન્ટીગ્રેડ રહેશે તેમજ લઘુતમ તાપમાન ૨૭ ડીગ્રી સેન્ટીગ્રેડ રહેવાની સંભાવના છે.

૨.પોરબંદર:- જુલ્લામા આગામી તારીખ ૨૫/૦૫/૨૦૧૯ થી ૨૯/૦૫/૨૦૧૯ દરમ્યાન આકાશ મુખ્યત્વે આગામી પાંચ દિવસોમાં ચોખ્ખું રહેશે, આવતા પાંચ દિવસોમાં વરસાદ ની કોઈ શક્યતાઓ રહેલી નથી. પવનની ઝડપ ૧૪ થી ૧૮ કીમી/કલાક રહેવાની શક્યતા રહેશે. આ સમયગાળામાં મહત્તમ તાપમાન ૩૬ થી ૩૭ ડીગ્રી સેન્ટીગ્રેડ રહેશે તેમજ લઘુતમ તાપમાન ૨૭ ડીગ્રી સેન્ટીગ્રેડ રહેવાની સંભાવના છે.

૨.ગીર સોમનાથ:- જીલ્લામા આગામી તારીખ ૨૫/૦૫/૨૦૧૯ થી ૨૯/૦૫/૨૦૧૯ દરમ્યાન આકાશ મુખ્યત્વે આગામી પાંચ દિવસોમાં ચોખ્ખું થી મોટેભાગે ચોખ્ખું રહેશે, આવતા પાંચ દિવસોમાં વરસાદ ની કોઈ શક્યતાઓ રહેલી નથી. પવનની ઝડપ ૧૫ થી ૧૮ કીમી/કલાક રહેવાની શક્યતા રહેશે. આ સમયગાળામાં મહત્તમ તાપમાન ૩૫ થી ૩૭ ડીગ્રી સેન્ટીગ્રેડ રહેશે તેમજ લઘુત્તમ તાપમાન ૨૭ થી ૨૮ ડીગ્રી સેન્ટીગ્રેડ રહેવાની સંભાવના છે.

જુનાગઢ

હવામાન પરીબળો / તારીખ	દિવસ-૧ ૨૫-૦૫-૨૦૧૯	દિવસ-૨ ૨૬-૦૫-૨૦૧૯	દિવસ-૩ ૨૭-૦૫-૨૦૧૯	દિવસ-૪ ૨૮-૦૫-૨૦૧૯	દિવસ-૫ ૨૯-૦૫-૨૦૧૯
વરસાદ (મી.મી)	૦૦	૦૦	૦૦	૦૦	૦૦
મહત્તમ તાપમાન (સે.ગ્રે)	૩૯	૩૯	૩૯	૩૯	૪૦
લઘુત્તમ તાપમાન (સે.ગ્રે)	૨૭	૨૭	૨૭	૨૭	૨૭
વાદળની સ્થિતિ	ચોખ્ખું	ચોખ્ખું	ચોખ્ખું	મોટેભાગે ચોખ્ખું	ચોખ્ખું
સવારના ભેજના ટકા	૮૭	૮૮	૯૦	૮૯	૮૭
બપોર બાદ ભેજના ટકા	૪૨	૪૦	૪૩	૪૭	૪૫
પવનની ગતિ (કિમી/કલાક)	૦૧૫	૦૧૬	૦૧૪	૦૧૫	૦૧૫
પવનની દિશા (ડીગ્રી)	૨૭૯	૨૬૯	૨૬૪	૨૭૬	૨૮૮

પોરબંદર

વરસાદ (મી.મી)	૦૦	૦૦	૦૦	૦૦	૦૦
મહત્તમ તાપમાન (સે.ગ્રે)	૩૭	૩૬	૩૭	૩૬	૩૭
લઘુત્તમ તાપમાન (સે.ગ્રે)	૨૭	૨૭	૨૭	૨૭	૨૭
વાદળની સ્થિતિ	ચોખ્ખું	ચોખ્ખું	ચોખ્ખું	ચોખ્ખું	ચોખ્ખું
સવારના ભેજના ટકા	૮૭	૮૭	૮૮	૮૭	૮૭
બપોર બાદ ભેજના ટકા	૫૬	૬૦	૫૯	૬૧	૬૦
પવનની ગતિ (કિમી/કલાક)	૦૧૫	૦૧૮	૦૧૬	૦૧૫	૦૧૪
પવનની દિશા (ડીગ્રી)	૨૭૧	૨૭૦	૨૬૩	૨૭૫	૨૮૩

ગીર સોમનાથ

વરસાદ (મી.મી)	૦૦	૦૦	૦૦	૦૦	૦૦
મહત્તમ તાપમાન (સે.ગ્રે)	૩૭	૩૬	૩૫	૩૫	૩૬
લઘુત્તમ તાપમાન (સે.ગ્રે)	૨૭	૨૮	૨૮	૨૮	૨૮
વાદળની સ્થિતિ	ચોખ્ખું	ચોખ્ખું	ચોખ્ખું	મોટેભાગે ચોખ્ખું	ચોખ્ખું
સવારના ભેજના ટકા	૯૦	૯૧	૯૨	૯૧	૯૧
બપોર બાદ ભેજના ટકા	૬૩	૬૦	૬૬	૭૦	૬૭
પવનની ગતિ (કિમી/કલાક)	૦૧૭	૦૧૭	૦૧૫	૦૧૭	૦૧૮
પવનની દિશા (ડિગ્રી)	૨૮૭	૨૭૪	૨૭૧	૨૮૦	૨૯૩

દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકિય વિભાગ

પાકનું નામ	પાક અવસ્થા	રોગ / જીવાત / જાત	કૃષિ સલાહ
			<ul style="list-style-type: none"> ❖ ઉનાળામાં જમીનની ઉંડી ખેડ કરવી જેથી રોગ અને કીટકોની જુદી જુદી અવસ્થાએ સુર્યતાપ સામે જમીન ખુલ્લી થતા નાશ પામે. ❖ કોમ્પોસ્ટ, દેશી ખાતર અને જીપ્સમ ભલામણ મુજબ યાસમાં ભરવા. ❖ ટમેટા, ભીંડી, બીજા શાકભાજી અને ફળપાકોમાં પીયત આપવું. વર્તમાન સુકી હવામાનની સ્થિતિમાં ભેજની જાળવણી માટે છોડની આજુબાજુ પ્લાસ્ટિક અથવા ઘાસચારાના પાકોનું આવરણ કરવું. ❖ હાલની હવામાનની પરિસ્થિતિનાં કારણે ઉનાળુ શાકભાજી, મગફળી, તલ અને ઘાસચારાના પાકોને સાંજના સમયે જરૂરિયાત મુજબ હળવું પીયત આપવું.
ઉનાળું મગફળી	ડોડવા / પરિપક્વ અવસ્થા	-	<ul style="list-style-type: none"> ❖ જો પાણીની અછત હોય તો મગફળીનાં પાકમાં ડોડવા અવસ્થા વખતે અવશ્ય પીયત આપવું. ❖ મગફળીનો પાક પરીપક્વ થાય એટલે પીયત બંધ કરવું. ડોડવા પરીપક્વ થયે પીયત આપવાથી મગફળીનો ઉગાવો થવાથી ઉત્પાદન અને ગુણવત્તા ઘટે છે અને સમયસર કાપણી કરવી.
ઉનાળું તલ	બચઢાં	થ્રીપ્સ	❖ તલના પાકમાં થ્રીપ્સ, નિયંત્રણ માટે અથવા ૧૫ મીલી

	અવસ્થા		<p>ડાયમીથોએટ અથવા ૧૫ મીલી મીથાઈલ-ઓ-ડેમેટોન દવા ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો.</p> <p>❖ વધારે ગરમીના દિવસોમાં પાકને વિકાસની અવસ્થાએ ટુંકા ગાળે પીયત આપવું.</p>
ઉનાળુ મગ	શીંગ / પરિપક્વ અવસ્થા	-	<p>❖ મગના પાકને દાણા ભરાવવાની અવસ્થાએ પીયત આપવું.</p> <p>❖ મગના પાકની પરીપક્વ શીંગોની વીણી કરવી.</p>
ઉનાળુ અડદ	શીંગ / પરિપક્વ અવસ્થા	-	<p>❖ અડદના પાકને દાણા ભરાવવાની અવસ્થાએ પીયત આપવું.</p> <p>❖ અડદના પાકની પરીપક્વ શીંગોની વીણી કરવી.</p>
આંબો	ફળ / પરિપક્વ અવસ્થા	-	<p>❖ સમયસર પીયત આપવું. શાખ પડે એટલે પરીપક્વ કેરી ઉતારી ગ્રેડીંગ કરી બજારમાં મોકલવી.</p>
શાકભાજી	વૃદ્ધી/કુલ/ ફળ	-	<p>❖ વેલાવાળા શાકભાજીના પાકમાં પાન કોરીયાનો ઉપદ્રવ ટાળવા માટે મુખ્ય પાકની ફરતે દિવેલા તથા ગલગોટાનું વાવેતર કરવાથી તેનું નિયંત્રણ કરી શકાય છે જો ઉપદ્રવ વધુ જોવા મળે તો ૧૦ મીલી મીથાઈલ ઓ ડેમેટોન (૨૫ ઈસી) અથવા ૧૦ મીલી ડાયમીથોએટ (૩૦ ઈસી) ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો ત્યાર બાદ પછીથી બીજો છંટકાવ ૧૫ દિવસ પછી કરવો.</p> <p>❖ શાકભાજીના પાકમાં ચુસીયા પ્રકારની જીવાતના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦ મીલી અથવા લીબોળીનું મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦મીલી ૧૦ લી. પાણીમાં ભેળવી છંટકાવ કરવો. જો વધુ ઉપદ્રવ હોય તો એસીટામ્રાપ્રીડ ૨૦ ટકા ૫ ગ્રામ/૧૦લી. અથવા દ્રાયઝોફોસ ૪૦ ટકા ૨૦ મીલી/ ૧૦ લી.પાણીમાં મીશ્ર કરી છંટકાવ કરવો.</p> <p>❖ મરચીના પાકમાં થ્રીપ્સના નિયંત્રણ માટે ૨૦મીલી પ્રોફેનોફોસ અને લિમ્બોળીનું તેલ ૫૦ મીલી ૧૦ લીટર પાણી મિશ્રણ કરી છાંટવી.</p> <p>❖ શાકભાજીના પાકની રોપણી બાદ ચુસીયા પ્રકારની જીવાતના નિયંત્રણ માટે દાણાદાર ફોરેટ ૧૦ જી ઓડ દીઠ ૨ ગ્રામ મુજબ આપીને પીયત આપવું.</p> <p>❖ મરચીના પાકમાં વાઈરસના કારણે પીળા થઈ કુકડાઈ ગયેલ છોડ ઉપાડી જમીનમાં દાટી નાશ કરવો અને ૫ મીલી ઈમીડાક્લોપ્રાઈડ ૧૦ લીટર પાણીમાં ભેળવી</p>

			<p>છંટકાવ કરવો.</p> <ul style="list-style-type: none"> ❖ રીંગણ અને ટમેટાના ફળ અને અને ડોકાની ઈયળોનો ઉપદ્રવ જણાય તો નુકશાન પામેલ ફળ અને ડોકાને તોડીને જમીનમા દાટી દેવા અને રીનાક્ષીપાયર દવા ૩મીલીદવા ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો. ❖ રીંગણ અને ટમેટાના પાકમા ફળમાખીના નિયંત્રણ માટે એસીટેમાપ્રીડ ૨૦ ટકા પગામ દવા ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો. ❖ શાકભાજી ને સમયસર ઉતારી ગ્રેડીંગ કરી બજારમાં મુકવા.
પશુપાલન	-	-	<ul style="list-style-type: none"> ❖ તાપમાન વધારે રહેવાના કારણે ઢોરને દિવસના સમયે છાયડામા રાખવા અને મિનરલ મીક્ષર્યુકત પીવાના સ્વચ્છ પાણીની વ્યવસ્થા કરવી. ❖ પશુઓને યોગ્ય આહાર આપવો અને પશુઓના રહેઠાણ ચોખ્ખા રાખવા. ❖ માખી મચ્છરથી રક્ષણ માટે ફીનાઈલ છાટવું. ❖ પશુઓને યોગ્ય આહારમાં ખનિજયુક્ત મિશ્રણ સાથે સંગ્રહિત ચારા આપો. ચારા ને સલામત સ્થળે સંગ્રહિત કરવો જોઈએ. ❖ ચારા માટે ચાફ કટર વાપરો. ❖ પશુઓમાં ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટાઝ ૨ મીલી/ ૧ લીટર પાણીમાં નાખીને છાટવી. ❖ દુધાળા પશુઓના આઉને ઝીંક ઓક્સાઈડ અથવા બોરીક પાવડરથી બરાબર સાફ કરવા.

JUNAGADH

Phone: 0285-2670139

Date: -24-05-2019

Significant weather for the preceding week

Rain fall (mm)	000.0
Maximum temperature °C	40.9
Minimum temperature °C	26.0
RH % (Morning/Afternoon)	80 – 32
Wind speed (Kmph)	7.5
Cloud cover	0 - 2

(Period from 25 to 29 May 2019)

Weather forecast for Junagadh Dist. until 08:30 hrs of 29-05-2019

Parameters	Day-1 25-05-19	Day-2 26-05-19	Day-3 27-05-19	Day-4 28-05-19	Day-5 29-05-19
Rain fall (mm)	00	00	00	00	00
Maximum temperature °C	39	39	39	39	40
Minimum temperature °C	27	27	27	28	27
Cloud cover (Okta)	0	0	0	1	0
RH I % (Morning)	87	88	90	89	87
RH II % (After noon)	42	40	43	47	45
Wind speed (Km/h)	015	016	014	015	015
Wind direction(Deg)	279	269	264	276	288

Agro meteorological Advisories

General Advice	Rain fall Temperature	There is forecasting of no rainfall during next five days. Maximum and minimum temperature likely to be in range of 39 - 40°C and 27 - 28°C,
-----------------------	--	---

	RH (%) Wind Direction and Wind Speed		respectively. The Relative Humidity is likely to be in the range of 90 – 40 %. Direction and speed of wind is likely to be mostly W and 14 - 16 km/h, respectively.
Crop	Variety/ Pest/Insect	Stage	Agro met Advisories
			<ul style="list-style-type: none"> ➤ To cultivate deep in the soil summer season so that diseases and pests are destroyed at different stages by the direct sunlight. ➤ Compost, FYM fertilizers and gypsum filling in the furrow as recommended. ➤ Apply irrigation in tomato, bhindi, other vegetables and fruit crops. In present dry weather condition, spread plastic or fodder as mulch in vegetable fields for conserving moisture. ➤ Irrigate summer vegetable and fodder crop during night hours.
Summer Groundnut	-	Pod development / Maturity	<ul style="list-style-type: none"> ➤ Apply irrigation at the time of pod development stage. ➤ Postponed the irrigation at the time of maturity. If irrigation is applied at the time of pod maturity, yield is reduced and also affect the quality of pod.
Summer Sesame	Thrips	Capsule formation	<ul style="list-style-type: none"> ➤ For the control of Thrips, spray 15 ml dimethoate OR 15 ml methyl-O-demeton mix in 10 liter of water. ➤ During hot summer, apply irrigation to the crop at the critical growth stage.

Summer green gram	-	Pod formation / Maturity	<ul style="list-style-type: none"> ➤ Apply irrigation at the time of pod development. ➤ Harvesting operation should be carried out at maturity stage.
Summer black gram	-	Pod formation / Maturity	<ul style="list-style-type: none"> ➤ Apply irrigation at the time of pod development. ➤ Harvesting operation should be carried out at maturity stage.
Mango	-	Fruiting / Maturity	<ul style="list-style-type: none"> ➤ Apply irrigation regularly and send mango fruit to the market often grading.
Vegetable crops	-	Sowing/ Flowering/ Fruiting	<ul style="list-style-type: none"> ➤ For the control of leaf webber in fruit leafy vegetable, sowing of trap crop (castor and galgota) around the vegetable crop. For the heavy incidence spray 10 ml methyl O demeton (25 EC) Or 10 ml diamethoate (30 EC) in 10 liter of water. ➤ For the control of thrips and white fly, spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. ➤ For the control of thrips in chilli, spray 20 ml profenophos in 10 litres of water. ➤ For the control of fruit and shoot borer, spray neem seed extract 5 % or neem seed oil 50 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. ➤ Rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of 5 mi Imidacloprid per 10 litre of water is

			<p>advised for control the vector, when sky remains clear.</p> <ul style="list-style-type: none"> ➤ Apply regular irrigation at 6 to 8 days to the crop at the flowering and fruiting stage. ➤ In brinjal and tomato crops constant monitoring of borers by using pheromone traps 4-6 per acre is advised, for the control of fruit borer, spray reenaxipayar @ 3.0ml in 10 lit of water. ➤ In brinjal and tomato crops, for the control of fruit fly, spray acetemaprid 5.0 ml in 10 lit of water. ➤ Keep vegetable crop weed free, apply irrigation regularly and send vegetable fruit to the market often grading.
Livestock	-	-	<ul style="list-style-type: none"> ➤ Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day. ➤ To provide adequate food and keep the animals clean shed. ➤ Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Give stored fodder with mineral mixture. Fodder should be stored at safe place. ➤ Feed silage & hay material along with concentrate feed. ➤ Use chaff cutter for fodder. ➤ For the control of tick and other parasites, spray deltemethrin or aemitaz 2 ml in 1 litre of water. ➤ Udder of milking animals must be properly

clean with zinc oxide or boric powder.

PORBANDAR

Phone: 0285-2670139

Date: - 24-05-2019

(Period from 25 to 29 May 2019)

Weather forecast for **Porbandar** Dist. until 08:30 hrs of 29-05-2019

Parameters	Day-1 25-05-19	Day-2 26-05-19	Day-3 27-05-19	Day-4 28-05-19	Day-5 29-05-19
Rain fall (mm)	00	00	00	00	00
Maximum temperature °C	37	36	37	36	37
Minimum temperature °C	27	27	27	27	27
Cloud cover (Okta)	0	0	0	0	0
RH I % (Morning)	87	87	88	87	87
RH II % (After noon)	56	60	59	61	60
Wind speed (Km/h)	015	018	016	015	014
Wind direction(Deg)	271	270	263	275	283

Agro meteorological Advisories

General Advice	Rain fall	<p>There is forecasting of no rainfall during next five days.</p> <p>Maximum and minimum temperature likely to be in range of 36 – 37 °C and 27°C, respectively.</p> <p>The Relative Humidity is likely to be in the range of 88 – 56 %.</p> <p>Direction and speed of wind is likely to be mostly W and 14 – 18 km/h, respectively.</p>	
	Temperature		
	RH (%)		
	Wind Direction and Wind Speed		
Crop	Variety/ Pest / Insect	Stage	Agro met Advisories
			➤ To cultivate deep in the soil summer season so that diseases and pests are destroyed at

			<p>different stages by the direct sunlight.</p> <ul style="list-style-type: none"> ➤ Compost, FYM fertilizers and gypsum filling in the furrow as recommended. ➤ Apply irrigation in tomato, bhindi, other vegetables and fruit crops. In present dry weather condition, spread plastic or fodder as mulch in vegetable fields for conserving moisture. ➤ Irrigate summer vegetable and fodder crop during night hours.
Summer Groundnut	-	Pod development / Maturity	<ul style="list-style-type: none"> ➤ Apply irrigation at the time of pod development stage. ➤ Postponed the irrigation at the time of maturity. If irrigation is applied at the time of pod maturity, yield is reduced and also affect the quality of pod.
Summer Sesame	Thrips	Capsule formation	<ul style="list-style-type: none"> ➤ For the control of Thrips, spray 15 ml dimethoate OR 15 ml methyl-O-demeton mix in 10 liter of water. ➤ During hot summer, apply irrigation to the crop at the critical growth stage.
Summer green gram	-	Pod development / Maturity	<ul style="list-style-type: none"> ➤ Apply irrigation at the time of pod development. ➤ Harvesting operation should be carried out at maturity stage.
Summer black gram	-	Pod development / Maturity	<ul style="list-style-type: none"> ➤ Apply irrigation at the time of pod development. ➤ Harvesting operation should be carried out at maturity stage.
Mango	-	Fruiting /	<ul style="list-style-type: none"> ➤ Apply irrigation regularly and send mango fruit

		Maturity	to the market often grading.
Vegetable crops	-	Sowing/ Flowering/ Fruiting	<ul style="list-style-type: none"> ➤ For the control of leaf webber in fruit leafy vegetable, sowing of trap crop (castor and galgota) around the vegetable crop. For the heavy incidence spray 10 ml methyl O demeton (25 EC) Or 10 ml diamethoate (30 EC) in 10 liter of water. ➤ For the control of thrips and white fly, spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. ➤ For the control of thrips in chilli, spray 20 ml profenophos in 10 litres of water. ➤ For the control of fruit and shoot borer, spray neem seed extract 5 % or neem seed oil 50 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. ➤ Rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of 5 ml Imidacloprid per 10 litre of water is advised for control the vector, when sky remains clear. ➤ Apply regular irrigation at 6 to 8 days to the crop at the flowering and fruiting stage. ➤ In brinjal and tomato crops constant monitoring of borers by using pheromone traps 4-6 per acre is advised, for the control of fruit borer, spray reanaxipayar @ 3.0ml in 10 lit of water.

			<ul style="list-style-type: none"> ➤ In brinjal and tomato crops, for the control of fruit fly, spray acetemaprid 5.0 ml in 10 lit of water. ➤ Keep vegetable crop weed free, apply irrigation regularly and send vegetable fruit to the market often grading.
Livestock	-	-	<ul style="list-style-type: none"> ➤ Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day. ➤ To provide adequate food and keep the animals clean shed. ➤ Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Give stored fodder with mineral mixture. Fodder should be stored at safe place. ➤ Feed silage & hay material along with concentrate feed. ➤ Use chaff cutter for fodder. ➤ For the control of tick and other parasites, spray deltemethrin or aemitaz 2 ml in 1 litre of water. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.

Gir Somnath

Phone: 0285-2670139

Date: - 24-05-2019

(Period from 25 to 29 May 2019)

Weather forecast for Gir Somnath Dist. until 08:30 hrs of 29-05-2019

Parameters	Day-1 25-05-19	Day-2 26-05-19	Day-3 27-05-19	Day-4 28-05-19	Day-5 29-05-19
Rain fall (mm)	00	00	00	00	00
Maximum temperature °C	37	36	35	35	36
Minimum temperature °C	27	28	28	28	28
Cloud cover (Okta)	0	0	0	1	0
RH I % (Morning)	90	91	92	91	91
RH II % (After noon)	63	60	66	70	67
Wind speed (Km/h)	017	017	015	017	018
Wind direction(Deg)	287	274	271	280	293

Agro meteorological Advisories

General Advice	Rain fall Temperature RH (%) Wind Direction and Wind Speed		<p>There is forecasting of no rainfall during next five days.</p> <p>Maximum and minimum temperature likely to be in range of 35 – 37 °C and 27 - 28°C, respectively.</p> <p>The Relative Humidity is likely to be in the range of 92 – 60 %.</p> <p>Direction and speed of wind is likely to be mostly W & NW and 15 – 18 km/h, respectively.</p>
Crop	Variety/ Pest / Insect	Stage	Agro met Advisories
			➤ To cultivate deep in the soil summer season so that diseases and pests are destroyed at

			<p>different stages by the direct sunlight.</p> <ul style="list-style-type: none"> ➤ Compost, FYM fertilizers and gypsum filling in the furrow as recommended. ➤ Apply irrigation in tomato, bhindi, other vegetables and fruit crops. In present dry weather condition, spread plastic or fodder as mulch in vegetable fields for conserving moisture. ➤ Irrigate summer vegetable and fodder crop during night hours.
Summer Groundnut	-	Pod development / Maturity	<ul style="list-style-type: none"> ➤ Apply irrigation at the time of pod development stage. ➤ Postponed the irrigation at the time of maturity. If irrigation is applied at the time of pod maturity, yield is reduced and also affect the quality of pod.
Summer Sesame	Thrips	Capsule formation	<ul style="list-style-type: none"> ➤ For the control of Thrips, spray 15 ml dimethoate OR 15 ml methyl-O-demeton mix in 10 liter of water. ➤ During hot summer, apply irrigation to the crop at the critical growth stage.
Summer green gram	-	Pod development / Maturity	<ul style="list-style-type: none"> ➤ Apply irrigation at the time of pod development. ➤ Harvesting operation should be carried out at maturity stage.
Summer black gram	-	Pod development / Maturity	<ul style="list-style-type: none"> ➤ Apply irrigation at the time of pod development. ➤ Harvesting operation should be carried out at maturity stage.
Mango	-	Fruiting /	<ul style="list-style-type: none"> ➤ Apply irrigation regularly and send mango fruit

		Maturity	to the market often grading.
Vegetable crops	-	Sowing/ Flowering/ Fruiting	<ul style="list-style-type: none"> ➤ For the control of leaf webber in fruit leafy vegetable, sowing of trap crop (castor and galgota) around the vegetable crop. For the heavy incidence spray 10 ml methyl O demeton (25 EC) Or 10 ml diamethoate (30 EC) in 10 liter of water. ➤ For the control of thrips and white fly, spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. ➤ For the control of thrips in chilli, spray 20 ml profenophos in 10 litres of water. ➤ For the control of fruit and shoot borer, spray neem seed extract 5 % or neem seed oil 50 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. ➤ Rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of 5 ml Imidacloprid per 10 litre of water is advised for control the vector, when sky remains clear. ➤ Apply regular irrigation at 6 to 8 days to the crop at the flowering and fruiting stage. ➤ In brinjal and tomato crops constant monitoring of borers by using pheromone traps 4-6 per acre is advised, for the control of fruit borer, spray reanaxipayar @ 3.0ml in 10 lit of water.

			<ul style="list-style-type: none"> ➤ In brinjal and tomato crops, for the control of fruit fly, spray acetemaprid 5.0 ml in 10 lit of water. ➤ Keep vegetable crop weed free, apply irrigation regularly and send vegetable fruit to the market often grading.
Livestock	-	-	<ul style="list-style-type: none"> ➤ Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day. ➤ To provide adequate food and keep the animals clean shed. ➤ Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Give stored fodder with mineral mixture. Fodder should be stored at safe place. ➤ Feed silage & hay material along with concentrate feed. ➤ Use chaff cutter for fodder. ➤ For the control of tick and other parasites, spray deltemethrin or aemitaz 2 ml in 1 litre of water. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.