

 	<p>ગ્રામીણ કૃષિ મૌસમ સેવા</p> <p>કૃષિ હવામાન સેલ</p> <p>કૃષિ વિજ્ઞાન વિભાગ</p> <p>જુનાગઢ કૃષિ યુનિવર્સિટી</p> <p>જુનાગઢ - ૩૬૨૦૦૧</p> <p>ઈમેઇલ:- amfujind@yahoo.com</p> <p>ફોન નં. ૦૨૮૫-૨૬૭૦૧૩૬</p>	
--	--	---

હવામાન આગાહી આધારીત દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારનાં જુનાગઢ જીલ્લા માટે કૃષિ સલાહ બુલેટીન
(કૃષિ હવામાન સેલ, જુનાગઢ કૃષિ યુનિવર્સિટી અને ભારત હવામાન વિભાગ દ્વારા સંપદ્ધ)

વાર:- મંગળવાર

તારીખ:- ૨૫/૦૬/૨૦૨૪

હવામાન આગાહી આધારીત બુલેટીન: ૦૨૫ (૨૦૨૪-૨૫)

પાછલા પાંચ દિવસનું હવામાન (તારીખ:- ૨૦ થી ૨૪ જૂન, ૨૦૨૪)					હવામાન પરિબળો	હવામાન આગાહી આગામી પાંચ દિવસ માટે (તારીખ:- ૨૬ થી ૩૦ જૂન, ૨૦૨૪) માન્ય ૩૦ જૂન, ૨૦૨૪. સવારના ૮:૩૦ વાગ્યાં સુધી				
૨૦/૦૬	૨૧/૦૬	૨૨/૦૬	૨૩/૦૬	૨૪/૦૬	તારીખ	૨૬/૦૬	૨૭/૦૬	૨૮/૦૬	૨૯/૦૬	૩૦/૦૬
૦	૦	૦	૦	૧૦૨.૬	વરસાદ(મીમી)	૭૬	૧૩	૧૬	૩૧	૪૪
૩૮.૫	૩૮.૩	૩૮.૦	૩૮.૭	૩૮.૬	મહત્વમાન (સે.ગ્રે)	૩૩	૩૩	૩૨	૩૨	૩૩
૨૮.૧	૨૮.૬	૨૮.૪	૨૮.૩	૨૮.૫	લધુત્વમાન (સે.ગ્રે)	૨૫	૨૪	૨૪	૨૪	૨૪
અંશતા વાદળાયું	અંશતા વાદળાયું	અંશતા વાદળાયું	અંશતા વાદળાયું	મોટેલાગે વાદળાયું	વાદળની સ્થિતિ	વાદળાયું	મોટેલાગે વાદળાયું	વાદળાયું	વાદળાયું	વાદળાયું
૭૬	૭૬	૭૮	૭૮	૮૫	સવારના સેજના ટકા	૮૨	૮૧	૭૬	૮૨	૮૪
૫૨	૪૬	૪૫	૪૮	૭૬	બપોરબાદ ના સેજના ટકા	૬૨	૪૬	૫૮	૫૦	૭૩
૧૪.૬	૧૮.૨	૧૫.૩	૬.૧	૫.૨	પવનની ગતે (કીમી/કલાક)	૨૨.૦	૧૭.૦	૨૧.૦	૨૦.૦	૨૪.૦
૨૩૦	૨૫૦	૨૭૦	૨૫૦	૨૬૦	પવનની દિશા	૨૬૨	૨૪૮	૨૪૮	૨૪૮	૨૪૮

સામાન્ય સલાહ

ભારત સરકારશીના હવામાન વિભાગ તરફથી મળેલ હવામાન આગાહીની માહીતી અનુસાર અત્રેનાં ગ્રામીણ કૃષિ મૌસમ સેવા યોજના, જુનાગઢ કૃષિ યુનિવર્સિટી, જુનાગઢ તરફથી જણાવવામા આવે છે કે દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારનાં જુનાગઢ જીલ્લામાં આગામી તારીખ ૨૬/૦૬/૨૦૨૪ થી ૩૦/૦૬/૨૦૨૪ દરમયાન આકાશ મુખ્યત્વે આગામી પાંચ દિવસોમાં મુખ્યત્વે મોટેલાગે વાદળાયું થી વાદળાયું રહેશે, જીલ્લામાં આવતીકાલે ભારે વરસાદ અને પાંચ દિવસો દરમયાન છુટા છવાયાં ભાગોમાં હળવા થી મધ્યમ વરસાદની શક્યતાઓ છે. પવનની ઝડપ ૧૭ થી ૨૪ કીમી/કલાક રહેવાની શક્યતા રહેશે. આ સમયગાળામાં મહત્વમાં તાપમાન ૩૨ થી ૩૩ ડિગ્રી સેન્ટીગ્રેડ રહેશે તેમજ લધુત્વમાન ૨૪ થી ૨૫ ડિગ્રી સેન્ટીગ્રેડ રહેવાની સંભાવના છે.

માન્ય વરસાદની સૂચી મુજબ જુનાગઢ જુહ્લામાં ૨૭ સપ્ટેમ્બર થી ૨૫ ઓક્ટોબર, ૨૦૨૩ નાં સમયગાળા દરમિયાન મધ્યમ લેજની સ્થિતિ જોવા મળેલ છે.

સાપ્તાહિક હવામાન આગાહીની સૂચી મુજબ સૌરાષ્ટ્ર અને કચ્છનાં વિસ્તારોમાં તા. ૩૦/૦૯/૨૦૨૪ થી તા. ૦૬/૦૭/૨૦૨૪ દરમિયાન વરસાદ સામાન્ય થી ઉપર & મહત્વમાન અને લધુતમ તાપમાન સામાન્ય રહેવાની શક્યતાઓ છે.


મેધદૂત એપ્લીકેશન:- કૃષિ હવામાન સલાહ સેવા અને હવામાન આગાહી અંગેની જાણકારી મેળવવા માટે ડાઉનલોડ કરો મેધદૂત એપ્લીકેશન કે જે ખેડૂતોને તેમની સ્થાનિક ભાષામાં સ્થાન, પાક અને પશુધન વિશે હવામાન આધારિત કૃષિ સલાહ પ્રદાન કરશે.

મેધદૂત એપ્લીકેશનને ડાઉનલોડ કરવા માટે નીચેની લિંક પર ક્લિક કરો.

એન્ડ્રોઈડ(મોબાઈલ) ઉપયોગકર્તા: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>

આઇ.ઓ.એસ. ઉપયોગકર્તા: <https://apps.apple.com/in/app/meghdoot/id1474048155>

- ✓ જો સૂકાયેલ કણસલાં કે કાપણી કરેલ ઉનાળું પાક/ધાસચારો ખેતરમાં પડેલ હોય તો યોગ્ય જગ્યાએ સંગ્રહ કરવો.
- ❖ ખેતરમાંથી જમીનના નમુના લેવા અને જમીન પૃથ્વીકરણ કરવા માટે લેબોરેટરીમાં મોકલી આપવા.
- ❖ ખેતરના સેઢા, પાળા અને વરસાદના પાણી નિકાલની ગટર ને ચોખ્માં કરી ને આગામી ચોમાસા માટે ખેતર ને તૈયાર કરવું.
- ❖ કોમ્પોસ્ટ, દેશી ખાતર અને જુખ્સમ ભલામણ મુજબ ચાસમાં ભરવા.
- ❖ આગામી ચોમાસાની ઋતુમાં વાવણી માટે ખેડૂતોને સારી ગુણવત્તા અને વધારે ઉત્પાદન આપતી જતોના પ્રમાણિત બીજની પસંદગી કરીને મેળવી લેવાની સલાહ આપવામાં આવે છે.
- ❖ વાવણી લાયક વરસાદ થયે ચોમાસું પાકનું વાવેતર કરવું.



પાકનું નામ	પાક અવસ્થા	રોગ/જીવાત/જાત	કૃષિ સલાહ
------------	------------	---------------	-----------

ચોમાસું મગફળી	વાવેતર / ઉગાવો	અર્ધવેલડી જાત- જીજી.૨૦ અથવા ઉભડી જાતો જીજી.૨ જીજી.૫ જીજી.૭ ટી.જી. ૨૬ / મુંડા, ચુસીયા જીવાતો, ઉગસુક, થડનો કોહવારો	<ul style="list-style-type: none"> ❖ વાવેતર માટે અર્ધવેલડી જાત- જીજી.૨૦ અથવા ઉભડી જાતો જીજી.૨ જીજી.૫ જીજી.૭ ટી.જી. ૨૬ ની પસંદગી કરવી. ❖ ચોમાસુ મગફળીના પાકને હેકટરે ૧૨.૫ કિ.ગ્રા નાઇટ્રોજન અને ૨૫ કિ.ગ્રા ફોસ્ફરસ ખાતર તરીકે વાવણી વખતે ચાસમાં ઓરીને આપવા. ❖ વાવેતર સમયે ડીએપી ૫૪.૩ કીલો અને ૬ કીલો યુરીયા/હે. અને એરેનીનો ખોળ ૫૦૦ કી.ગ્રા./હે. અથવા જીપ્સમ ૫૦૦ કી.ગ્રા./હે. સાથે એરેનીનો ખોળ ૫૦૦ કી.ગ્રા./હે પાયામા આપવો. ❖ વાવતી વખતે બીયારણને થાયરમ અથવા મેંકોઝેબ અથવા વીટાવેક્ષ દવાનો ૨-૩ ગ્રામ/કી. બીજને પટ આપવો. ❖ મુંડાના નિયંત્રણ માટે કલોરપાચરીફોસ ૨૦ % છ.સી. ૨૫ મિલી/કિલો બીજ અથવા ઈમીડાકલોપ્રીડ ૬૦૦ % એફ.એસ. ૪ મિલી/કિલો બીજ મુજબ બીજ માવજત આપવી. ❖ બીવેરિયા બાસીયાના અથવા મેટારીઝીયમ એનીસોપ્ટી નામની ફૂગનો પાવડર વાવેતર પહેલા ૫ કિલો પ્રતિ હેકટર મુજબ એરેનીના ખોળ સાથે ચાસમાં આપવાથી મુંડાનું નિયંત્રણ કરી શકાય છે. અથવા કાર્બોફ્યુરાન ૩૪ નામની દાણાદાર દવાને ૩૩ કિલો/હેકટર મુજબ ચાસમાં આપવી. ❖ ચુસીયા જીવાતોના નિયંત્રણ માટે ઈમીડાકલોપ્રીડ ૬૦૦ % એફ.એસ. ૪ મિલી / કિલો અથવા થાયામીથોકઝામ ૭૦ % ડબલ્યુ.એસ. ૧ ગ્રામ / કિલો બીજ મુજબ બીજ માવજત આપવી. ❖ વાવણી લાયક પૂરતા વરસાદ થયે પાકનું વાવેતર સમયસર કરવું.
કપાસ	વાવેતર	ગુજરાત હાઈબ્રીડ કપાસ-૬, ૮, ૧૨ (બોલગાડ્-૨) કે બી.ટી. કપાસ / સુકારો અને મૂળનો કોહવારો	<ul style="list-style-type: none"> ❖ કપાસ વાવણી માટે નવી બહાર પાડેલ વધુ ઉત્પાદન આપતી બીઠી હાઈબ્રીડ કપાસ ગુજરાત હાઈબ્રીડ કપાસ-૬, ૮, ૧૨ (બોલગાડ્-૨) પસંદ કરવી કે બી.ટી. કપાસનું વિશ્વાષપાત્ર બીયારણ જ ખરીદવું. ❖ કપાસનું વિશ્વાષપાત્ર સર્ટીફાઈડ બીજ ખરીદવું. ❖ પાકની ફેરબદલી કરવી. કપાસની સાથે મઠનો આંતરપાક લેવો. ❖ ઈક્કડના પાકનો લીલો પડવાશ કરવો. ❖ સ્યુડોમોનાસ ફલુરોસન્સ ૧%, ૧૦ ગ્રામ અથવા કાર્બોકઝીન ૩૭.૫ % + થાયરમ ૩૭.૫ % ડી. એસ., ૩.૫ ગ્રામ પ્રતિ કીલો બીજને પટ આપવો. ❖ વાવણી લાયક પૂરતા વરસાદ થયે પાકનું વાવેતર સમયસર કરવું.
સોયાબીન	વાવેતર	ગુજ. સોયાબીન - ૧, ૨ &	<ul style="list-style-type: none"> ❖ સોયાબીન પાકનાં વાવેતર માટે સારી ગુણવત્તા અને વધારે ઉત્પાદન આપતી જાતોનાં પ્રમાણિત બીજની પસંદગી કરીને મેળવી લેવું.

		જીજેએસ - 3 / ચુસીયા જીવાતો, કોલાર રોટ, ચારડોલ રોટ	<ul style="list-style-type: none"> ❖ પાકમાં વાવેતર સમયે પાચામાં એસએસપી ૩૭૫ કીલો અને ૬૫.૨ કીલો ચુસીયા/હે આપવું. ❖ ઈમીડાકલોપ્રીડ ૪૮ એફ.એસ. @૩ મીલી/કિગ્રા સાથે બીજ માવજત આપવી. ❖ ફલુકાપાયરોક્સેડ ૩૩૩ ગ્રામ/લી એફએસ ને ૧.૦ મીલી/કિલો અથવા કાબોકઝીન ૩૭.૫% + થાયરમ ૩૭.૫% ડબલ્યુએસ ની ૩ ગ્રામ/કિલો બીયારણ મુજબ બીજ માવજત આપવી. ❖ વાવણી લાયક પૂરતા વરસાદ થયે પાકનું વાવેતર સમયસર કરવું.
આંબા	ફળ પરિપક્વતા	-	<ul style="list-style-type: none"> ❖ આંબાનાં બગીચામાં નિયમિત પીયત આપવું. ❖ શાખ પડે એટલે પરીપક્વ કેરી ઉતારી ગ્રેડીંગ કરી બજારમાં વહેંચાણ માટે મોકલવી.
શાકભાજુ	વૃદ્ધી / કુલ / ફળ	-	<ul style="list-style-type: none"> ❖ પાકમાં દવા છાંટવાનું અને ખાતર આપવનું કામ આ સમયગાળામાં કરવું. ❖ પાકમાં હાથથી નિંદામણ અને આંતરખેડની કામગીરી કરવી. ❖ શાકભાજુ પાકોને સમયસર પીયત આપવું. ❖ શાકભાજુના પાકોને નીદામણ મુકત કરવું અને નિયમિત પિયત આપવું. શાકભાજુ ને સમયસર ઉતારી ગ્રેડીંગ કરી બજારમાં મુકવા.
પશુપાલન		-	<ul style="list-style-type: none"> ❖ પશુઓને યોગ્ય આહાર આપવો & પીવા માટે ચોખ્યું ઠંડુ પાણી આપવું અને પશુઓના રહેઠાણ ચોખ્યા રાખવા. ❖ માખી મચ્છરથી રક્ષણ માટે ફીનાઈલ છાટવું. ❖ પશુઓને યોગ્ય આહારમાં ખનિજયુક્ત મિશ્રણ સાથે સંગ્રહિત ચારા આપો. ચારા ને સલામત સ્થળે સંગ્રહિત કરવો જોઈએ. ❖ ચારા માટે ચાફ કટર વાપરો. ❖ પશુઓમાં ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટાઇ ૨ મીલી/ ૧ લીટર પાણીમાં નાખીને છાટવી. ❖ દુધાળા પશુઓના આઉને ઝીંક ઓક્સાઈડ અથવા બોરીક પાવડરથી બરાબર સાફ કરવા.



ગ્રામીણ કૃષિ મૌસમ સેવા
કૃષિ હવામાન સેલ
કૃષિ વિજ્ઞાન વિભાગ
જૂનાગઢ કૃષિ યુનિવર્સિટી
જૂનાગઢ - ૩૬૨૦૦૧
ઈમેઇલ:-amfujnd@yahoo.com ફોન નં. ૦૨૮૫-૨૬૭૦૧૩૯



હવામાન આગાહી આધારીત દક્ષિણ સૌરાષ્ટ્ર એત આબોહવાકીય વિસ્તારનાં પોરબંદર જીલ્લા માટે કૃષિ સલાહ બુલેટીન
(કૃષિ હવામાન સેલ, જૂનાગઢ કૃષિ યુનિવર્સિટી અને ભારત હવામાન વિભાગ દ્વારા સંપદ્ધ)

વાર:- મંગળવાર

તારીખ: ૨૫/૦૬/૨૦૨૪

હવામાન આગાહી આધારીત બુલેટીન: ૦૨૫ (૨૦૨૪-૨૫)

પાછલા પાંચ દિવસનું હવામાન (તારીખ:- ૨૦ થી ૨૪ જૂન, ૨૦૨૪)					હવામાન પરિબળો	હવામાન આગાહી આગામી પાંચ દિવસ માટે (તારીખ:- ૨૫ થી ૩૦ જૂન, ૨૦૨૪)				
૨૦/૦૬	૨૧/૦૬	૨૨/૦૬	૨૩/૦૬	૨૪/૦૬	તારીખ	૨૬/૦૬	૨૭/૦૬	૨૮/૦૬	૨૯/૦૬	૩૦/૦૬
૦	૦	૦	૦	૪.૬	વરસાદ(મિમી)	૧૪	૨	૮	૧૬	૨૦
૩૪.૮	૩૫.૦	૩૫.૨	૩૫.૪	૩૫.૬	મહત્વમાન તાપમાન (°સે.ગ્રે)	૩૩	૩૪	૩૪	૩૪	૩૪
૨૯.૮	૩૦.૦	૨૯.૮	૨૯.૬	૨૯.૦	લઘુત્વમાન તાપમાન (°સે.ગ્રે)	૨૯	૨૯	૨૯	૨૯	૨૯
અંશતા વાદળાયું	અંશતા વાદળાયું	મોટેભાગે ચોખ્યાં	મોટેભાગે વાદળાયું	મોટેભાગે વાદળાયું	વાદળની સ્થિતિ	વાદળાયું	મોટેભાગે વાદળાયું	મોટેભાગે વાદળાયું	વાદળાયું	વાદળાયું
૭૬	૭૪	૭૫	૭૫	૬૦	સવારના ભેજના ટકા	૮૧	૮૦	૭૬	૮૦	૮૩
૬૭	૭૪	૭૨	૭૪	૭૨	બપોરબાદ ના ભેજના ટકા	૬૯	૫૫	૬૪	૬૫	૭૧
૧૫.૮	૨૪.૧	૧૬.૭	૧૧.૧	૭.૪	પવનની ગતિ (કિમી/કલાક)	૨૮.૦	૧૬.૦	૨૩.૦	૨૪.૦	૨૮.૦
૨૫૦	૨૫૦	૨૭૦	૩૨૦	૫૦	પવનની દિશા	૨૮૮	૨૪૮	૨૪૮	૨૪૮	૨૪૮

સામાન્ય સલાહ

ભારત સરકારશ્રીના હવામાન વિભાગ તરફથી મળેલ હવામાન આગાહીની માહીતી અનુસાર અત્રેના ગ્રામીણ કૃષિ મૌસમ સેવા યોજના, જૂનાગઢ કૃષિ યુનિવર્સિટી, જૂનાગઢ તરફથી જણાવવામા આવે છે કે દક્ષિણ સૌરાષ્ટ્ર એત આબોહવાકીય વિસ્તારના પોરબંદર જીલ્લામાં આગામી તારીખ ૨૫/૦૬/૨૦૨૪ થી ૩૦/૦૬/૨૦૨૪ દરમ્યાન આકાશ મુખ્યત્વે આગામી પાંચ દિવસોમાં મુખ્યત્વે મોટેભાગે વાદળાયું થી વાદળાયું રહેશે, જીલ્લામાં આગામી પાંચ દિવસો દરમિયાન છુટા છવાયાં ભાગોમાં હળવા થી મધ્યમ વરસાદની શક્યતાઓ છે. પવનની ઝડપ ૧૮ થી ૨૮ કિમી/કલાક રહેવાની શક્યતા રહેશે. આ સમયગાળામાં મહત્વમાન ૩૩ થી ૩૪ ડિગ્રી સેન્ટીગ્રેડ રહેશે તેમજ લઘુત્વમાન ૨૯ ડિગ્રી સેન્ટીગ્રેડ રહેવાની સંભાવના છે.

માન્ય વરસાદની સૂચી મુજબ પોરબંદર જુહ્લામાં ૨૭ સપ્ટેમ્બર થી ૨૫ ઓક્ટોબર, ૨૦૨૩ નાં સમયગાળા દરમિયાન મધ્યામ સૂકી રિઝિટ જોવા મળેલ છે.

સાપ્તાહિક હવામાન આગાહીની સૂચી મુજબ સૌરાષ્ટ્ર અને કચ્છનાં વિસ્તારોમાં તા. ૩૦/૦૬/૨૦૨૪ થી તા. ૦૬/૦૭/૨૦૨૪ દરમિયાન વરસાદ સામાન્ય થી ઉપર & મહત્વમાન અને લઘુત્વમાન સામાન્ય રહેવાની શક્યતાઓ છે.


મેધદૂત એપ્લીકેશન:-કૃષિ હવામાન સલાહ સેવા અને હવામાન આગાહી અંગેની જાણકારી મેળવવા માટે ડાઉનલોડ કરો મેધદૂત એપ્લીકેશન કે જે ખેડૂતોને તેમની સ્થાનિક ભાષામાં સ્થાન, પાક અને પશુધન વિશે હવામાન આધારિત કૃષિ સલાહ પ્રદાન કરશે.

મેધદૂત એપ્લીકેશનને ડાઉનલોડ કરવા માટે નીચેની લિંક પર ક્લિક કરો.

એન્ડ્રોઇડ(મોબાઇલ)ઉપયોગકર્તા: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>

આઈ.ઓ.એસ.ઉપયોગકર્તા: - <https://apps.apple.com/in/app/meghdoot/id1474048155>

- ✓ જો સૂકાયેલ કણસલાં કે કાપણી કરેલ ઉનાળું પાક/ધાસચારો ખેતરમાં પડેલ હોય તો યોગ્ય જગ્યાએ સંગ્રહ કરવો.
- ❖ ખેતરમાંથી જમીનના નમુના લેવા અને જમીન પૃથ્વીકરણ કરવા માટે લેબોરેટરીમાં મોકલી આપવા.
- ❖ ખેતરના સેઢા, પાળા અને વરસાદના પાણી નિકાલની ગટર ને ચોખ્માં કરી ને આગામી ચોમાસા માટે ખેતર ને તૈયાર કરવું.
- ❖ કોપ્પોસ્ટ, દેશી ખાતર અને જુખ્સમ ભલામણ મુજબ ચાસમાં ભરવા.
- ❖ આગામી ચોમાસાની ઋતુમાં વાવણી માટે ખેડૂતોને સારી ગુણવત્તા અને વધારે ઉત્પાદન આપતી જાતોના પ્રમાણિત બીજની પસંદગી કરીને મેળવી લેવાની સલાહ આપવામાં આવે છે.



પાકનું નામ	પાક અવસ્થા	રોગ/જીવાત/જાત	કૃષિ સલાહ
------------	------------	---------------	-----------

<p style="color: red;">ચોમાસું મગફળી</p>	<p style="color: orange;">વાવેતર / ઉગાવો</p>	<p style="color: blue;">અર્ધવેલડી જત- જુજુ.૨૦ અથવા ઉભડી જતો જુજુ.૨ જુજુ.૫ જુજુ.૭ ટી.જી. ૨૬ / મુંડા, ચુસીયા જીવાતો, ઉગાસુક, થડનો કોહવારો</p>	<ul style="list-style-type: none"> ❖ વાવેતર માટે અર્ધવેલડી જત- જુજુ.૨૦ અથવા ઉભડી જતો જુજુ.૨ જુજુ.૫ જુજુ.૭ ટી.જી. ૨૬ ની પસંદગી કરવી. ❖ ચોમાસું મગફળીના પાકને હેકટરે ૧૨.૫ કિ.ગ્રા નાઇટ્રોજન અને ૨૫ કિ.ગ્રા ફોસ્ફરસ ખાતર તરીકે વાવણી વખતે ચાસમાં ઓરીને આપવા. ❖ વાવેતર સમયે ડીએપી ૫૪.૩ કીલો અને ૬ કીલો યુરીયા/હે. અને એરેનીનો ખોળ ૫૦૦ કી.ગ્રા./હે. અથવા જુપ્સમ ૫૦૦ કી.ગ્રા./હે. સાથે એરેનીનો ખોળ ૫૦૦ કી.ગ્રા./હે પાયામા આપવો. ❖ વાવતી વખતે બીયારણને થાયરમ અથવા મેંકોઝેબ અથવા વીટાવેક્ષ દવાનો ૨-૩ ગ્રામ/કી. બીજને પટ આપવો. ❖ મુંડાના નિયંત્રણ માટે કલોરપાયરીઝેસ ૨૦ છ.સી. % ૨૫ મિલી/કિલો બીજ અથવા ઈમીડાકલોપ્રીડ ૬૦૦ એફ.એસ. %૪ મિલી/કિલો બીજ મુજબ બીજ માવજત આપવી. ❖ બીવેરિયા બાસીયાના અથવા મેટારીઝીયમ એનીસોપ્ટી નામની ફૂગનો પાવડર વાવેતર પહેલા ૫ કિલો પ્રતિ હેકટર મુજબ એરેનીના ખોળ સાથે ચાસમાં આપવાથી મુંડાનું નિયંત્રણ કરી શકાય છે. અથવા કાર્બોફ્યુરાન ૩૪ નામની દાણાદાર દવાને ૩૩ કિલો/હેકટર મુજબ ચાસમાં આપવી. ❖ ચુસીયા જીવાતોના નિયંત્રણ માટે ઈમીડાકલોપ્રીડ ૬૦૦ એફ.એસ. %૪ મિલી/કિલો અથવા થાયામીથોકઝામ ૭૦ ડબલ્યુ.એસ. %૧ ગ્રામ/કિલો બીજ મુજબ બીજ માવજત આપવી. ❖ વાવણી લાયક પૂરતા વરસાદ થયે પાકનું વાવેતર સમયસર કરવું.
<p style="color: red;">કપાસ</p>	<p style="color: blue;">વાવેતર</p>	<p style="color: blue;">ગુજરાત હાઈબ્રીડ કપાસ- ૫, ૮, ૧૨ (બોલગાડ્-૨) કે બી.ટી. કપાસ / સુકારો અને મૂળનો કોહવારો</p>	<ul style="list-style-type: none"> ❖ કપાસ વાવણી માટે નવી બહાર પાડેલ વધુ ઉત્પાદન આપતી બીટી હાઈબ્રીડ કપાસ ગુજરાત હાઈબ્રીડ કપાસ-૫, ૮, ૧૨ (બોલગાડ્-૨) પસંદ કરવી કે બી.ટી. કપાસનું વિશ્વાખપાત્ર બીયારણ જ ખરીદવું. ❖ કપાસનું વિશ્વાખપાત્ર સર્ટિફિકેડ બીજ ખરીદવું. ❖ પાકની ફેરબદલી કરવી. કપાસની સાથે મઠનો આંતરપાક લેવો. ❖ ઈક્કડના પાકનો લીલો પડવાશ કરવો. ❖ સ્યુડોમોનાસ ફલુરોસન્સ ૧%, ૧૦ ગ્રામ અથવા કાર્બોકઝીન ૩૭.૫ % + થાયરમ ૩૭.૫ % ડી. એસ., ૩.૫ ગ્રામ પ્રતિ કીલો બીજને પટ આપવો. ❖ વાવણી લાયક પૂરતા વરસાદ થયે પાકનું વાવેતર સમયસર કરવું.

સોયાબીન	વાવેતર	ગુજ. સોયાબીન - ૧, ૨ & જીજેસ - ૩ / ચુસીયા જીવાતો, કોલાર રોડ, ચારકોલ રોડ	<ul style="list-style-type: none"> ❖ સોયાબીન પાકનાં વાવેતર માટે સારી ગુણવત્તા અને વધારે ઉત્પાદન આપતી જાતોનાં પ્રમાણિત બીજની પસંદગી કરીને મેળવી લેવું. ❖ પાકમાં વાવેતર સમયે પાચામાં એસએસપી ૩૭૫ કીલો અને ૬૫.૨ કીલો ચુરીયા/હે આપવું. ❖ ઇમીડાકલોપ્રીડ ૪૮ એક્સ.એસ. @૩ મીલી/કિગ્રા સાથે બીજ માવજત આપવી. ❖ ફલુકાપાચરોક્સેડ ૩૩૩ ગ્રામ/લી એક્સ.એસ ને ૧.૦ મીલી/કિલો અથવા કાબોકગીન ૩૭.૫% + થાયરમ ૩૭.૫% ડબલ્યુએસ ની ૩ ગ્રામ/કિલો બીચારણ મુજબ બીજ માવજત આપવી. ❖ વાવણી લાયક પૂરતા વરસાદ થયે પાકનું વાવેતર સમયસર કરવું.
આંબા	ફળ પરિપક્વતા	-	<ul style="list-style-type: none"> ❖ આંબાનાં બગીચામાં નિયમિત પીયત આપવું. ❖ શાખ પડે એટલે પરીપક્વ કેરી ઉતારી ગ્રેડીંગ કરી બજારમાં વહેંચાણ માટે મોકલવી.
શાકભાજુ	વૃદ્ધી / કુલ / ફળ	-	<ul style="list-style-type: none"> ❖ પાકમાં દવા છાંટવાનું અને ખાતર આપવનું કામ આ સમયગાળામાં કરવું. ❖ પાકમાં હાથથી નિંદામણ અને આંતરખેડની કામગીરી કરવી. ❖ શાકભાજુ પાકોને સમયસર પીયત આપવું. ❖ શાકભાજુના પાકોને નીદામણ મુક્ત કરવું અને નિયમિત પિયત આપવું. શાકભાજુ ને સમયસર ઉતારી ગ્રેડીંગ કરી બજારમાં મુકવા.
પશુપાલન	-	-	<ul style="list-style-type: none"> ❖ પશુઓને યોગ્ય આહાર આપવો & પીવા માટે યોખ્યું પાણી આપવું અને પશુઓના રહેઠાણ યોખ્યા રાખવા. ❖ માઝી મચ્છરથી રક્ષણ માટે ફીનાઈલ છાટવું. ❖ પશુઓને યોગ્ય આહારમાંખનિજ્યુક્ત મિશ્રણ સાથે સંગ્રહિત ચારા આપો. ચારા ને સલામત સ્થળે સંગ્રહિત કરવો જોઈએ. ❖ ચારા માટે ચાફ કટર વાપરો. ❖ પશુઓમાં ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટાઇ ૨ મીલી/ ૧ લીટર પાણીમાં નાખીને છાટવી. ❖ દુધાળા પશુઓના આઉને ઝીંક ઓક્સાઈડ અથલા બોરીક પાવડરથી બરાબર સાફ કરવા.



ગ્રામીણ કૃષિ મૌસમ સેવા
કૃષિ હવામાન સેલ
કૃષિ વિજ્ઞાન વિભાગ
જુનાગઢ કૃષિ યુનિવર્સિટી
જુનાગઢ - ૩૬૨૦૦૧
ઈમેઇલ:- amfujind@yahoo.com ફોન નં. **૦૨૮૫-૨૬૭૦૧૩૯**



હવામાન આગાહી આધારીત દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારનાં ગીર સોમનાથ જીલ્લા માટે કૃષિ સલાહ બુલેટીન
(કૃષિ હવામાન સેલ, જુનાગઢ કૃષિ યુનિવર્સિટી અને ભારત હવામાન વિભાગ દ્વારા સંપર્ક)

વાર:- મંગળવાર

તારીખ:- ૨૫/૦૬/૨૦૨૪

હવામાન આગાહી આધારીત બુલેટીન: ૦૨૫ (૨૦૨૪-૨૫)

પાછલા પાંચ દિવસનું હવામાન (તારીખ:- ૨૦ થી ૨૪ જૂન, ૨૦૨૪)					હવામાન પરિબળો	હવામાન આગાહી આગામી પાંચ દિવસ માટે (તારીખ:- ૨૬ થી ૩૦ જૂન, ૨૦૨૪) માન્ય ૩૦ જૂન, ૨૦૨૪. સવારના ૮:૩૦ વાગ્યાં સુધી				
૨૦/૦૬	૨૧/૦૬	૨૨/૦૬	૨૩/૦૬	૨૪/૦૬	તારીખ	૨૬/૦૬	૨૭/૦૬	૨૮/૦૬	૨૯/૦૬	૩૦/૦૬
૧.૩	૦	૦	૦	૦.૨	વરસાદ(મિમી)	૭૦	૨૦	૨૪	૨૨	૨૭
૩૩.૬	૩૩.૮	૩૪.૪	૩૪.૪	૩૪.૨	મહત્વમાન (ઓસે.ગ્રે)	૩૫	૩૪	૩૪	૩૨	૩૪
૩૦.૩	૩૦.૩	૩૦.૩	૨૬.૮	૨૭.૮	લઘુત્વમાન (ઓસે.ગ્રે)	૩૦	૨૬	૨૬	૨૬	૨૬
મોટેભાગે વાદળાયું	અંશતા: વાદળાયું	અંશતા: વાદળાયું	મોટેભાગે વાદળાયું	મોટેભાગે વાદળાયું	વાદળાની સ્થિતિ	વાદળાયું	મોટેભાગે વાદળાયું	વાદળાયું	વાદળાયું	વાદળાયું
૮૩	૮૪	૮૪	૮૭	૮૫	સવારના ભેજના ટકા	૮૨	૮૩	૮૧	૮૫	૮૫
૮૧	૭૬	૭૭	૭૨	૮૩	બપોરબાદ ના ભેજના ટકા	૬૬	૫૫	૫૩	૬૭	૭૨
૧૧.૧	૨૦.૪	૧૧.૧	૫.૬	-	પવનની ગતિ (કિમી/કલાક)	૨૨.૦	૧૬.૦	૨૩.૦	૨૦.૦	૨૫.૦
૨૫૦	૨૫૦	૨૭૦	૨૭૦	-	પવનની દિશા (ડિગ્રી)	૨૬૨	૨૪૬	૨૪૮	૨૪૮	૨૪૮

સામાન્ય સલાહ

ભારત સરકારશ્રીના હવામાન વિભાગ તરફથી મળેલ હવામાન આગાહીની માહીતી અનુસાર અત્રેના ગ્રામીણ કૃષિ મૌસમ સેવા યોજના, જુનાગઢ કૃષિ યુનિવર્સિટી, જુનાગઢ તરફથી જણાવવામા આવે છે કે દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના ગીર સોમનાથ જીલ્લામાં આગામી તારીખ ૨૬/૦૬/૨૦૨૪ થી ૩૦/૦૬/૨૦૨૪ દરમ્યાન આકાશ મુખ્યત્વે આગામી દિવસોમાં મુખ્યત્વે મોટેભાગે વાદળાયું થી વાદળાયું રહેશે, જીલ્લામાં આવતીકાલે ભારે વરસાદ અને પાંચ દિવસો દરમિયાન છુટા છવાયાં ભાગોમાં હળવા થી મધ્યમ વરસાદની શક્યતાઓ છે. પવનની ઝડપ ૧૮ થી ૨૫ કીમી/કલાક રહેવાની શક્યતાઓ રહેશે. આ સમયગાળામાં મહત્વમાન તાપમાન ૩૨ થી ૩૫ ડિગ્રી સેન્ટીગ્રેડ રહેશે તેમજ લઘુત્વમાન તાપમાન ૨૮ થી ૩૦ ડિગ્રી સેન્ટીગ્રેડ રહેવાની સંભાવના છે.

માન્ય વરસાદની સૂરી મુજબ ગોર્ખાથ જુલ્લામાં ૨૭ સપ્ટેમ્બર થી ૨૫ ઓક્ટોબર, ૨૦૨૩ નાં સમગ્રગતાની દરમિયાન મધ્યમ લેજની સ્થિતિ જોવા મળેલ છે.

સાપ્તાહિક હવામાન આગાહીની સૂરી મુજબ સૌરાષ્ટ્ર અને કચ્છનાં વિસ્તારોમાં તા. ૩૦/૦૬/૨૦૨૪ થી તા. ૦૬/૦૭/૨૦૨૪ દરમિયાન વરસાદ અને સામાન્ય થી ઉપર & મહત્વમાન અને લધુતમાં તાપમાન સામાન્ય રહેવાની શક્યતાઓ છે.

મેધદૂત એપ્લીકેશન:- કૃષિ હવામાન સલાહ સેવા અને હવામાન આગાહી અંગેની જાણકારી મેળવવા માટે ડાઉનલોડ કરો મેધદૂત એપ્લીકેશન કે જે ડેડોને તેમની સ્થાનિક ભાષામાં સ્થાન, પાક અને પશુધન વિશે હવામાન આધારિત કૃષિ સલાહ પ્રદાન કરશે.

મેધદૂત એપ્લીકેશનને ડાઉનલોડ કરવા માટે નીચેની લિંક પર ક્લિક કરો.

એન્ડ્રોઈડ(મોબાઇલ)ઉપયોગકર્તા: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>

આઈ.ઓ.એસ.ઉપયોગકર્તા: - <https://apps.apple.com/in/app/meghdoot/id1474048155>

✓ જો સૂકાયેલ કણસલાં કે કાપણી કરેલ ઉનાળું પાક/ધાસચારો ઘેતરમાં પડેલ હોય તો યોગ્ય જ્યાએ સંગ્રહ કરવો.


❖ ઘેતરમાંથી જમીનના નમુના લેવા અને જમીન પૃથ્વકરણ કરવા માટે લેબોરેટરીમાં મોકલી આપવા.

❖ ઘેતરના સેઢા, પાળા અને વરસાદના પાણી નિકાલની ગટર ને ચોખ્માં કરી ને આગામી ચોમાસા માટે ઘેતર ને તૈયાર કરવું.

❖ ક્રોમ્પોસ્ટ, દેશી ખાતર અને જુખ્સમ ભલામણ મુજબ ચાસમાં ભરવા.

❖ આગામી ચોમાસાની ઝતુમાં વાવણી માટે ડેડોને સારી ગુણવત્તા અને વધારે ઉત્પાદન આપતી જતોના પ્રમાણિત બીજની પસંદગી કરીને મેળવી લેવાની સલાહ આપવામાં આવે છે.

❖ વાવણી લાયક વરસાદ થયે ચોમાસું પાકનું વાવેતર કરવું.



પાકનું નામ	પાક અવસ્થા	રોગ/જીવાત/ જાત	હૃદિ સલાહ
ચોમાસું મગફળી	વાવેતર / ઉગાવો	અર્ધવેલડી જાત- જુજુ.૨૦ અથવા ઉલભી જાતો જુજુ.૨ જુજુ.૫ જુજુ.૭ ટી.જુ. ૨૬ / મુંડા, ચુસીયા જીવાતો, ઉગસુક, થડનો કોહવારો	<ul style="list-style-type: none"> ❖ વાવેતર માટે અર્ધવેલડી જાત- જુજુ.૨૦ અથવા ઉલભી જાતો જુજુ.૨ જુજુ.૫ જુજુ.૭ ટી.જુ. ૨૬ ની પસંદગી કરવી. ❖ ચોમાસું મગફળીના પાકને હેકટરે ૧૨.૫ કિ.ગ્રા નાઇટ્રોજન અને ૧૫ કિ.ગ્રા ફોસ્ફરસ ખાતર તરીકે વાવણી વખતે ચાસમાં ઓરીને આપવા. ❖ વાવેતર સમયે ડીએપી ૫૪.૩ કીલો અને ૬ કીલો યુરીયા/હે. અને એરેનીનો ખોળ ૫૦૦ કી.ગ્રા./હે. અથવા જુખ્સમ ૫૦૦ કી.ગ્રા./હે. સાથે એરેનીનો ખોળ ૫૦૦ કી.ગ્રા./હે પાયામા આપવો. ❖ વાવતી વખતે બીયારણને થાયરમ અથવા મેંકોઝેબ અથવા વીટાવેક્ષ દવાનો ૨-૩ ગ્રામ/કી. બીજને પટ આપવો. ❖ મુંડાના નિયંત્રણ માટે કલોરપાયરીફોસ ૨૦ % છ.સી. ૧૫ મિલી/કિલો બીજ અથવા ઈમીડાકલોપ્રીડ ૬૦૦ % એફ.એસ. ૪ મિલી/કિલો બીજ મુજબ બીજ માવજત આપવી. ❖ બીવેરિયા બાસીયાના અથવા મેટારીઝીયમ એનીસોપ્ટી નામની ફૂગનો પાવડર વાવેતર પહેલા ૫ કિલો પ્રતિ હેકટર મુજબ એરેનીના ખોળ સાથે ચાસમાં આપવાથી મુંડાનું નિયંત્રણ કરી શકાય છે. અથવા કાર્બોફ્યુરાન ૩૪ નામની દાણાદાર દવાને ૩૩ કિલો/હેકટર મુજબ ચાસમાં આપવી. ❖ ચુસીયા જીવાતોના નિયંત્રણ માટે ઈમીડાકલોપ્રીડ ૬૦૦ % એફ.એસ. ૪ મિલી /કિલો અથવા થાયામીથોકામ ૭૦ % ડબલ્યુ.એસ. ૧ ગ્રામ / કિલો બીજ મુજબ બીજ માવજત આપવી. ❖ વાવણી લાયક પૂરતા વરસાદ થયે પાકનું વાવેતર સમયસર કરવું.
કપાસ	વાવેતર	ગુજરાત હાઈબ્રીડ કપાસ- ૬, ૮, ૧૨ (બોલગાઈ-૨) કે બી.ટી. કપાસ / સુકારો અને મૂળનો કોહવારો	<ul style="list-style-type: none"> ❖ કપાસ વાવણી માટે નવી બહાર પાડેલ વધુ ઉત્પાદન આપતી બીટી હાઈબ્રીડ કપાસ ગુજરાત હાઈબ્રીડ કપાસ-૬, ૮, ૧૨ (બોલગાઈ-૨) પસંદ કરવી કે બી.ટી. કપાસનું વિશ્વાષપાત્ર બીયારણ જ ખરીદવું. ❖ કપાસનું વિશ્વાષપાત્ર સર્ટિફિકેડ બીજ ખરીદવું. ❖ પાકની ફેરબદલી કરવી. કપાસની સાથે મઠનો આંતરપાક લેવો. ❖ ઈક્કડના પાકનો લીલો પડવાશ કરવો.

			<ul style="list-style-type: none"> ❖ સ્યુડોમોનાસ ફલુરોસન્સ ૧%, ૧૦ ગ્રામ અથવા કાબોકઝીન ૩૭.૫ % + થાયરમ ૩૭.૫ % ડી. એસ., ૩.૫ ગ્રામ પ્રતિ કીલો બીજને પટ આપવો. ❖ વાવણી લાયક પૂરતા વરસાદ થયે પાકનું વાવેતર સમયસર કરવું.
સોયાબીન	વાવેતર	ગુજ. સોયાબીન - ૧, ૨ & જીજેએસ - ૩ / ચુસીયા જીવાતો, કોલાર રોટ, ચારકોલ રોટ	<ul style="list-style-type: none"> ❖ સોયાબીન પાકનાં વાવેતર માટે સારી ગુણવત્તા અને વધારે ઉત્પાદન આપતી જાતોનાં પ્રમાણિત બીજની પસંદગી કરીને મેળવી દેવું. ❖ પાકમાં વાવેતર સમયે પાયામાં એસએસપી ૩૭૫ કીલો અને ૬૫.૨ કીલો યુરીયા/હે આપવું. ❖ ઈમીડાકલોપ્રીડ ૪૮ એફ.એસ. @૩ મિલી/કિગ્રા સાથે બીજ માવજત આપવી. ❖ ફલુકઅપાયરોક્સેડ ૩૩૩ ગ્રામ/લી એફએસ ને ૧.૦ મિલી/કિલો અથવા કાબોકઝીન ૩૭.૫% + થાયરમ ૩૭.૫% ડબલ્યુએસ ની ૩ ગ્રામ/કિલો બીયારણ મુજબ બીજ માવજત આપવી. ❖ વાવણી લાયક પૂરતા વરસાદ થયે પાકનું વાવેતર સમયસર કરવું.
અંબા	ફળ પરિપક્વતા	-	<ul style="list-style-type: none"> ❖ અંબાનાં બગીચામાં નિયમિત પીયત આપવું. ❖ શાખ પડે એટલે પરીપક્વ કેરી ઉતારી ગ્રેડિંગ કરી બજારમાં વહેંચાણ માટે મોકલવી.
શાકભાજુ	દુંધણી / કુલ / ફળ	-	<ul style="list-style-type: none"> ❖ પાકમાં દવા છાંટવાનું અને ખાતર આપવનું કામ આ સમયગાળામાં કરવું. ❖ પાકમાં હાથથી નિંદામણની કામગીરી કરવી. ❖ શાકભાજુ પાકોને સમયસર પીયત આપવું. ❖ શાકભાજુના પાકોને નીદામણ મુકત કરવું અને નિયમિત પિયત આપવું. શાકભાજુ ને સમયસર ઉતારી ગ્રેડિંગ કરી બજારમાં મુકવા.
પશુપાલન	-	-	<ul style="list-style-type: none"> ❖ પશુઓને યોગ્ય આહાર આપવો & પીવા માટે ચોખ્યું ઠંડું પાણી આપવું અને પશુઓના રહેઠાણ ચોખ્યા રાખવા. ❖ માખી મચ્છરથી રક્ષણ માટે ફીનાઈલ છાટવું. ❖ પશુઓને યોગ્ય આહારમાં ખનિજયુક્ત મિશ્રણ સાથે સંગ્રહિત ચારા આપો. ચારા ને સલામત સ્થળે સંગ્રહિત કરવો જોઈએ. ❖ ચારા માટે ચાઙ કટર વાપરો.

- ❖ પશુઓમાં ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટાજ ૨ મીલી/ ૧ લીટર પાણીમાં નાખીને છાટવી.
- ❖ દુધાળા પશુઓના આઉને ઝીંક ઓક્સાઇડ અથલા બોરીક પાવડરથી બરાબર સાફ કરવા.

યાંત્રિક અધિકારી
ગ્રામીણ કૃષિ મૌસમ સેવા
જ.કૃ.યુ.

Gramin Krushi Mausam Seva (GKMS)
Agrometeorological Cell
Department of Agronomy
Junagadh Agricultural University
Junagadh-362001

E-mail: amfujnd@yahoo.com Phone: 0285-2670139



Agromet Advisory Service Bulletin for Junagadh District of South Saurashtra Agro-Climatic Zone
(Issued jointly by Junagadh Agricultural University and India Meteorological Department)

Day:- Tuesday

Date: 25-06-2024

Agromet Advisory Service bulletin. No:- 025 (2024-25)

Observed Weather during last Five days (Dated 20 th to 24 th June, 2024)					Weather Parameters	Weather Forecast upcoming Five days (Valid for 26 th to 30 th June, 2024) <u>until 08:30 hrs of 30-06-2024</u>				
20/06	21/06	22/06	23/06	24/06	Date	26/06	27/06	28/06	29/06	30/06
0	0	0	0	102.9	Rainfall(mm)	79	13	19	31	44
38.5	38.3	38.0	38.7	39.9	T _{max} (°C)	33	33	32	32	33
29.1	29.6	29.4	29.3	24.5	T _{min} (°C)	25	24	24	24	24
3	3	3	4	8	Cloud Cover	8	7	8	8	8
79	76	78	78	85	RH-I (%)	82	81	79	82	84
52	49	45	48	79	RH-II (%)	62	49	58	60	73
14.9	18.2	15.3	9.1	5.2	Wind speed (km/hr)	22.0	17.0	21.0	20.0	24.0
230	250	270	250	290	Wind direction	292	248	248	248	248

Agro meteorological Advisories

General Advice	Rain fall	As per IMD Forecast, There is forecasting of very heavy rainfall tomorrow and light to moderate rainfall occurs during next five days. Sky will be remaining mainly cloudy to cloudy.
	Temperature	Maximum and minimum temperature likely to be in range of 32 - 33°C and 24 - 25°C , respectively.
	RH (%)	The Relative Humidity is likely to be in the range of 49 – 84 % .
	Wind Direction and Wind Speed	Direction and speed of wind is to be mainly W and 17 – 24 km/h , respectively.

■ **Standardized Precipitation Index:** For the period of ~~28th September to 25th October, 2023.~~ Junagadh district considered under the mildly wet condition.

■ As per ERFS products, **above normal** rainfall and T_{\max} . & T_{\min} . **normal** may occur in Saurashtra & Kutch region during **30th June to 06th July, 2024.**


■ **MEGHDOOT Mobile Application**:- Download the MEGHDOOT mobile application for Agromet Advisory Services and information on weather forecasting, which will provide weather based agricultural advice to farmers for their location, crops and livestock in their local language.

Click on the following link to download the MEGHDOOT application:

Android users: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>

IOS users: <https://apps.apple.com/in/app/meghdoot/id1474048155>

- ✓ If dried hay or harvested summer crop / fodder is lying in the field, store it in the right place.
- ❖ To take the soil samples from the field and send to the soil testing laboratory for the soil analysis.
- ❖ Cleaning ridge, fencing and drainage system of the field and preparing the field for the next monsoon.
- ❖ Compost, FYM fertilizers and gypsum filling in the furrow as recommended.
- ❖ Farmers are advised to procure good quality and select the certified seed with high yielding variety of Kharif crops for sowing next monsoon season.
- ❖ Timely sowing should be done after sufficient rainfall.



Crop	Variety/ Pest/Insect	Stage	Agro met Advisories
Kharif Groundnut	GG.20 or erect GG.2, GG.5, GG.7 and TG-26 / white grub, sucking pests, Collar rot, stem rot	Sowing / Germination	<ul style="list-style-type: none"> ➤ For pre-sowing of Kharif groundnut select semi-spread GG.20 or erect GG.2, GG.5, GG.7 and TG-26 varieties. ➤ In Kharif groundnut apply 12.5 kg N/ha and 25 kg P/ha at the time of sowing. ➤ Apply 54.3 kg/ha DAP, 6 kg/ha Urea and 500 kg/ha Castor cake as basal dose. ➤ Seed should be treatment with Dithane M-45/ mencozeb/vitavax @ 2 - 3 grams per kg of seeds before sowing. ➤ Seed treatment with chlorpyriphos 20EC @ 25 ml/kg seed or imidacloprid 600 FS @ 4 ml/kg seed for white grub. ➤ Soil application of Beauveria bassiana or Metarizium anisopliae 1.15 WP @ 5 kg/ha (Min. 2 x 10⁶ cfu/g) along with castor cake (300 kg/ha) before sowing for white grub. or Soil application of carbofuran 3 % CG @ 33 kg/ha for white grub.

			<ul style="list-style-type: none"> ➤ Seed treatment with imidacloprid 600 FS @ 4 ml/kg seed or thiamethoxam 70 WS @ 1 g/kg seed for sucking pests. ➤ Timely sowing should be done after sufficient rainfall.
Cotton	Gujarat Hybrid Cotton-6, 8, 12' (Bolegaurd-2) or Govt. approved varieties of B.T. cotton / Wilt and root rot	Sowing	<ul style="list-style-type: none"> ➤ Farmers are advised to procure the seed of high yielding variety of BT hybrid Cotton 'Gujarat Hybrid Cotton-6, 8, 12' (Bolegaurd-2) or Govt. approved varieties of B.T. cotton for sowing. ➤ Select the certified seed of cotton for cultivation. ➤ Follow crop rotation. Inter cropping with Vigna aconitifolia. ➤ Green manuring with Sesbania acubeata. ➤ Treat the seeds with Pseudomonas fluorescens 1% (10 gm/ kg seed) or Carboxin 37.5% + thiram 37.5% DS @ 3.5 g/Kg seed. ➤ Timely sowing should be done after sufficient rainfall.
Soyabean	Guj. Soyabean – 1, 2 & GJS – 3 / Sucking pests, Collar rot, Charcoal rot	Sowing	<ul style="list-style-type: none"> ➤ To procure good quality and select the certified seed with high yielding variety of soyabean crop. ➤ In soybean apply 375 kg SSP/ha & 65.2 kg urea/ha at the time of sowing. ➤ Seed treatment with Imidacloprid 48 FS @ 3 ml/kg seed.

			<ul style="list-style-type: none"> ➤ Treat the seeds with Fluxapyroxad 333 g/l FS 1.0 ml/ kg seeds or Carboxin 37.5% + Thiram 37.5% WS 2.25 g/Kg seed. ➤ Timely sowing should be done after sufficient rainfall.
Mango	-	Fruit maturity	<ul style="list-style-type: none"> ➤ Apply irrigation regularly in mango orchard. ➤ To pick the mango fruit at physiological maturity stage and send it to the market often grading for selling.
Vegetable crops	-	Sowing/ Flowering/ Fruiting	<ul style="list-style-type: none"> ➤ To apply fertilizer and any agro-chemicals to the crops during this period. ➤ To the spray of agrochemicals during this period. ➤ Hand weeding and interculturing operation should be carry out. ➤ To apply irrigation to the crop at regular interval. ➤ Keep vegetable crop weed free, apply irrigation regularly and send vegetable fruit to the market often grading.
Livestock	-	-	<ul style="list-style-type: none"> ➤ To provide adequate food & clean cool water for drinking and keep the animals clean shed. ➤ Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Give stored fodder with mineral mixture. Fodder should be stored at safe place. ➤ Use chaff cutter for fodder. ➤ For the control of tick and other parasites, spray deltemethrin or aemitaz 2 ml in 1 litre of water. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.



Gramin Krushi Mausam Seva (GKMS)
Agrometeorological Cell
Department of Agronomy
Junagadh Agricultural University
Junagadh-362001

E-mail: amfujnd@yahoo.com Phone: 0285-2670139



Agromet Advisory Service Bulletin for Porbandar District of South Saurashtra Agro-Climatic Zone

(Issued jointly by Junagadh Agricultural University and India Meteorological Department)

Day:- Tuesday

Date: 25-06-2024


Agromet Advisory Service bulletin. No:- 025 (2024-25)

Observed Weather during last Five days (Dated 20 th to 24 th June,2024)					Weather Parameters	Weather Forecast upcoming Five days (Valid for 26 th to 30 th June,2024)				
20/06	21/06	22/06	23/06	24/06	Date	26/06	27/06	28/06	29/06	30/06
0	0	0	0	4.9	Rainfall(mm)	14	2	8	19	20
34.8	35.0	35.2	35.4	35.6	Tmax (°C)	33	34	34	34	34
29.8	30.0	29.8	28.6	26.0	Tmin (°C)	29	29	29	29	29
4	4	2	7	7	Cloud Cover	8	6	7	8	8
76	74	75	75	90	RH-I (%)	81	80	79	80	83
67	74	72	74	72	RH-II (%)	69	55	64	65	71
18.5	24.1	16.7	11.1	7.4	Wind speed (km/hr)	28.0	19.0	23.0	24.0	28.0
250	250	270	320	50	Wind direction	288	248	248	248	248

Agro meteorological Advisories

General Advice	Rain fall	As per IMD Forecast, There is forecasting of light to moderate rainfall occurs at isolated places during next five days. Sky will be remaining mainly cloudy to cloudy.
	Temperature	Maximum and minimum temperature likely to be in range of 33 - 34°C and 29°C , respectively.
	RH (%)	The Relative Humidity is likely to be in the range of 55 – 83 % .
	Wind Direction and Wind Speed	Direction and speed of wind is to be mainly W and 19 – 28 km/h , respectively.

- ❖ **Standardized Precipitation Index:** For the period **28th September to 25th October, 2023.**
Porbandar district considered under the mildly dry condition.
- ❖ **As per ERFS products, above normal rainfall and T_{max.} & T_{min.} normal may occur in Saurashtra & Kutch region during 30th June to 06th July, 2024.**
- ❖ **MEGHDOOT Mobile Application:** Download the MEGHDOOT mobile application for Agromet Advisory Services and information on weather forecasting, which will provide weather based agricultural advice to farmers for their location, crops and livestock in their local language.
Click on the following link to download the MEGHDOOT application:
Android users:<https://play.google.com/store/apps/details?id=com.aas.meghdoot>
IOS users:<https://apps.apple.com/in/app/meghdoot/id1474048155>
- ✓ If dried hay or harvested summer crop / fodder is lying in the field, store it in the right place.
- ❖ Take the soil samples from the field and send to the soil testing laboratory for the soil analysis.
- ❖ Cleaning ridge, fencing and drainage system of the field and preparing the field for the next monsoon.
- ❖ Compost, FYM fertilizers and gypsum filling in the furrow as recommended.
- ❖ Farmers are advised to procure good quality and select the certified seed with high yielding variety of Kharif crops for sowing next monsoon season.
- ❖ Timely sowing should be done after sufficient rainfall.



Crop	Variety/ Pest / Insect	Stage	Agro met Advisories
------	---------------------------	-------	---------------------

Kharif Groundnut	GG.20 or erect GG.2, GG.5, GG.7 and TG- 26 / white grub, sucking pests, Collar rot, stem rot	Sowing / Germination	<ul style="list-style-type: none"> ➤ For pre-sowing of Kharif groundnut select semi-spread GG.20 or erect GG.2, GG.5, GG.7 and TG-26 varieties. ➤ In Kharif groundnut apply 12.5 kg N/ha and 25 kg P/ha at the time of sowing. ➤ Apply 54.3 kg/ha DAP, 6 kg/ha Urea and 500 kg/ha Castor cake as basal dose. ➤ Seed should be treatment with Dithane M-45/ mencozeb/vitavax @ 2 - 3 grams per kg of seeds before sowing. ➤ Seed treatment with chlorpyriphos 20EC @ 25 ml/kg seed or imidacloprid 600 FS @ 4 ml/kg seed for white grub. ➤ Soil application of Beauveria bassiana or Metarizium anisopliae 1.15 WP @ 5 kg/ha (Min. 2 x 106 cfu/g) along with castor cake (300 kg/ha) before sowing for white grub. or Soil application of carbofuran 3 % CG @ 33 kg/ha for white grub. ➤ Seed treatment with imidacloprid 600 FS @ 4 ml/kg seed or thiamethoxam 70 WS @ 1 g/kg seed for sucking pests. ➤ Timely sowing should be done after sufficient rainfall.
Cotton	Gujarat Hybrid Cotton-6, 8, 12' (Bolegaurd-2) or Govt.	Sowing	<ul style="list-style-type: none"> ➤ Farmers are advised to procure the seed of high yielding variety of BT hybrid Cotton 'Gujarat Hybrid Cotton-6, 8, 12' (Bolegaurd-2) or Govt. approved varieties of B.T. cotton for sowing.

	approved varieties of B.T. cotton / Wilt and root rot		<ul style="list-style-type: none"> ➤ Select the certified seed of cotton for cultivation. ➤ Follow crop rotation. Inter cropping with Vigna aconitifolia. ➤ Green manuring with Sesbania acubeata. ➤ Treat the seeds with Pseudomonas fluorescens 1% (10 gm/ kg seed) or Carboxin 37.5% + thiram 37.5% DS @ 3.5 g/Kg seed. ➤ Timely sowing should be done after sufficient rainfall.
Soyabean	Guj. Soyabean – 1, 2 & GJS – 3 / Sucking pests, Collar rot, Charcoal rot	Sowing	<ul style="list-style-type: none"> ➤ To procure good quality and select the certified seed with high yielding variety of soyabean crop. ➤ In soybean apply 375 kg SSP/ha & 65.2 kg urea/ha at the time of sowing. ➤ Seed treatment with Imidacloprid 48 FS @ 3 ml/kg seed. ➤ Treat the seeds with Fluxapyroxad 333 g/l FS 1.0 ml/ kg seeds or Carboxin 37.5% + Thiram 37.5% WS 2.25 g/Kg seed. ➤ Timely sowing should be done after sufficient rainfall.
Mango	-	Fruit maturity	<ul style="list-style-type: none"> ➤ Apply irrigation regularly in mango orchard. ➤ To pick the mango fruit at physiological maturity stage and send it to the market often grading for selling.
Vegetable crops	-	Sowing/ Flowering/ Fruiting	<ul style="list-style-type: none"> ➤ To apply fertilizer and any agro-chemicals to the crops during this period. ➤ To the spray of agrochemicals during this period.

			<ul style="list-style-type: none"> ➤ Hand weeding and interculturing operation should be carry out. ➤ To apply irrigation to the crop at regular interval. ➤ Keep vegetable crop weed free, apply irrigation regularly and send vegetable fruit to the market often grading.
Livestock	-	-	<ul style="list-style-type: none"> ➤ To provide adequate food & clean cool water for drinking and keep the animals clean shed. ➤ Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Give stored fodder with mineral mixture. Fodder should be stored at safe place. ➤ Feed silage & hay material along with concentrate feed. ➤ Use chaff cutter for fodder. ➤ For the control of tick and other parasites, spray deltemethrin or aemitaz 2 ml in 1 litre of water. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.

  आजादी का अमृत महोत्सव	<p>Gramin Krushi Mausam Seva (GKMS) Agrometeorological Cell Department of Agronomy Junagadh Agricultural University Junagadh-362001</p> <p>E-mail: amfujnd@yahoo.com Phone: 0285-2670139</p>	 
--	---	--

Agromet Advisory Service Bulletin for Gir Somnath District of South Saurashtra Agro-Climatic Zone
(Issued jointly by Junagadh Agricultural University and India Meteorological Department)

Day:- Tuesday

Date: 25-06-2024

Agromet Advisory Service Bulletin. No:- 025 (2024-25)

Observed Weather during last Five days (Dated 20 th to 24 th June,2024)					Weather Parameters	Weather Forecast upcoming Five days (Valid for 26 th to 30 th June,2024) until 08:30 hrs of 30-06-2024				
20/06	21/06	22/06	23/06	24/06	Date	26/06	27/06	28/06	29/06	30/06
1.3	0	0	0	0.2	Rainfall(mm)	70	20	24	22	27
33.6	33.8	35.4	35.4	34.2	Tmax (°C)	35	34	34	32	34
30.3	30.3	30.3	29.8	27.8	Tmin (°C)	30	29	29	29	29
6	4	4	6	7	Cloud cover	8	6	8	8	8
83	84	84	87	85	RH-I (%)	82	83	81	85	85
81	79	77	72	83	RH-II (%)	66	55	63	67	72
11.1	20.4	11.1	5.6	-	Wind speed (km/hr)	22.0	19.0	23.0	20.0	25.0
250	250	270	270	-	Wind direction	292	249	248	248	248

Agro meteorological Advisories

General Advice	Rain fall	As per IMD Forecast, There is forecasting of very heavy rainfall tomorrow and light to moderate rainfall occurs during next five days. Sky will be remaining mainly cloudy to cloudy.
	Temperature	Maximum and minimum temperature likely to be in range of 32 – 35°C and 29 -30°C , respectively.
	RH (%)	The Relative Humidity is likely to be in the range of 55 – 85 % .
	Wind Direction and Wind Speed	Direction and speed of wind is to be mainly W and around 19 – 25 km/h , respectively.

+ **Standardized Precipitation Index:** For the period **28th September to 25th October, 2023**.

Gir Somnath district considered under the mildly wet condition.

+ **As per ERFS products, above normal rainfall and T_{max}. & T_{min}. normal may occur in Saurashtra & Kutch region during 30th June to 06th July, 2024.**


+ **MEGHDOOT Mobile Application:** Download the MEGHDOOT mobile application for Agromet Advisory Services and information on weather forecasting, which will provide weather based agricultural advice to farmers for their location, crops and livestock in their local language.

Click on the following link to download the MEGHDOOT application:

Android users: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>

IOS users: <https://apps.apple.com/in/app/meghdoot/id1474048155>

- ✓ If dried hay or harvested summer crop / fodder is lying in the field, store it in the right place.
- ❖ To take the soil samples from the field and send to the soil testing laboratory for the soil analysis.
- ❖ Cleaning ridge, fencing and drainage system of the field and preparing the field for the next monsoon.
- ❖ Compost, FYM fertilizers and gypsum filling in the furrow as recommended.
- ❖ Farmers are advised to procure good quality and select the certified seed with high yielding variety of Kharif crops for sowing next monsoon season.
- ❖ Timely sowing should be done after sufficient rainfall.



Crop	Variety/ Pest / Insect	Stage	Agro met Advisories
Kharif Groundnut	GG.20 or erect GG.2, GG.5, GG.7 and TG-26 / white grub, sucking pests, Collar rot, stem rot	Sowing / Germination	<ul style="list-style-type: none"> ➤ For pre-sowing of Kharif groundnut select semi-spread GG.20 or erect GG.2, GG.5, GG.7 and TG-26 varieties. ➤ In Kharif groundnut apply 12.5 kg N/ha and 25 kg P/ha at the time of sowing. ➤ Apply 54.3 kg/ha DAP, 6 kg/ha Urea and 500 kg/ha Castor cake as basal dose.

			<ul style="list-style-type: none"> ➤ Seed should be treatment with Dithane M-45/mencozeb/vitavax @ 2 - 3 grams per kg of seeds before sowing. ➤ Seed treatment with chlorpyriphos 20EC @ 25 ml/kg seed or imidacloprid 600 FS @ 4 ml/kg seed for white grub. ➤ Soil application of Beauveria bassiana or Metarizium anisopliae 1.15 WP @ 5 kg/ha (Min. 2 x 106 cfu/g) along with castor cake (300 kg/ha) before sowing for white grub. or Soil application of carbofuran 3 % CG @ 33 kg/ha for white grub. ➤ Seed treatment with imidacloprid 600 FS @ 4 ml/kg seed or thiamethoxam 70 WS @ 1 g/kg seed for sucking pests. ➤ Timely sowing should be done after sufficient rainfall.
Cotton	Gujarat Hybrid Cotton-6, 8, 12' (Bolegaurd-2) or Govt. approved varieties of B.T. cotton / Wilt and root rot	Sowing	<ul style="list-style-type: none"> ➤ Farmers are advised to procure the seed of high yielding variety of BT hybrid Cotton 'Gujarat Hybrid Cotton-6, 8, 12' (Bolegaurd-2) or Govt. approved varieties of B.T. cotton for sowing. ➤ Select the certified seed of cotton for cultivation. ➤ Follow crop rotation. Inter cropping with Vigna aconitifolia. ➤ Green manuring with Sesbania acubeata.

			<ul style="list-style-type: none"> ➤ Treat the seeds with Pseudomonas fluorescens 1% (10 gm/ kg seed) or Carboxin 37.5% + thiram 37.5% DS @ 3.5 g/Kg seed. ➤ Timely sowing should be done after sufficient rainfall.
Soyabean	Guj. Soyabean – 1, 2 & GJS – 3 / Sucking pests, Collar rot, Charcoal rot	Sowing	<ul style="list-style-type: none"> ➤ To procure good quality and select the certified seed with high yielding variety of soyabean crop. ➤ In soybean apply 375 kg SSP/ha & 65.2 kg urea/ha at the time of sowing. ➤ Seed treatment with Imidacloprid 48 FS @ 3 ml/kg seed. ➤ Treat the seeds with Fluxapyroxad 333 g/l FS 1.0 ml/ kg seeds or Carboxin 37.5% + Thiram 37.5% WS 2.25 g/Kg seed. ➤ Timely sowing should be done after sufficient rainfall.
Mango	-	Fruit maturity	<ul style="list-style-type: none"> ➤ Apply irrigation regularly in mango orchard. ➤ To pick the mango fruit at physiological maturity stage and send it to the market often grading for selling.
Vegetable crops	-	Sowing/ Flowering/ Fruiting	<ul style="list-style-type: none"> ➤ To apply fertilizer and any agro-chemicals to the crops during this period. ➤ To the spray of agrochemicals during this period. ➤ Hand weeding and interculturing operation should be carry out. ➤ To apply irrigation to the crop at regular interval.

			<ul style="list-style-type: none"> ➤ Keep vegetable crop weed free, apply irrigation regularly and send vegetable fruit to the market often grading.
Livestock	-	-	<ul style="list-style-type: none"> ➤ To provide adequate food & clean cool water for drinking and keep the animals clean shed. ➤ Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Give stored fodder with mineral mixture. Fodder should be stored at safe place. ➤ Feed silage & hay material along with concentrate feed. ➤ Use chaff cutter for fodder. ➤ For the control of tick and other parasites, spray deltemethrin or aemitzaz 2 ml in 1 litre of water. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.

Technical Officer
 Gramin Krishi Mausam Seva (GKMS)
 J.A.U., Junagadh