

પત્રક-અ

હવામાન આગાહી આધારીત કૃષિ સલાહ બુદેટીન

(સમય તા : ૨૬.૦૭.૧૪ થી તા : ૩૦.૦૭.૧૪)

પાછળા અઠવાડીયાનું હવામાન

હવામાન પરીબળો	કૃષિ આબોહવાક્ય વિસ્તાર દક્ષિણ સૌરાષ્ટ્ર
વરસાદ (મીમી)	087.0
મહત્વમાન તાપમાન (સે.ગે.)	32.0
લઘુત્વમાન તાપમાન (સે.ગ્રે.)	23.8
ભેજ(%) (સવાર/બપોરબાદ)	94-83
પવનની ગતિ (કિમી/કલાક)	11.7

પત્રક-૬ : જિલ્લાવાર હવામાન આગાહી (તા. ૩૦.૦૭.૧૪ સુધી)

હવામાન પરીબળોઅ	તારીખ	દક્ષિણ સૌરાષ્ટ્ર	
		જુનાગઢ	પોરબંદર
વરસાદ (મીમી)	૨૬.૭.૧૪	6	5
	૨૭.૭.૧૪	10	6
	૨૮.૭.૧૪	5	3
	૨૯.૭.૧૪	0	3
	૩૦.૭.૧૪	0	3
મહત્વમાન તાપમાન (સે.ગ્રે.)	૨૬.૭.૧૪	28	28
	૨૭.૭.૧૪	28	28
	૨૮.૭.૧૪	29	29
	૨૯.૭.૧૪	29	29
	૩૦.૭.૧૪	29	29
લઘૃત્મ તાપમાન (સે.ગ્રે.)	૨૬.૭.૧૪	24	26
	૨૭.૭.૧૪	24	26
	૨૮.૭.૧૪	24	27
	૨૯.૭.૧૪	25	27
	૩૦.૭.૧૪	25	27
વાદળની સ્થિતિ (ઓક્ટો)	૨૬.૭.૧૪	મોટેભાગે વાદળછાયુ	મોટેભાગે વાદળછાયુ
	૨૭.૭.૧૪	અશંત: વાદળછાયુ	અશંત: વાદળછાયુ
	૨૮.૭.૧૪	મોટેભાગે વાદળછાયુ	મોટેભાગે વાદળછાયુ
	૨૯.૭.૧૪	મોટેભાગે વાદળછાયુ	મોટેભાગે વાદળછાયુ
	૩૦.૭.૧૪	મોટેભાગે વાદળછાયુ	મોટેભાગે વાદળછાયુ
મહત્વમાન ભેજ (%)	૨૬.૭.૧૪	78	78
	૨૭.૭.૧૪	80	80
	૨૮.૭.૧૪	80	80
	૨૯.૭.૧૪	79	79
	૩૦.૭.૧૪	81	81
લઘૃત્મ ભેજ (%)	૨૬.૭.૧૪	67	66
	૨૭.૭.૧૪	71	70
	૨૮.૭.૧૪	72	71
	૨૯.૭.૧૪	74	75
	૩૦.૭.૧૪	77	77
પવનની ગતિ (કિગ્રી/કલાક)	૨૬.૭.૧૪	027	025
	૨૭.૭.૧૪	031	027
	૨૮.૭.૧૪	035	032
	૨૯.૭.૧૪	039	036
	૩૦.૭.૧૪	036	033
પવનની દિશા (ડીગ્રી)	૨૬.૭.૧૪	240	240
	૨૭.૭.૧૪	250	250
	૨૮.૭.૧૪	250	250
	૨૯.૭.૧૪	240	240
	૩૦.૭.૧૪	240	240

પત્રક-ક

જિલ્લાવાર કૃષિ સલાહ

પાકનું નામ	કૃષિ સલાહ
દક્ષિણ ભેત આબોહવાક્ય વિસ્તાર	<p>ચોમાસુ મગફળી</p> <ul style="list-style-type: none"> ❖ વરાપના સમયે પાકમાં જરૂરીયાત મુજબ સમયાંતરે આતરખેડ અને નિંદામણ જેવી કામગીરી હાથ ધરવી. ❖ વાદળછાયા વાતાવરણના લીધે પાકમાં ઉગસુકનો રોગ જોવા મળે શકે છે તો તેના નિયંત્રણ માટે બીજ ને થાયરમ નો પટ આપવો અથવા જમીનમાં ક કીગ્રાહે. ટ્રાઇકોડરમા પાવડરનો છંટકાવ કરવો. ❖ મગફળી પીળી પડી જાય તો ૧૦૦ ગ્રામ ફેરસ સલ્ફેટ અને ૧૦ ગ્રામ લીન્યુના કુલને ૧૦ લીટર પાણીમાં ઓગાળીને છંટકાવ કરવો. જરૂર પડે તો ૮ થી ૧૦ દિવસે ફરી છટકાવ કરવો.
	<p>કપાસ</p> <ul style="list-style-type: none"> ❖ વરાપના સમયે પાકમાં જરૂરીયાત મુજબ સમયાંતરે આતરખેડ અને નિંદામણ જેવી કામગીરી હાથ ધરવી. ❖ આગોત્ર વાવેતર કરેલ કપાસમાં યુરીયા ૨૫ કિ.ગ્રા અને અમોનીયમ સલ્ફેટ ૨૫ કિ.ગ્રા. જમીનમાં ભેજ હોય ત્યારે આપવુ.
	<p>સોયાબીન</p> <ul style="list-style-type: none"> ❖ વરાપના સમયે પાકમાં જરૂરીયાત મુજબ સમયાંતરે આતરખેડ અને નિંદામણ જેવી કામગીરી હાથ ધરવી.
શાકભાજુ	<ul style="list-style-type: none"> ❖ વાતાવરણમાં ભેજ વધવાને કારણે કીટક અને રોગનું પ્રમાણ વધે નહી તેના માટે યોગ્ય કાળજી લેવી. ચોમાસુ શાકભાજુ જોવાકે રીંગણા, ટમેટા, મરચા વગેરે માટે ધરુવાડી તૈયાર કરવા. ❖ અગામી પાચ દિવસમાં ઘણા સ્થળો પર મધ્યમ વરસાદ પડવાની સંભાવના હોય તો પિયત ટાળવુ તેમજ દવા જેવા રાસાયણીક દવાનો ઉપયોગ કરવો નહી.
પશુ	<ul style="list-style-type: none"> ❖ પશુઓને યોગ્ય આહાર આપવો. પશુઓને માખી અને મછરથી બચાવવા માટે રહેવાની જગ્યા સ્વચ્છ કરવી.

JUNAGADH

Phone: 0285-2670139

Date: 25-07-2014

(Period from 26 to 30 July -2014)

Significant weather for the preceding week

Rain fall (mm)	087.0
Maximum temperature °C	30.0-33.5
Minimum temperature °C	22.5-24.6
RH % (Morning/Afternoon)	94-83
Wind speed (Kmph)	9.6-13.0
Cloud cover	6-8

Weather forecast for Junagadh Dist. until 08:30 hrs of 30-07-2014

Parameters	Day-1 26-07-14	Day-2 27-07-14	Day-3 28-07-14	Day-4 29-07-14	Day-5 30-07-14
Rain fall (mm)	6	10	5	0	0
Maximum temperature °C	28	28	29	29	29
Minimum temperature °C	24	24	24	25	25
Cloud cover (Okta)	6	4	5	6	5
RH % (Morning/After noon)	78-67	80-71	80-72	79-74	81-77
Wind speed (Kmph)	027	031	035	039	036
Wind direction(Deg)	240	250	250	240	240

Agro meteorological Advisories

Crop	Stage	Agromet Advisories
Kharif Groundnut	Vegetative	<ul style="list-style-type: none"> During vaspa condition Interculturing and hand weeding practices should be carried out. Due to cloudy whether condition, there is chance for the incidence of collar rot of groundnut, for the control of seed treatment and soil application with <i>Trichoderama</i> at 6 kg/ha. For the control of yellowish of groundnut, spray 100 gm ferrous sulphate or 10 gm lemon flower in 10 liter of water at 8 to 10 days interval.
Cotton	Vegetative	<ul style="list-style-type: none"> During vaspa condition Interculturing and hand weeding practices should be carried out. Apply urea 25 kg and ammonium sulphate fertilizer 25 kg/ha to pre monsoon sown cotton crop.
Soybean	Vegetative	<ul style="list-style-type: none"> During vaspa condition Interculturing and hand weeding practices should be carried out.
Vegetable	Seedling/ Nursery	<ul style="list-style-type: none"> There may be chances of pest/ disease infestation in vegetables due to increase in relative humidity so take appropriate precautions against them. Prepared the nursery for kharif vegetables like brinjal, tomato, chilli etc. Due to moderate rainfall forecast during next five days at most places avoid irrigation and chemical application.
Livestock	-	<ul style="list-style-type: none"> Keep animal shed hygienic and clean to avoid flies and mosquitoes.

PORBANDAR

Phone : 0285-2670139

Date : 25-07-2014

(Period from 26 to 30 July -2014)

Weather forecast for Porbandar Dist. until 08:30 hrs of 30-07-2014

Parameters	Day-1 26-07-14	Day-2 27-07-14	Day-3 28-07-14	Day-4 29-07-14	Day-5 30-07-14
Rain fall (mm)	5	6	3	3	3
Maximum temperature °C	28	28	29	29	29
Minimum temperature °C	26	26	27	27	27
Cloud cover (Okta)	6	4	5	5	6
RH % (Morning/After noon)	78-66	80-70	80-71	79-75	81-77
Wind speed (Kmph)	025	027	032	036	033
Wind direction(Deg)	240	250	250	240	240

[Agro meteorological Advisories](#)

Crop	Stage	Agromet Advisories
Kharif Groundnut	Vegetative	<ul style="list-style-type: none"> During vaspa condition Interculturing and hand weeding practices should be carried out. Due to cloudy whether condition, there is chance for the incidence of collar rot of groundnut, for the control of seed treatment and soil application with <i>Tricoderama</i> at 6 kg/ha. For the control of yellowish of groundnut, spray 100 gm ferrous sulphate or 10 gm lemon flower in 10 liter of water at 8 to 10 days interval.
Cotton	Vegetative	<ul style="list-style-type: none"> During vaspa condition Interculturing and hand weeding practices should be carried out. Apply urea 25 kg and ammonium sulphate fertilizer 25 kg/ha to pre monsoon sown cotton crop.
Soybean	Vegetative	<ul style="list-style-type: none"> During vaspa condition Interculturing and hand weeding practices should be carried out.
Vegetable	Seedling/ Nursery	<ul style="list-style-type: none"> There may be chances of pest/ disease infestation in vegetables due to increase in relative humidity so take appropriate precautions against them. Prepared the nursery for kharif vegetables like brinjal, tomato, chilli etc. Due to moderate rainfall forecast during next five days at most places avoid irrigation and chemical application.
Livestock	-	<ul style="list-style-type: none"> Keep animal shed hygienic and clean to avoid flies and mosquitoes.