



ગ્રામીણ કૃષિ મૌસમ સેવા
કૃષિ હવામાન સેલ, કૃષિ વિજ્ઞાન વિભાગ
જુનાગઢ કૃષિ યુનિવર્સિટી
જુનાગઢ - ૩૬૨૦૦૧

ઈમેઇલ:- amfuind@yahoo.com ફોન નં. ૦૨૮૫-૨૬૭૦૧૩૮



હવામાન આગાહી આધારીત દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારનાં જુનાગઢ જીલ્લા માટે કૃષિ સલાહ બુલેટીન
(કૃષિ હવામાન સેલ, જુનાગઢ કૃષિ યુનિવર્સિટી અને ભારત હવામાન વિભાગ દ્વારા સંપત્ત)

(#આગાહી નો અમૃત મહોત્સવ#)

વાર:- શુક્રવાર

તારીખ:- ૨૬/૧૧/૨૦૨૧

હવામાન આગાહી આધારીત બુલેટીન: ૦૬ (૨૦૨૧-૨૨)

પાછલા પાંચ દિવસનું હવામાન (તારીખ:- ૨૧ થી ૨૫ નવેમ્બર, ૨૦૨૧)					હવામાન પરિબળો	હવામાન આગાહી આગામી પાંચ દિવસ માટે (તારીખ:- ૨૭ નવેમ્બર થી ૦૧ ડિસેમ્બર, ૨૦૨૧) માન્ય ૦૧ ડિસેમ્બર, ૨૦૨૧. સવારના ૮:૩૦ વાગ્યાં સુધી				
૨૧/૧૧	૨૨/૧૧	૨૩/૧૧	૨૪/૧૧	૨૫/૧૧	તારીખ	૨૭/૧૧	૨૮/૧૧	૨૯/૧૧	૩૦/૧૧	૦૧/૧૨
૦	૦	૦	૦	૦	વરસાદ (મિમી)	૦	૦	૦	૦	૦
૩૨.૬	૩૩.૬	૩૪.૫	૩૪.૫	૩૩.૨	મહત્વમાન તાપમાન (°સે.ગ્ર)	૩૫	૩૪	૩૪	૩૪	૩૩
૨૩.૬	૨૩.૦	૨૧.૪	૧૭.૮	૧૭.૦	લઘુત્વમાન તાપમાન (°સે.ગ્ર)	૧૪	૧૫	૧૫	૧૫	૧૪
ચોખ્યું	ચોખ્યું	ચોખ્યું	ચોખ્યું	ચોખ્યું	વાદળની સ્થિતિ	અંશત વાદળાચ્યું	મોટેભાગે વાદળાચ્યું	અંશત વાદળાચ્યું	મોટેભાગે વાદળાચ્યું	વાદળાચ્યું
૫૬	૭૭	૮૮	૬૬	૫૦	સવારના લેજના ટકા	૩૮	૪૧	૪૪	૩૮	૭૦
૫૩	૫૮	૪૯	૪૦	૩૧	બપોરબાદ ના લેજના ટકા	૧૮	૧૬	૨૧	૨૪	૨૫
૫.૬	૫.૧	૨.૩	૨.૫	૨.૭	પવનની ગતિ (કિમી/કલાક)	૮.૬	૮.૪	૧૦.૧	૧૪.૬	૧૬.૮
૫૦	૫૦	૫૦	૩૨૦	૫૦	પવનની દિશા	૨૭	૪૫	૬૮	૭૨	૫૮


સામાન્ય સલાહ

ભારત સરકારશ્રીના હવામાન વિભાગ તરફથી મળેલ હવામાન આગાહીની માહીતી અનુસાર અતેના ગ્રામીણ કૃષિ મૌસમ સેવા યોજના, જુનાગઢ કૃષિ યુનિવર્સિટી, જુનાગઢ તરફથી જણાવવામા આવે છે કે દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના જુનાગઢ જીલ્લામાં આગામી તારીખ ૨૭/૧૧/૨૦૨૧ થી ૦૧/૧૨/૨૦૨૧ દરમયાન આકાશ મુખ્યત્વે આગામી પાંચ દિવસોમાં અંશત વાદળાચ્યું થી વાદળાચ્યું રહેશે, જીલ્લામાં આગામી પાંચ દિવસ દરમયાન વરસાદની શક્યતા નથી. પવનની ઝડપ ૮.૪ થી ૧૬.૮ કિમી/કલાક રહેવાની શક્યતા રહેશે. આ સમયગાળામાં મહત્વમાન તાપમાન ૩૩ થી ૩૫ ડિગ્રી સેન્ટીગ્રેડ રહેશે તેમજ લઘુત્વમાન પદ્ધતિ ૧૪ થી ૧૫ ડિગ્રી સેન્ટીગ્રેડ રહેવાની સંભાવના છે.

- માન્ય વરસાદની સૂચી મુજબ જુનાગઢ જીવલામાં ૧૪ ઓક્ટોબર થી ૧૦ નવેમ્બર, ૨૦૨૧ ના સમયગાળા દરમિયાન વધુ હળવી સૂકી સ્થિતિ જોવા મળેલ છે.
- સાપ્તાહિક વરસાદ આગાહીની સૂચી મુજબ દક્ષિણ સૌરાષ્ટ્રનાં વિસ્તારમાં તા. ૦૧/૧૨/૨૦૨૧ થી તા. ૦૭/૧૨/૨૦૨૧ સામાન્ય થી ઉપર વરસાદની શક્યતાઓ છે.
- મેધ્દૂત એપ્લિકેશન:- કૃષિ હવામાન સલાહ સેવા અને હવામાન આગાહી અંગેની જાણકારી મેળવવા માટે ડાઉનલોડ કરો મેધ્દૂત એપ્લિકેશન કે જે ખેડૂતોને તેમની સ્થાનિક ભાષામાં સ્થાન, પાક અને પશુધન વિશે હવામાન આધારિત કૃષિ સલાહ પ્રદાન કરશે.
- મેધ્દૂત એપ્લિકેશનને ડાઉનલોડ કરવા માટે નીચેની લિંક પર ક્લિક કરો.
એન્ડ્રોઇડ(મોબાઇલ)ઉપયોગકર્તાનું:
<https://play.google.com/store/apps/details?id=com.aas.meghdoot>
આઈ.ઓ.એસ. ઉપયોગકર્તાનું:
<https://apps.apple.com/in/app/meghdoot/id1474048155>

- કોરોના (કોવીડ-૧૯) રોગચાળાની મહામારી ના લીધે, ખેડૂતમિત્રો એ ખેતર પર ખેતીની કામગીરી દરમિયાન ખેત સાધનો / ઓજારો અને ખેત મજૂરો સાથે એકબીજાથી સામાજિક અંતર રાખવું અને તે માટે ની સાવધાની અને સલામતી રાખવી તેમજ અગમયેતી જાળવવી.
- ખેડૂતોને ચોમાસું પાકોની કાપણી થયાં બાદ શિયાળુ પાકોનાં વાવેતર માટે ખેતરો તૈયાર કરવાની સલાહ આપવામાં આવે છે.
- શિયાળુ વાવેતર કરેલ પાકોમાં આંતરખેડ કરવી અને હાથથી નિંદામણની કામગીરી કરવી.
- શિયાળુ પાકોમાં ભલામણ મુજબ ખાતર અને કૃષિ રસાયણ આપવું.

NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 46
(12.11.2021 to 18.11.2021) over Agricultural region



Agricultural vigour is good over isolated pockets of Gujarat.

Source: <https://www.star.nesdis.noaa.gov>

પાકનું નામ	પાક અવસ્થા	રોગ / જીવાત / જાત	કૃષિ સલાહ
ધઉં	વાવેતર / ઉગાવો / વૃક્ષિ	ગુ.જુ.ધઉં - ૪૬૩, લોક - ૧ અને ૧૧, ગુ. ધઉં - ૩૬૬, ૪૮૬, ૧૭૩ અને ૧૧	<ul style="list-style-type: none"> ❖ ૨૬ નવેમ્બર થી ૧૦ ડિસેમ્બરનાં સમયગાળા દરમિયાન ધઉંનું મોડું વાવેતર કરવું. ❖ ઉધઈનો ઉપક્રમ ઘટાડવા અગાઉનાં પાકનાં અવશેષો પાંડડાં, મૂળ અને ડાળીઓનો નાશ કરવો. ❖ સારુ કહોવાચેલું છાણિયું ખાતર વાપરવું. ખાતર તરીકે દિવેલા, લીંબોળી અને કરંજ વગેરેના ખોળનો ઉપયોગ કરવો. ❖ ધઉંનાં ૧૦૦ કિ.ગ્રા. બિયારણ દીઠ બાયકેન્થીન ૧૦ ઈસી ૨૦૦ મિ.લિ. અથવા ફીપ્રોનીલ ૫ એસસી ૬૦૦ મિ.લિ. અથવા કલોરપાયરીફોસ ૨૦ ઈસી ૪૦૦ મિ.લિ. ૫ લિટર પાણીમાં મિશ્રણ કરી પટ આપવો. ❖ પાકને એકસરખાં સમયગાળામાં પીયત આપવું. ❖ ધઉંના ઉભા પાકમાં ઉધઈનો ઉપક્રમ શરૂ થતો જણાય તો તુરત જ એક હેક્ટર પાક વિસ્તાર માટે ફીપ્રોનીલ ૫ એસસી ૧.૬૦૦ લિટર અથવા કલોરપાયરીફોસ ૨૦ ઈસી ૧.૫ લિટર દવાને ૧૦૦ કિ.ગ્રા. રેતી સાથે મિશ્રણ કરી બરાબર લેળવીને દવાની માવજત આપેલ રેતી ધઉંના ઉભા પાકમાં પુંખવી અને ત્યાર બાદ પાકને હળવું પિયત આપવું. ❖ વાવેતર બાદ ૩૦ થી ૩૫ દિવસે નિદામણ નિયંત્રણ માટે ૨૦ મીલી ૨.૪ ડી. અથવા મેટસલ્ફ્યુરન મિથાઇલ ૦.૮ ગ્રામ ૧૦ લિટર પાણીમાં લેળવી છંટકાવ કરવો. ❖ પાકમાં હાથથી નિંદામણ ની કામગીરી કરવી.
ચણા	ઉગાવો / વૃક્ષિ	મોલો	<ul style="list-style-type: none"> ❖ પાકને એકસરખાં સમયગાળામાં પીયત આપવું. ❖ મોલોનાં પરિક્ષણ માટે પીળા ચીકણા પીંજર પ્રતિ હેક્ટેરે ૧૦ પ્રમાણે ગોઠવવા. ❖ પાકમાં હાથથી નિંદામણ અને આંતરખેડની કામગીરી કરવી.
ધાણા	ઉગાવો / વૃક્ષિ	મોલો	<ul style="list-style-type: none"> ❖ વધારે પડતું પાણી આપવું નહિ. ❖ મોલો સામે રક્ષણ મેળવવા થાયામેથોકઝામ ૭૦ ડબલ્યુએસ દવા ૪.૨ ગ્રામ/કિ.ગ્રા. બીજને પટ આપવો. ❖ પાકને એકસરખાં સમયગાળામાં પીયત આપવું. ❖ પાકમાં હાથથી નિંદામણ ની કામગીરી કરવી.

લસણ	ઉગાવો / વૃક્ષિ	ચુસીયા	<ul style="list-style-type: none"> ❖ પાકને એકસરખાં સમયગાળામાં પીયત આપવું. ❖ આ ચુસીયા જીવાતના અસરકારક નિયંત્રણ માટે બ્યુવેરીયા બાજીયાના ૧.૧૫ વે.પા. નો પ્રથમ છંટકાવ જીવાત દેખાય ત્યારે ૦.૦૦૩૫% (૩૦ ગ્રામ/૧૦ લીટર પાણીમાં)થી બીજો ૦.૦૦૭ ટકા (૬૦ ગ્રામ/૧૦ લીટર પાણી) અને ત્રીજો ૦.૦૦૯ ટકા (૮૦ ગ્રામ/૧૦ લીટર પાણી) ૧૦ દિવસના અંતરે કરવા. ૨. ❖ આ ચુસીયા જીવાતના રાસાયણિક નિયંત્રણ માટે ડાયમીથોએટ ૩૦ છ.સી. ૧૦ મિ.લી. અથવા કવીનાલફોસ ૨૫ છ.સી. ૨૦ મિ.લી. દવા ૧૦ લીટર પાણીમાં લેળવી દવાની અસરકારકતા વધારવા તેમાં સ્ટીકર ઉમેરી દ્રાવણનો છંટકાવ કરવો. ❖ પાકમાં હાથથી નિંદામણ ની કામગીરી કરવી.
ચોમાસું મગફળી	લણણી	-	<ul style="list-style-type: none"> ❖ મગફળીનાં પાકની લણણીની કામગીરી આગામી હવામાન પરિસ્થિતિ મુજબ કરવી.
કપાસ	ફૂલ / ઝીંડવા વિકાસ / વીણી	ગુલાબી ઈયળ	<ul style="list-style-type: none"> ❖ ફૂલ અવસ્થાથી ગુલાબી ઈયળની મોજણીની શરૂઆત કરવી. જો છોડ પર બંધ પાંખડીવાળા ફૂલ જોવા મળે તો એને વીણી લઇ તેનો નાશ કરવો. ❖ એક એકર વિસ્તાર માંથી ૨૦ લીલા જીડવા તોડી ગુલાબી ઈયળની ક્ષમ્યમાત્રાની ચકાસણી કરવી. ❖ ગુલાબી ઈયળની ક્ષમ્યમાત્રા ૧૦% નુકશાન કરેલ ફૂલ અથવા ૧૦% નુકશાન કરેલ જીડવા અથવા ૮ નર ફૂદા પકડાયેલાદ્રેપ/રાત્રી સતત ૩ દિવસ માટે છે. ❖ જો ગુલાબી ઈયળ ક્ષમ્યમાત્રા વટાવે તો થાયોડીકાર્બ ૭૫ ડબલ્યુ. પી. ૨૦ ગ્રામ અથવા કવીનાલફોસ ૨૦ એ. ફ. ૨૦ મિ.લી./૧૦ લીટર પાણીમાં લેળવી પાક પર છંટકાવ કરવો. ❖ બર્ડ પર્ચીસ ગોઠવવા. ❖ કલોરોપાયરીફોસ ૨૦% ઈ. સી. ૨૫ મી. લી. અથવા થાયોડીકાર્બ ૭૫% ડબલ્યુ. પી. ૨૦ ગ્રામ./૧૦ લીટર પાણીમાં લેળવી ભમરી છોડ્યાનાં ૧૦ દિવસ પછી છંટકાવ કરવો. ❖ સારો અને નુકશાનવાળો કપાસ જુદી વીણવો. સારા કપાસનો સંગ્રહ કરવો કે બજારમાં વેચવો જ્યારે નુકશાનવાળા કપાસનો નાશ કરવો.

			<ul style="list-style-type: none"> ❖ કપાસ જેમ જેમ પાકતો જાય તેમ તેમ વીણી કરવી અને વીણી સવારના સમયે કરવી. ❖ પાકને જરૂરીયાત મુજબ પીયત આપવુ.
તુવેર	ડાળી / ફૂલ	તુવેર નો વ્યંધત્વ વાયરસ , ચુસીયા	<ul style="list-style-type: none"> ❖ પ્રોપરગાઇટ પ્યાંસી અથવા સ્પાયરોમેસીફેન ૨૨.૬ એસસી ૧૦ મિલી/૧૦ લી. નો છંટકાવ કરી ને તુવેર નો વ્યંધત્વ વાયરસ અટકાવી શકાય છે. ❖ પાકમાં હાથથી નિંદામણ અને આંતરખેડની કામગીરી કરવી. ❖ પાકને જરૂરીયાત મુજબ પીયત આપવુ. ❖ તુવેરના પાકમાં ૫૦ % ફૂલ બેસવાની અવસ્થાએ ચુસીયા જીવાતના નિયત્રણ માટે ૧૫ મીલી પોલીટ્રીન અથવા ૨૦ ગ્રામ એસીફેટ અથવા ૫ મીલી ઈન્ડોક્ષાકાર્બ અથવા ૨૫ મીલી પ્રોફેનોફોસ દવાને ૧૦લીટર પાણીમાં લેળવી છંટકાવ કરવો.
એરંડા	ફૂલ / ગળ બંધાવા	અળ અને પાનના ટપકાનો રોગ	<ul style="list-style-type: none"> ❖ અળ અને પાનનાં ટપકાનાં રોગ માટે મેન્કોઝેબ ૭૫% વે.પા. ૨૬ ગ્રામ/૧૦ લી. અથવા કોપર ઓક્સિક્લોરાઇડ ૫૦% વે.પા. ૪૦ ગ્રામ/૧૦ લી. મુજબ છંટકાવ કરવો. ❖ પાકમાં હાથથી નિંદામણ અને આંતરખેડની કામગીરી કરવી. ❖ પાકને જરૂરીયાત મુજબ પીયત આપવુ. ❖ વાદળાયું હવામાન સ્થિતિને કારણે એરંડામાં ઘોડીયા ઈયળના નિયત્રણ માટે ૨૦ મીલી ક્વીનાલ્ફોસ (૧.૫ % ભુકી) અથવા ૪૦ ગ્રામ કાર્બોરીલ ૧૦ લી. પાણીમાં લેળવી છંટકાવ કરવો.
શાકભાજુ	વૃધ્ધી/ફૂલ/ ફળ	-	<ul style="list-style-type: none"> ❖ આ સમયગાળા દરમિયાન દવા છંટકાવની કામગીરી કરવી. ❖ પાકમાં હાથથી નિંદામણ અને આંતરખેડની કામગીરી કરવી. ❖ શાકભાજુ પાકોને સમયસર પીયત આપવુ. ❖ શાકભાજુના પાકોને નીદામણ મુકત કરવુ અને નિયમીત પીયત આપવુ. શાકભાજુ ને સમયસર ઉતારી ગ્રેડિંગ કરી બજારમાં મુકવા.
પશુપાલન	-	-	<ul style="list-style-type: none"> ❖ પશુઓને યોગ્ય આહાર આપવો & પીવા માટે ચોખ્યું પાણી આપવું અને પશુઓના રહેઠાણ ચોખ્યા રાખવા. ❖ માખી મચ્છરથી રક્ષણ માટે ફીનાઈલ છાટવુ. ❖ પશુઓને યોગ્ય આહારમાં ખનિજચુક્ત મિશ્રણ સાથે સંગ્રહિત ચારા આપો. ચારા ને સલામત સ્થળે સંગ્રહિત કરવો જોઈએ.

- ❖ ચારા માટે ચાફ કટર વાપરો.
- ❖ પશુઓમાં ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટાજ ર મીલી/ ૧ લીટર પાણીમાં નાખીને છાટવી.
- ❖ દુધાળા પશુઓના આઉને જીંક ઓક્સાઇડ અથલા બોરીક પાવડરથી બરાબર સાફ કરવા.



ગ્રામીણ કૃષિ મૌસમ સેવા
કૃષિ હવામાન સેલ, કૃષિ વિજ્ઞાન વિભાગ
જુનાગઢ કૃષિ યુનિવર્સિટી
જુનાગઢ - ૩૬૨૦૦૧

ઈમેઇલ:- amfujind@yahoo.com ફોન નં. ૦૨૮૫-૨૬૭૦૧૩૬



હવામાન આગાહી આધારીત દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારનાં પોરબંદર જુલ્લા માટે કૃષિ સલાહ બુલેટીન
(કૃષિ હવામાન સેલ, જુનાગઢ કૃષિ યુનિવર્સિટી અને ભારત હવામાન વિભાગ દ્વારા સંપદ્ધ)

(#આગાહી નો અમૃત મહોત્સવ#)

વાર:- શુક્રવાર

તારીખ:- ૨૫/૧૧/૨૦૨૧

હવામાન આગાહી આધારીત બુલેટીન: ૦૫૬ (૨૦૨૧-૨૨)

પાછલા પાંચ દિવસનું હવામાન (તારીખ:- ૨૧ થી ૨૫ નવેમ્બર, ૨૦૨૧)					હવામાન પરિબળો		હવામાન આગાહી આગામી પાંચ દિવસ માટે (તારીખ:- ૨૭ નવેમ્બર થી ૦૧ ડિસેમ્બર, ૨૦૨૧)				
૨૧/૧૧	૨૨/૧૧	૨૩/૧૧	૨૪/૧૧	૨૫/૧૧	તારીખ	૨૭/૧૧	૨૮/૧૧	૨૯/૧૧	૩૦/૧૧	૦૧/૧૨	
૦	૦	૦	૦	૦	વરસાદ (મિમી)	૦	૦	૦	૦	૦	
૩૨.૬	૩૪.૨	૩૪.૭	૩૪.૮	૩૩.૪	મહત્વમાન તાપમાન (°સ.ગ્ર.)	૩૪	૩૪	૩૫	૩૫	૩૪	
૨૩.૮	૨૩.૮	૨૧.૦	૧૭.૬	૧૭.૦	લઘુત્વમાન તાપમાન (°સ.ગ્ર.)	૧૮	૧૭	૧૮	૧૮	૧૭	
મોટેભાગે ચોખ્યું	મોટેભાગે વાદળાયું	મોટેભાગે ચોખ્યું	ચોખ્યું	ચોખ્યું	વાદળની સ્થિતિ	અંશત વાદળાયું	મોટેભાગે વાદળાયું	અંશત વાદળાયું	મોટેભાગે વાદળાયું	વાદળાયું	
૬૫	૬૨	૬૭	૬૧	૬૩	સવારના લેજના ટકા	૪૪	૪૮	૪૧	૪૬	૪૮	
૬૦	૪૮	૫૦	૪૩	૩૬	અપોરબાદ ના લેજના ટકા	૨૭	૨૬	૨૮	૩૧	૩૦	
૧૪.૮	૧૧.૧	-	૫.૬	૩.૭	પવનની ગતિ (કિમી/કલાક)	૧૨.૪	૧૩.૫	૧૨.૮	૧૩.૩	૨૦.૭	
૫૦	૫૦	-	૧૪૦	૩૬૦	પવનની દિશા	૨૨	૪૫	૭૦	૭૧	૪૫	

સામાન્ય સલાહ

ભારત સરકારશીના હવામાન વિભાગ તરફથી મળેલ હવામાન આગાહીની માહીતી અનુસાર અત્રેના ગ્રામીણ કૃષિ મૌસમ સેવા યોજના, જુનાગઢ કૃષિ યુનિવર્સિટી, જુનાગઢ તરફથી જણાવવામા આવે છે કે દક્ષિણ સૌરાષ્ટ્ર ખેત

આબોહવાકીય વિસ્તારના પોરબંદર જીલ્લામાં આગામી તારીખ ૨૭/૧૧/૨૦૨૧ થી ૦૧/૧૨/૨૦૨૧ દરમયાન આકાશ મુખ્યત્વે આગામી પાંચ દિવસોમાં અંશત વાદળણાયું થી વાદળણાયું રહેશે, જીલ્લામાં આગામી પાંચ દિવસો દરમિયાન વરસાદની શક્યતા નથી. પવનની ઝડપ ૧૨.૪ થી ૨૦.૭ કીમી/કલાક રહેવાની શક્યતા રહેશે. આ સમયગાળામાં મહત્વમાન તાપમાન ૩૪ થી ૩૫ ડિગ્રી સેન્ટીગ્રેડ રહેશે તેમજ લઘુત્તમ તાપમાન ૧૭ થી ૧૮ ડિગ્રી સેન્ટીગ્રેડ રહેવાની સંભાવના છે.

- + માન્ય વરસાદની સૂચી મુજબ પોરબંદર જીલ્લામાં ૧૪ ઓક્ટોબર થી ૧૦ નવેમ્બર, ૨૦૨૧ ના સમયગાળા દરમિયાન હળવી ભેજની સ્થિતિ જોવા મળેલ છે.
- + સાન્તાહિક વરસાદ આગાહીની સૂચી મુજબ દક્ષિણ સૌરાષ્ટ્રનાં વિસ્તારમાં તા. ૦૧/૧૨/૨૦૨૧ થી તા. ૦૭/૧૨/૨૦૨૧ સામાન્ય થી નીચે વરસાદની શક્યતાઓ છે.
- + મેધદૂત એપ્લીકેશન:- કૃષિ હવામાન સલાહ સેવા અને હવામાન આગાહી અંગેની જાણકારી મેળવવા માટે **ડાઉનલોડ** કરો મેધદૂત એપ્લીકેશન કે જે એડૂતોને તેમની સ્થાનિક ભાષામાં સ્થાન, પાક અને પશુધન વિશે હવામાન આધારિત કૃષિ સલાહ પ્રદાન કરશે.
- + મેધદૂત એપ્લીકેશનને **ડાઉનલોડ** કરવા માટે નીચેની લિંક પર ક્લિક કરો.
એન્ડ્રોઇડ(મોબાઇલ)ઉપયોગકર્તાઓ:


<https://play.google.com/store/apps/details?id=com.aas.meghdoot>

આઈ.ઓ.એ.સ. ઉપયોગકર્તાઓ:-

<https://apps.apple.com/in/app/meghdoot/id1474048155>

- ❖ કોરોના (કોવીડ-૧૯) રોગચાળાની મહામારી ના લીધે, એડૂતમિત્રો એ ખેતર પર ખેતીની કામગીરી દરમિયાન ખેત સાધનો / ઓજારો અને ખેત મજૂરો સાથે એકબીજાથી સામાજિક અંતર રાખવું અને તે માટે ની સાવધાની અને સલામતી રાખવી તેમજ અગમયેતી જાળવવી.
- ❖ એડૂતોને ચોમાસું પાકોની કાપણી થયાં બાદ શિયાળુ પાકોનાં વાવેતર માટે ખેતરો તૈયાર કરવાની સલાહ આપવામાં આવે છે.
- ❖ શિયાળુ વાવેતર કરેલ પાકોમાં આંતરએડ કરવી અને હાથથી નિંદામણની કામગીરી કરવી.
- ❖ શિયાળુ પાકોમાં ભલામણ મુજબ ખાતર અને કૃષિ રસાયણ આપવું.

NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 46
(12.11.2021 to 18.11.2021) over Agricultural region



Agricultural vigour is good over isolated pockets of Gujarat.

Source: <https://www.star.nesdis.noaa.gov>

પાકનું નામ	પાક અવસ્થા	રોગ / જીવાત / જાત	કૃષિ સલાહ
ઘઉં	વાવેતર / ઉગાવો / વૃદ્ધિ	ગુ.જુ.ઘઉં - ૪૬૩, લોક - ૧ અને ૧૧, ગુ. ઘઉં - ૩૬૬, ૪૮૬, ૧૭૩ અને ૧૧	<ul style="list-style-type: none"> ❖ ૨૬ નવેમ્બર થી ૧૦ ડિસેમ્બરનાં સમયગાળા દરમિયાન ઘઉંનું મોડું વાવેતર કરવું. ❖ ઉધર્ણો ઉપક્રમ ઘટાડવા અગાઉનાં પાકનાં અવશેષો પાંડાં, મૂળ અને ડાળીઓનો નાશ કરવો. ❖ સાચું કહેવાયેલું છાણિયું ખાતર વાપરવું. ખાતર તરીકે દિવેલા, લીંબોળી અને કરંજ વગેરેના ખોળનો ઉપયોગ કરવો. ❖ ઘઉંનાં ૧૦૦ કિ.ગ્રા. બિયારણ દીઠ બાયફેન્થીન ૧૦ ઈસી ૨૦૦ મિ.લિ. અથવા ફીપ્રોનીલ ૫ એસસી ૬૦૦ મિ.લિ. અથવા કલોરપાયરીફોસ ૨૦ ઈસી ૪૦૦ મિ.લિ. ૫ લિટર પાણીમાં મિશ્રણ કરી પટ આપવો. ❖ પાકને એકસરખાં સમયગાળામાં પીયત આપવું. ❖ ઘઉંના ઉભા પાકમાં ઉધર્ણો ઉપક્રમ શરૂ થતો જણાય તો તુરત ૪ એક હેક્ટાર પાક વિસ્તાર માટે ફીપ્રોનીલ ૫ એસસી ૧.૬૦૦ લિટર અથવા કલોરપાયરીફોસ ૨૦ ઈસી ૧.૫ લિટર દવાને ૧૦૦ કિ.ગ્રા. રેતી સાથે મિશ્રણ કરી બરાબર લેળવીને દવાની માવજત આપેલ રેતી ઘઉંના ઉભા પાકમાં પુંખવી અને ત્યાર બાદ પાકને હળવું પિયત આપવું.

			<ul style="list-style-type: none"> ❖ વાવેતર બાદ 30 થી 35 દિવસે નિંદામણ નિયંત્રણ માટે 20 મીલી ૨,૪ ડી. અથવા મેટસલ્ફ્યુરન મિથાઇલ ૦.૮ ગ્રામ ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો. ❖ પાકમાં હાથથી નિંદામણ ની કામગીરી કરવી.
ચણા	ઉગાવો / વૃદ્ધિ	મોલો	<ul style="list-style-type: none"> ❖ પાકને એકસરખાં સમયગાળામાં પીયત આપવું. ❖ મોલોનાં પરિક્ષણ માટે પીળા ચીકણા પીંજર પ્રતિ ફેક્ટેરે ૧૦ પ્રમાણે ગોઠવવા. ❖ પાકમાં હાથથી નિંદામણ અને આંતરખેડની કામગીરી કરવી.
જીરુ	વાવેતર / ઉગાવો	ચુસીયા	<ul style="list-style-type: none"> ❖ જીરુમાં થાયોમેથોકામ 70 ડબ્લ્યુએસ દવા ૪.૨ ગ્રામ અથવા ઇમીડાકલોપ્રીડ 70 ડબ્લ્યુએસ દવા ૧૦ ગ્રામ/કિ.ગ્રા. બીજને પટ આપીને વાવણી કરવાથી થ્રિપ્સ સામે રક્ષણ મળે છે. ❖ થાયરમ ૩ ગ્રામ/કિલો મુજબ બીજ માવજત આપવી. ❖ પાકને એકસરખાં સમયગાળામાં પીયત આપવું. ❖ ચુસીયાના નિયંત્રણ માટે એસીફેટ ૭૫ એસીપી ૧૦ ગ્રામ ૧૦ લીટર પાણીમાં ઉમેરી છંટકાવ કરવો. ❖ પાકમાં હાથથી નિંદામણ અને આંતરખેડની કામગીરી કરવી.
લસણ	ઉગાવો / વૃદ્ધિ	ચુસીયા	<ul style="list-style-type: none"> ❖ પાકને એકસરખાં સમયગાળામાં પીયત આપવું. ❖ આ ચુસીયા જીવાતના અસરકારક નિયંત્રણ માટે બ્યુવેરીયા બાઝીયાના ૧.૧૫ વે.પા. નો પ્રથમ છંટકાવ જીવાત દેખાય ત્યારે ૦.૦૦૩૫% (૩૦ ગ્રામ/૧૦ લીટર પાણીમાં)બ બીજો ૦.૦૦૭ ટકા (૬૦ ગ્રામ/૧૦ લીટર પાણી) અને ત્રીજો ૦.૦૦૯ ટકા (૮૦ ગ્રામ/૧૦ લીટર પાણી) ૧૦ દિવસના અંતરે કરવા. ૨. ❖ આ ચુસીયા જીવાતના રાસાયણિક નિયંત્રણ માટે ડાયમીથોએટ ૩૦ ઇ.સી. ૧૦ મી.લી. અથવા કવીનાલજોસ ૨૫ ઇ.સી. ૨૦ મી.લી. દવા ૧૦ લીટર પાણીમાં ભેળવી દવાની અસરકારકતા વધારવા તેમાં સ્ટીકર ઉમેરી દ્રાવણનો છંટકાવ કરવો. ❖ પાકમાં હાથથી નિંદામણ ની કામગીરી કરવી.

ચોમાસું મગફળી	લણણી		<ul style="list-style-type: none"> ❖ મગફળીનાં પાકની લણણીની કામગીરી આગામી હવામાન પરિસ્થિતિ મુજબ કરવી.
કપાસ	કૂલ / ઝીંડવા વિકાસ / વીણી	ગુલાબી ઈયળ	<ul style="list-style-type: none"> ❖ કૂલ અવસ્થાથી ગુલાબી ઈયળની મોજણીની શરૂઆત કરવી. જો છોડ પર બંધ પાંખડીવાળા કૂલ જોવા મળે તો એને વીણી લઇ તેનો નાશ કરવો. ❖ એક એકર વિસ્તાર માંથી ૨૦ લીલા ઝીંડવા તોડી ગુલાબી ઈયળની ક્ષમ્યમાત્રાની ચકાસણી કરવી. ❖ ગુલાબી ઈયળની ક્ષમ્યમાત્રા ૧૦% નુકશાન કરેલ કૂલ અથવા ૧૦% નુકશાન કરેલ ઝીંડવા અથવા ૮ નર ફૂદા પકડાયેલા/દ્રેપ/રાત્રી સતત ૩ દિવસ માટે છે. ❖ જો ગુલાબી ઈયળ ક્ષમ્યમાત્રા વટાવે તો થાયોડીકાર્બ ૭૫ ડબ્લ્યુ. પી. ૨૦ ગ્રામ અથવા કવીનાલફોસ ૨૦ એ. ફ. ૨૦ મી.લી./૧૦ લીટર પાણીમાં બેળવી પાક પર છંટકાવ કરવો. ❖ બર્ડ પર્ચિસ ગોઠવવા. ❖ પરજીવી ભમરી ટ્રાઇકોગામાં બેકટરી % ૫૦૫૦૦૦/ એકર છોડવી. ❖ કલોરોપાયરીફોસ ૨૦% ઇ. સી. ૨૫ મી. લી. અથવા થાયોડીકાર્બ ૭૫% ડબ્લ્યુ. પી. ૨૦ ગ્રામ./૧૦ લીટર પાણીમાં બેળવી ભમરી છોડયાનાં ૧૦ દિવસ પછી છંટકાવ કરવો. ❖ સારો અને નુકશાનવાળો કપાસ જુદો વીણાવો. સારા કપાસનો સંગ્રહ કરવો કે બજારમાં વેચવો જયારે નુકશાનવાળા કપાસનો નાશ કરવો. ❖ કપાસ જેમ જેમ પાકતો જાય તેમ તેમ વીણી કરવી અને વીણી સવારના સમયે કરવી. ❖ પાકને જરૂરીયાત મુજબ પીયત આપવુ.
તુવેર	ડાળી / કૂલ	તુવેર નો વ્યંઘત્વ વાયરસ, યુસીયા	<ul style="list-style-type: none"> ❖ પ્રોપરગાઈટ પ્યાંડિકેન ૨૨.૬ એસસી ૧૦ મિલી/૧૦ લી. નો છંટકાવ કરી ને તુવેર નો વ્યંઘત્વ વાયરસ અટકાવી શકાય છે. ❖ પાકમાં હાથથી નિંદામણ અને આંતરખેડની કામગીરી કરવી. ❖ પાકને જરૂરીયાત મુજબ પીયત આપવુ.

			<ul style="list-style-type: none"> ❖ તુવેરના પાકમાં ૫૦ % કુલ બેસવાની અવસ્થાએ ચુસીયા જીવાતના નિયત્રણ માટે ૧૫ મીલી પોલીટ્રીન અથવા ૨૦ ગ્રામ એસીફેટ અથવા ૫ મીલી ઈન્ડોક્ષાકાર્બ અથવા ૨૫ મીલી પ્રોફેનોફોસ દવાને ૧૦લીટર પાણીમાં લેળવી છંટકાવ કરવો.
એરંડા	કૂલ / ગળ બંધાવા	ઝર અને પાનના ટપકાનો રોગ	<ul style="list-style-type: none"> ❖ ઝર અને પાનનાં ટપકાનાં રોગ માટે મેન્જોઝેબ ૭૫% વે.પા. ૨૬ ગ્રામ/૧૦ લી. અથવા કોપર ઓક્સિક્લોરાઇડ ૫૦% વે.પા. ૪૦ ગ્રામ/૧૦ લી. મુજબ છંટકાવ કરવો. ❖ પાકમાં હાથથી નિંદામણ અને આંતરખેડની કામગીરી કરવી. ❖ પાકને જરૂરીયાત મુજબ પીયત આપવુ. ❖ વાદળછાયું હવામાન સ્થિતિને કારણે એરંડામાં ઘોડીયા ઈયળના નિયંત્રણ માટે ૨૦ મીલી ક્વીનાલ્ફોસ (૧.૫ % લુકી) અથવા ૪૦ ગ્રામ કાર્બોરીલ ૧૦ લી. પાણીમાં લેળવી છંટકાવ કરવો.
શાકભાજુ	વૃધ્ધી/કુલ/ફળ	-	<ul style="list-style-type: none"> ❖ આ સમયગાળા દરમિયાન દવા છંટકાવની કામગીરી કરવી. ❖ પાકમાં હાથથી નિંદામણ અને આંતરખેડની કામગીરી કરવી. ❖ શાકભાજુ પાકોને સમયસર પીયત આપવુ. ❖ શાકભાજુના પાકોને નીદામણ મુક્ત કરવુ અને નિયમીત પીયત આપવુ. શાકભાજુ ને સમયસર ઉતારી ગ્રેડિંગ કરી બજારમાં મુકવા.
પશુપાલન	-	-	<ul style="list-style-type: none"> ❖ પશુઓને યોગ્ય આહાર આપવો & પીવા માટે ચોખ્યું પાણી આપવું અને પશુઓના રહેઠાણ ચોખ્યા રાખવા. ❖ માખી મચ્છરથી રક્ષણ માટે ફીનાઈલ છાટવુ. ❖ પશુઓને યોગ્ય આહારમાં ખનિજયુક્ત મિશ્રણ સાથે સંગ્રહિત ચારા આપો. ચારા ને સલામત સ્થળે સંગ્રહિત કરવો જોઈએ. ❖ ચારા માટે ચાફ કટર વાપરો. ❖ પશુઓમાં ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેશ્ટીન અથવા એમીટાજ ૨ મીલી/ ૧ લીટર પાણીમાં નાખીને છાટવી. ❖ દુધાળા પશુઓના આઉને જીંક ઓક્સાઇડ અથલા બોરીક પાવડરથી બરાબર સાફ કરવા.



ગ્રામીણ કૃષિ મૌસમ સેવા
કૃષિ હવામાન સેલ, કૃષિ વિજ્ઞાન વિભાગ
જુનાગઢ કૃષિ યુનિવર્સિટી
જુનાગઢ - ૩૬૨૦૦૧

ઈમેઇલ:- amfujind@yahoo.com ફોન નં. ૦૨૮૫-૨૬૭૦૧૩૬



હવામાન આગાહી આધારીત દક્ષિણ સૌરાષ્ટ્ર પેટ આબોહવાકીય વિસ્તારનાં ગીર સોમનાથ જીલ્લા માટે કૃષિ સલાહ બુલેટીન

(કૃષિ હવામાન સેલ, જુનાગઢ કૃષિ યુનિવર્સિટી અને ભારત હવામાન વિભાગ દ્વારા સંપદ)

(#આગાહી નો અમૃત મહોત્સવ#)

વાર:- શુક્રવાર

તારીખ:- ૨૬/૧૧/૨૦૨૧

હવામાન આગાહી આધારીત બુલેટીન: ૦૬૮ (૨૦૨૧-૨૨)

પાછળા પાંચ દિવસનું હવામાન (તારીખ:- ૨૧ થી ૨૫ નવેમ્બર, ૨૦૨૧)					હવામાન પરિબળો	હવામાન આગાહી આગામી પાંચ દિવસ માટે (તારીખ:- ૨૭ નવેમ્બર થી ૦૧ ડિસેમ્બર, ૨૦૨૧)				
૨૧/૧૧	૨૨/૧૧	૨૩/૧૧	૨૪/૧૧	૨૫/૧૧	તારીખ	૨૭/૧૧	૨૮/૧૧	૨૯/૧૧	૩૦/૧૧	૦૧/૧૨
૦	૦	૦	૦	૦	વરસાદ (મિમી)	૦	૦	૦	૦	૦
૩૪.૦	૩૪.૦	૩૨.૨	૩૨.૭	૩૪.૬	મહત્વમાન (૦૦૦૦૦૦)	૩૪	૩૪	૩૪	૩૪	૩૨
૨૪.૬	૨૫.૨	૨૪.૭	૨૨.૬	૨૧.૧	લઘુત્વમાન (૦૦૦૦૦૦)	૧૮	૧૮	૧૮	૧૮	૧૮
ચોખ્યું	ચોખ્યું	ચોખ્યું	ચોખ્યું	ચોખ્યું	વાદળની સ્થિતિ	અંશત વાદળથાયું	મોટેભાગે વાદળથાયું	મોટેભાગે વાદળથાયું	મોટેભાગે વાદળથાયું	વાદળથાયું
૬૧	૬૩	૬૨	૬૦	૬૧	સવારના ભેજના ટકા	૫૩	૪૮	૫૩	૪૫	૬૩
૭૮	૭૩	૬૧	૬૬	૫૮	બપોરબાદ ના ભેજના ટકા	૨૫	૨૭	૨૮	૩૩	૩૩
૧.૬	૧.૬	૧.૬	૩.૭	૩.૭	પવનની ગતિ (કિમી/કલાક)	૮.૭	૮.૨	૧૧.૬	૧૭.૫	૧૮.૧
૮૮	૮૮	૮૮	૮૮	૮૮	પવનની દિશા (ડિગ્રી)	૧૫૮	૨૨	૪૫	૭૬	૪૮

સામાન્ય સલાહ

ભારત સરકારશ્રીના હવામાન વિભાગ તરફથી મળેલ હવામાન આગાહીની માહીતી અનુસાર અત્રેના ગ્રામીણ કૃષિ મૌસમ સેવા યોજના, જુનાગઢ કૃષિ યુનિવર્સિટી, જુનાગઢ તરફથી જણાવવામા આવે છે કે દક્ષિણ સૌરાષ્ટ્ર પેટ આબોહવાકીય વિસ્તારનાં ગીર સોમનાથ જીલ્લામાં આગામી તારીખ ૨૭/૧૧/૨૦૨૧ થી ૦૧/૧૨/૨૦૨૧ દરમયાન આકાશ મુખ્યત્વે આગામી પાંચ દિવસોમાં અંશત વાદળથાયું થી વાદળથાયું રહેશે, જીલ્લામાં આગામી પાંચ દિવસો દરમયાન વરસાદની શક્યતાઓ નથી. પવનની ઝડપ ૮.૭ થી ૧૮.૧ કિમી/કલાક રહેવાની શક્યતા રહેશે. આ સમયગાળામાં મહત્વમાન તાપમાન ૩૨ થી ૩૫ ડિગ્રી સેન્ટીગ્રેડ રહેશે તેમજ લઘુત્વમાન ઠી થી ૧૮ ડિગ્રી સેન્ટીગ્રેડ રહેવાની સંભાવના છે.


- માન્ય વરસાદની સૂચી મુજબ ગીર સોમનાથ જીવલામાં ૧૪ ઓક્ટોબર થી ૧૦ નવેમ્બર, ૨૦૨૧ ના સમયગાળા દરમિયાન હળવી ભેજની સ્થિતિ જોવા મળેલ છે.
- સાપ્તાહિક વરસાદ આગાહીની સૂચી મુજબ દક્ષિણ સૌરાષ્ટ્રમાં વિસ્તારમાં તા. ૦૧/૧૨/૨૦૨૧ થી તા. ૦૭/૧૨/૨૦૨૧ સામાન્ય થી ઉપર વરસાદની શક્યતાઓ છે.

- મેધ્ડૂત એપ્લીકેશન:** કૃષિ હવામાન સલાહ સેવા અને હવામાન આગાહી અંગેની જાણકારી મેળવવા માટે ડાઉનલોડ કરો મેધ્ડૂત એપ્લીકેશન કે જે ખેડૂતોને તેમની સ્થાનિક ભાષામાં સ્થાન, પાક અને પશુધન વિશે હવામાન આધારિત કૃષિ સલાહ પ્રદાન કરશે.
- મેધ્ડૂત એપ્લીકેશનને ડાઉનલોડ કરવા માટે નીચેની લિંક પર ક્લિક કરો.**
એન્ડ્રોઇડ(મોબાઇલ)ઉપયોગકર્તાઃ
<https://play.google.com/store/apps/details?id=com.aas.meghdoot>

આઈ.ઓ.એસ. ઉપયોગકર્તાઃ-

<https://apps.apple.com/in/app/meghdoot/id1474048155>

- કોરોના (કોવિડ-૧૯) રોગચાળાની મહામારી ના લીધે, ખેડૂતમિત્રો એ ખેતર પર ખેતીની કામગીરી દરમિયાન ખેત સાધનો / ઓજારો અને ખેત મજુરો સાથે એકબીજાથી સામાજિક અંતર રાખવું અને તે માટે ની સાવધાની અને સલામતી રાખવી તેમજ અગમચેતી જળવવી.
- ખેડૂતોને ચોમાસું પાકોની કાપણી થયાં બાદ શિયાળું પાકોનાં વાવેતર માટે ખેતરો તૈયાર કરવાની સલાહ આપવામાં આવે છે.
- શિયાળું વાવેતર કરેલ પાકોમાં આંતરખેડ કરવી અને હાથથી નિંદામણની કામગીરી કરવી.
- શિયાળું પાકોમાં ભલામણ મુજબ ખાતર અને કૃષિ રસાયણ આપવું.



પાકનું નામ	પાક અવસ્થા	રોગ / જીવાત / જાત	કૃષિ સલાહ
ઘઉં	વાવેતર / ઉગાવો / વૃક્ષિ	મોડુ વાવેતર :- ગુ.જ.ઘઉં - ૧૭૩, લોક - ૧, ગુ. ઘઉં - ૧૧ અને એમ પી - ૩૨૮૮	<ul style="list-style-type: none"> ❖ ૨૬ નવેમ્બર થી ૧૦ ડિસેમ્બરનાં સમયગાળા દરમિયાન ઘઉંનું મોડુ વાવેતર કરવું. ❖ ઉધર્ણો ઉપદ્રવ ઘટાડવા અગાઉનાં પાકનાં અવશેષો પાંડાં, મૂળ અને ડાળીઓનો નાશ કરવો. ❖ સારુ કહોવાયેલું છાણિયું ખાતર વાપરવું. ખાતર તરીકે દિવેલા, લીંબોળી અને કરંજ વગેરેના ખોળનો ઉપયોગ કરવો. ❖ ઘઉંનાં ૧૦૦ કિ.ગ્રા. બિયારણ દીઠ બાયફેન્થીન ૧૦ ઈસી ૨૦૦ મિ.લિ. અથવા ફીપ્રોનીલ ૫ એસસી ૬૦૦ મિ.લિ. અથવા કલોરપાયરીફોસ ૨૦ ઈસી ૪૦૦ મિ.લિ. ૫ લિટર પાણીમાં મિશ્રણ કરી પટ આપવો. ❖ પાકને એકસરખાં સમયગાળામાં પીયત આપવું. ❖ ઘઉંના ઉભા પાકમાં ઉધર્ણો ઉપદ્રવ શરૂ થતો જણાય તો તુરત ૪ એક હેક્ટર પાક વિસ્તાર માટે ફીપ્રોનીલ ૫ એસસી ૧.૬૦૦ લિટર અથવા કલોરપાયરીફોસ ૨૦ ઈસી ૧.૫ લિટર દવાને ૧૦૦ કિ.ગ્રા. રેતી સાથે મિશ્રણ કરી બરાબર લેળવીને દવાની માવજત આપેલ રેતી ઘઉંના ઉભા પાકમાં પુંખવી અને ત્યાર બાદ પાકને હળવું પિયત આપવું. ❖ વાવેતર બાદ ૩૦ થી ૩૫ દિવસે નિદામણ નિયંત્રણ માટે ૨૦ મીલી ૨.૪ ડી. અથવા મેટસલ્ફ્યુરન મિથાઈલ ૦.૮ ગ્રામ ૧૦ લિટર પાણીમાં લેળવી છંટકાવ કરવો. ❖ પાકમાં હાથથી નિંદામણ ની કામગીરી કરવી.
ચણા	ઉગાવો / વૃક્ષિ	મોલો	<ul style="list-style-type: none"> ❖ પાકને એકસરખાં સમયગાળામાં પીયત આપવું. ❖ મોલોનાં પરિક્ષણ માટે પીળા ચીકણા પીંજર પ્રતિ હેક્ટેરે ૧૦ પ્રમાણે ગોઠવવા. ❖ પાકમાં હાથથી નિંદામણ અને આંતરખેડની કામગીરી કરવી.
ઘાણા	ઉગાવો / વૃક્ષિ	મોલો	<ul style="list-style-type: none"> ❖ વધારે પડતું પાણી આપવું નહિં. ❖ મોલો સામે રક્ષણ મેળવવા થાયામેથોકામ ૭૦ ડબલ્યુએસ દવા ૪.૨ ગ્રામ/કિ.ગ્રા. બીજને પટ આપવો.

			<ul style="list-style-type: none"> ❖ પાકને એકસરખાં સમયગાળામાં પીયત આપવુ. ❖ પાકમાં હાથથી નિંદામણ ની કામગીરી કરવી.
લસણ	ઉગાવો / વૃક્ષિ	ચુસીયા	<ul style="list-style-type: none"> ❖ પાકને એકસરખાં સમયગાળામાં પીયત આપવુ. ❖ આ ચુસીયા જીવાતના અસરકારક નિયંત્રણ માટે બ્યુવેરીયા બાજીયાના ૧.૧૫ વે.પા. નો પ્રથમ છંટકાવ જીવાત દેખાય ત્યારે ૦.૦૦૩૫% (૩૦ ગ્રામ/૧૦ લીટર પાણીમાં)બ બીજો ૦.૦૦૭ ટકા (૬૦ ગ્રામ/૧૦ લીટર પાણી) અને ત્રીજો ૦.૦૦૯ ટકા (૮૦ ગ્રામ/૧૦ લીટર પાણી) ૧૦ દિવસના અંતરે કરવા. 2. ❖ આ ચુસીયા જીવાતના રાસાયણિક નિયંત્રણ માટે ડાયમીથોએટ ૩૦ છ.સી. ૧૦ મિ.લી. અથવા કવીનાલફોસ ૨૫ છ.સી. ૨૦ મિ.લી. દવા ૧૦ લીટર પાણીમાં લેળવી દવાની અસરકારકતા વધારવા તેમાં સ્ટીકર ઉમેરી દ્રાવણનો છંટકાવ કરવો.
તુવેર	ડાળી / ફૂલ	તુવેર નો વ્યંધત્વ વાયરસ , ચુસીયા	<ul style="list-style-type: none"> ❖ પ્રોપરગાઇટ ૫૭ ઈસી અથવા સ્પાયરોમેસીફેન ૨૨.૬ એસસી ૧૦ મિલી/૧૦ લી. નો છંટકાવ કરી ને તુવેર નો વ્યંધત્વ વાયરસ અટકાવી શકાય છે. ❖ પાકમાં હાથથી નિંદામણ અને આંતરખેડની કામગીરી કરવી. ❖ પાકને જરૂરીયાત મુજબ પીયત આપવુ. ❖ તુવેરના પાકમાં ૫૦ % કુલ બેસવાની અવસ્થાએ ચુસીયા જીવાતના નિયત્રણ માટે ૧૫ મીલી પોલીટ્રીન અથવા ૨૦ ગ્રામ એસીફેટ અથવા ૫ મીલી ઈન્ડોક્ષાકાર્બ અથવા ૨૫ મીલી પ્રોફેનોફોસ દવાને ૧૦લીટર પાણીમાં લેળવી છંટકાવ કરવો.
શાકભાજુ	વૃધ્ઘણી/કુલ/ ફળ	-	<ul style="list-style-type: none"> ❖ આ સમયગાળા દરમિયાન દવા છંટકાવની કામગીરી કરવી. ❖ પાકમાં હાથથી નિંદામણ અને આંતરખેડની કામગીરી કરવી. ❖ શાકભાજુ પાકોને સમયસર પીયત આપવુ. ❖ શાકભાજુના પાકોને નીદામણ મુકત કરવુ અને નિયમીત પીયત આપવુ. શાકભાજુ ને સમયસર ઉતારી ગ્રેડિંગ કરી બજારમાં મુકવા.

પશુપાલન	-	-	<ul style="list-style-type: none"> ❖ પશુઓને યોગ્ય આહાર આપવો & પીવા માટે ચોખ્યું પાણી આપવું અને પશુઓના રહેઠાણ ચોખ્મા રાખવા. ❖ માખી મચ્છરથી રક્ષણ માટે ફીનાઈલ છાટવું. ❖ પશુઓને યોગ્ય આહારમાં ખનિજયુક્ત મિશ્રણ સાથે સંગ્રહિત ચારા આપો. ચારા ને સલામત સ્થળે સંગ્રહિત કરવો જોઈએ. ❖ ચારા માટે ચાફ કટર વાપરો. ❖ પશુઓમાં ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટાજ ૨ મીલી/ ૧ લીટર પાણીમાં નાખીને છાટવી. ❖ દુધાળા પશુઓના આઉને ઝીંક ઓક્સાઇડ અથલા બોરીક પાવડરથી બરાબર સાફ કરવા.
---------	---	---	--

યાંત્રિક અધિકારી
ગ્રામીણ કૃષિ મૌસમ સેવા
જી.કૃ.યુ.

	Gramin Krushi Mausam Seva (GKMS) Agro meteorological Cell Department of Agronomy Junagadh Agricultural University Junagadh-362001 E-mail: amfujnd@yahoo.com Phone: 0285-2670139	
--	--	--

Agromet Advisory Service Bulletin for Junagadh District of South Saurashtra Agro-Climatic Zone

(Issued jointly by Junagadh Agricultural University and India Meteorological Department)

(#AZADI KA AMRIT MAHOTSAV#)

Day:- Friday

Date: 26-11-2021

Agromet Advisory Service bulletin. No:- 069 (2021-22)

Observed Weather during last Five days (Dated 21 st to 25 th November, 2021)					Weather Parameters	Weather Forecast upcoming Five days (Valid for 27 th Nov. to 01 st Dec., 2021) until 08:30 hrs of 01-12-2021				
21/11	22/11	23/11	24/11	25/11	Date	27/11	28/11	29/11	30/11	01/12
0	0	0	0	0	Rainfall(mm)	0	0	0	0	0
32.6	33.6	34.5	34.5	33.2	T _{max} (°C)	35	34	34	34	33
23.6	23.0	21.4	17.8	17.0	T _{min} (°C)	14	15	15	15	14
7	0	0	0	0	Cloud Cover	4	6	4	7	8
69	77	89	66	50	RH-I (%)	38	41	44	39	70
53	58	49	40	31	RH-II (%)	18	19	21	24	25
5.9	5.1	2.3	2.5	2.7	Wind speed (km/hr)	8.6	8.4	10.1	14.9	16.8
50	50	50	320	50	Wind direction	27	45	68	72	58

Agro meteorological Advisories

General Advice	Rain fall	As per IMD Forecast, There is forecasting of no rainfall occurs during next five days. Sky will be remain partly cloudy to cloudy.
	Temperature	Maximum and minimum temperature likely to be in range of 33 - 35°C and 14 - 15°C, respectively.
	RH (%)	The Relative Humidity is likely to be in the range of 18 – 70 %.
	Wind Direction and Wind Speed	Direction and speed of wind is NE & E and 8.4 – 16.8 km/h, respectively.

❖ **Standardized Precipitation Index:** - For the period of **14th October to 10th November, 2021** for Junagadh district considered under the mildly dry condition.

❖ **AS per ERFS products Above Normal rainfall may occur in South Saurashtra region during 01st to 07th December, 2021.**

❖ **MEGHDOOT Mobile Application:**- **Download** the MEGHDOOT mobile application for Agromet Advisory Services and information on weather forecasting, which will provide weather based agricultural advice to farmers for their location, crops and livestock in their local language.


Click on the following link to download the MEGHDOOT application:

Android users: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>

IOS users: <https://apps.apple.com/in/app/meghdoot/id1474048155>

- ❖ **Due to pandemic of CORONA (COVID-19), Farmers should maintain social distancing and safety precaution while handling farm machines and labour in the field.**
- ❖ **Farmers are advised to prepare their field after harvesting of kharif crops for the sowing of rabi season crops timely.**
- ❖ **Interculturing and hand weeding should be carried out in rabi sowing crops.**
- ❖ **To apply fertilizer and agrochemicals as per recommendation dose in rabi crops.**

NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 46
(12.11.2021 to 18.11.2021) over Agricultural region



Agricultural vigour is good over isolated pockets of Gujarat.

Source: <https://www.star.nesdis.noaa.gov>

Crop	Variety/ Pest/Insect	Stage	Agro met Advisories
Wheat	Late Sowing:- GJW-173, Lok-1, GW-11 and MP - 3288	Sowing / Germination / Vegetative	<ul style="list-style-type: none"> ➤ Late sowing of wheat crop done during 26th Nov. to 10th December duration. ➤ To control Termite, Removal and destruction of preceding crop plant parts. ➤ Seed treatment with bifenthrin 10 EC 200 ml or fipronil 5 SC 600 ml or chlorpyriphos 20 EC 400 ml in 5 litre of water for 100 kg of wheat seed. ➤ Apply irrigation to the crop at regular interval. ➤ Apply chlorpyriphos 20 EC 1.5 litre or fipronil 5 SC 1.6 litre in 100 kg of sand, mix it and spread it and give the light irrigation or through irrigation (chemigation). ➤ At 30-35 days after sowing spray 20 ml 2-4-D ethyle easter or Metasulfuron 0.8 gm/10 liters of water to control broad leafy weeds in wheat crop.
Chickpea	Aphid	Germination / Vegetative	<ul style="list-style-type: none"> ➤ Apply irrigation to the crop at regular interval.

			<ul style="list-style-type: none"> ➤ Set up yellow stick trap @ 10/ha to monitor aphid population. ➤ Hand weeding and Interculturing operation should be carry out.
Coriander	Aphid	Germination / Vegetative	<ul style="list-style-type: none"> ➤ Over irrigation should be avoided. ➤ Seed treatment of thiamethoxam 70 WS 4.2 gram/kg against aphid. ➤ Apply irrigation to the crop at regular interval. ➤ Hand weeding operation should be carry out.
Garlic	Thrips	Germination / Vegetative	<ul style="list-style-type: none"> ➤ Apply irrigation to the crop at regular interval. ➤ For management of thrips, apply schedule spraying of Beauveria bassiana 1.15 WP (min 2 x 106 cfu/g), first spray at initiation of pest infestation 0.0035% (30 g/10 l of water). Subsequent second 0.007% (60 g/10 l of water) and third 0.009% (80 g/10 l of water) spray at 10 days interval. ➤ To chemical control of thrips, spray Dimethoate 30 EC (10 ml/10 l. of water) or Quinalphos 25 EC (20 ml/10 l. of water). ➤ Hand weeding operation should be carry out.
Kharif Groundnut	-	Threshing	<ul style="list-style-type: none"> ➤ To done the operation of threshing in groundnut as per upcoming weather condition.
Cotton	Pink bollworm	Flowering / Boll formation, development / Picking	<ul style="list-style-type: none"> ➤ Assess ETL by plucking 20 green bolls from randomly selected plants across one acre. ➤ ETL of pink bollworm -10% infested flowers or 10% infested bolls or 8 male moths catches/trap/night for 3 consecutive nights ➤ If ETL crossed, spray Thiodicarb 75 WP 20g or Quinalphos 20AF 20ml per 10 liters of water.

			<ul style="list-style-type: none"> ➤ Set up bird perches. ➤ Release parasitoid Trichogramma bactrae @60000/acre if available ➤ Spray Chlorpyrifos 20%EC 25ml Or Thiodicarb 75 WP 20g per 10 litres of water after 10 days of parasitoid release. ➤ Picking of clean and infested cotton may be carried out separately. Clean cotton may be stored or marketed. Infested cotton should be destroyed. ➤ Picking of cotton should be carried out during morning hours. ➤ Apply irrigation to the crop at critical stage.
Pigeon pea	Sterility mosaic virus, Sucking pest	Branching / Flowering	<ul style="list-style-type: none"> ➤ To Spray Propergite 57 EC or Spiromesifen 22.9 SC 10 ml/10 l control of the sterility mosaic virus. ➤ Hand weeding and interculturing operation should be carry out. ➤ Apply irrigation to the crop at critical stage. ➤ For the control of sucking pest at 50 % flowering, spray 15 ml politrin OR 20 gm acephate OR 50 ml endoxacarb OR 25 ml profenophos in 10 liter of water.
Castor	Blight and Leaf spot	Flowering / Capsule formation	<ul style="list-style-type: none"> ➤ To control of blight and leaf spot, spray with Mancozeb 75 % WP 26 g/10 lit or copper oxichloride 50 % WP 40g/10 lit. ➤ Hand weeding and interculturing operation should be carry out. ➤ Apply irrigation to the crop at critical stage. ➤ Due to cloudy weather condition for the control of castor semi looper, spray 20 ml

			quinalphos (1.5 % powder) OR 40 gm carbaryl in 10 liter of water.
Vegetable crops	-	Sowing/ Flowering/ Fruiting	<ul style="list-style-type: none"> ➤ To the spray of agrochemicals during this period. ➤ Hand weeding and interculturing operation should be carry out. ➤ Apply irrigation to the crop at regular interval. ➤ Keep vegetable crop weed free, apply irrigation regularly and send vegetable fruit to the market often grading.
Livestock	-	-	<ul style="list-style-type: none"> ➤ To provide adequate food & clean water for drinking and keep the animals clean shed. ➤ Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Give stored fodder with mineral mixture. Fodder should be stored at safe place. ➤ Feed silage & hay material along with concentrate feed. ➤ Use chaff cutter for fodder. ➤ For the control of tick and other parasites, spray deltemethrin or aemitzaz 2 ml in 1 litre of water. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.



Gramin Krushi Mausam Seva (GKMS)
Agro meteorological Cell
Department of Agronomy
Junagadh Agricultural University
Junagadh-362001

E-mail: amfujnd@yahoo.com

Phone: 0285-2670139



Agromet Advisory Service Bulletin for Porbandar District of South Saurashtra Agro-Climatic Zone
(Issued jointly by Junagadh Agricultural University and India Meteorological Department)

(#AZADI KA AMRIT MAHOTSAV #)

Day:- Friday

Date: 26-11-2021

Agromet Advisory Service bulletin. No:- 069 (2021-22)


Observed Weather during last Five days (Dated 21 st to 25 th November, 2021)					Weather Parameters	Weather Forecast upcoming Five days (Valid for 27 th Nov. to 01 st Dec., 2021) until 08:30 hrs of 01-12-2021				
21/11	22/11	23/11	24/11	25/11	Date	27/11	28/11	29/11	30/11	01/12
0	0	0	0	0	Rainfall(mm)	0	0	0	0	0
32.9	34.2	34.7	34.8	33.5	Tmax (°C)	34	34	35	35	34
23.8	23.8	21.0	17.9	17.0	Tmin (°C)	18	17	18	18	17
2	3	2	0	0	Cloud Cover	3	7	4	7	8
65	62	67	81	93	RH-I (%)	44	48	41	49	48
60	48	50	43	36	RH-II (%)	27	29	29	31	30
14.8	11.1	-	5.6	3.7	Wind speed (km/hr)	12.4	13.5	12.8	13.3	20.7
50	50	-	140	360	Wind direction	22	45	70	71	45

Agro meteorological Advisories

General Advice	Rain fall	As per IMD Forecast, There is forecasting of no rainfall occurs during next five days. Sky will be remain partly cloudy to cloudy.
	Temperature	Maximum and minimum temperature likely to be in range of 34 - 35°C and 17 - 18°C, respectively.
	RH (%)	The Relative Humidity is likely to be in the range of 27 – 49 %.
	Wind Direction and Wind Speed	Direction and speed of wind is to be N, NE & E and 12.4 – 20.7 km/h, respectively.

- ❖ **Standardized Precipitation Index:** - For the period **14th October to 10th November, 2021** for Porbandar district considered under the mildly wet condition.
- ❖ **AS per ERFS products** **Above Normal rainfall may occur in South Saurashtra region during 01st to 07th December, 2021.**
- ❖ **MEGHDOOT Mobile Application:** - **Download** the MEGHDOOT mobile application for Agromet Advisory Services and information on weather forecasting, which will provide weather based agricultural advice to farmers for their location, crops and livestock in their local language.
Click on the following link to download the MEGHDOOT application:
Android users: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>
IOS users: <https://apps.apple.com/in/app/meghdoot/id1474048155>
- ❖ **Due to pandemic of CORONA (COVID-19), Farmers should maintain social distancing and safety precaution while handling farm machines and labour in the field.**
- ❖ **Farmers are advised to prepare their field after harvesting of kharif crops for the sowing of rabi season crops timely.**
- ❖ **Interculturing and hand weeding should be carried out in rabi sowing crops.**
- ❖ **To apply fertilizer and agrochemicals as per recommendation dose in rabi crops.**

**NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 46
(12.11.2021 to 18.11.2021) over Agricultural region**



Agricultural vigour is good over isolated pockets of Gujarat.

Source: <https://www.star.nesdis.noaa.gov>

Crop	Variety/ Pest / Insect	Stage	Agro met Advisories
Wheat	Late Sowing:- GJW-173, Lok-1, GW-	Sowing / Germination / Vegetative	<ul style="list-style-type: none"> ➤ Late sowing of wheat crop done during 26th Nov. to 10th December duration. ➤ To control Termite, Removal and destruction of preceding crop plant parts.

	11 and MP - 3288		<ul style="list-style-type: none"> ➤ Apply well decomposed FYM to soil. Use castor, neem or karanj cake as organic manure. ➤ Seed treatment with bifenthrin 10 EC 200 ml or fipronil 5 SC 600 ml or chlorpyriphos 20 EC 400 ml in 5 litre of water for 100 kg of wheat seed. ➤ Apply irrigation to the crop at regular interval. ➤ Apply chlorpyriphos 20 EC 1.5 litre or fipronil 5 SC 1.6 litre in 100 kg of sand, mix it and spread it and give the light irrigation or through irrigation (chemigation). ➤ At 30-35 days after sowing spray 20 ml 2-4-D ethyle easter or Metasulfuron 0.8 gm/10 liters of water to control broad leafy weeds in wheat crop.
Chickpea	Aphid	Germination / Vegetative	<ul style="list-style-type: none"> ➤ Apply irrigation to the crop at regular interval. ➤ Set up yellow stick trap @ 10/ha to monitor aphid population. ➤ Hand weeding and Interculturing operation should be carry out.
Cumin	Guj. Cumin – 3 and 4 / Thrips	Germination / Establishment	<ul style="list-style-type: none"> ➤ Seed treatment of thiamethoxam 70 WS 4.2 gram or imidacloprid 70 WS 10 gram/kg against thrips. ➤ Seed treatment of thiram 3 g/kg seed. ➤ Apply irrigation to the crop at regular interval.

			<ul style="list-style-type: none"> ➤ Spray acephate 75 SP 10 gram per 10 liter of water to control thrips. ➤ Hand weeding and interculturing operation should be carry out.
Garlic	Thrips	Germination / Vegetative	<ul style="list-style-type: none"> ➤ Apply irrigation to the crop at regular interval. ➤ For management of thrips, apply schedule spraying of Beauveria bassiana 1.15 WP (min 2×10^6 cfu/g), first spray at initiation of pest infestation 0.0035% (30 g/10 l of water). Subsequent second 0.007% (60 g/10 l of water) and third 0.009% (80 g/10 l of water) spray at 10 days interval. ➤ To chemical control of thrips, spray Dimethoate 30 EC (10 ml/10 l. of water) or Quinalphos 25 EC (20 ml/10 l. of water). ➤ Hand weeding operation should be carry out.
Kharif Groundnut	-	Threshing	<ul style="list-style-type: none"> ➤ To done the operation of threshing in groundnut as per upcoming weather condition.
Cotton	Pink bollworm	Flowering / Boll formation, development / Picking	<ul style="list-style-type: none"> ➤ Assess ETL by plucking 20 green bolls from randomly selected plants across one acre. ➤ ETL of pink bollworm -10% infested flowers or 10% infested bolls or 8 male moths catches/trap/night for 3 consecutive nights ➤ If ETL crossed, spray Thiodicarb 75 WP 20g or Quinalphos 20AF 20ml per 10 liters of water. ➤ Set up bird perches. ➤ Release parasitoid Trichogramma bactrae @60000/acre if available

			<ul style="list-style-type: none"> ➤ Spray Chlorpyrifos 20%EC 25ml Or Thiodicarb 75 WP 20g per 10 litres of water after 10 days of parasitoid release. ➤ Picking of clean and infested cotton may be carried out separately. Clean cotton may be stored or marketed. Infested cotton should be destroyed. ➤ Picking of cotton should be carried out during morning hours. ➤ Apply irrigation to the crop at critical stage.
Pigeon pea	Sterility mosaic virus	Branching / Flowering	<ul style="list-style-type: none"> ➤ To Spray Propergite 57 EC or Spiromesifen 22.9 SC 10 ml/10 l control of the sterility mosaic virus. ➤ Hand weeding and interculturing operation should be carry out. ➤ Apply irrigation to the crop at critical stage. ➤ For the control of sucking pest at 50 % flowering, spray 15 ml politrin OR 20 gm acephate OR 50 ml endoxacarb OR 25 ml profenophos in 10 liter of water.
Castor	Blight and Leaf spot	Flowering / Capsule formation	<ul style="list-style-type: none"> ➤ To control of blight and leaf spot, spray with Mancozeb 75 % WP 26 g/10 lit or copper oxichloride 50 % WP 40g/10 lit. ➤ Hand weeding and interculturing operation should be carry out. ➤ Apply irrigation to the crop at critical stage. ➤ Due to cloudy weather condition for the control of castor semi looper, spray 20 ml quinalphos (1.5 % powder) OR 40 gm carbaryl in 10 liter of water.
Vegetable crops	-	Sowing/ Flowering/	<ul style="list-style-type: none"> ➤ To the spray of agrochemicals during this period.

		Fruiting	<ul style="list-style-type: none"> ➤ Hand weeding and interculturing operation should be carry out. ➤ Apply irrigation to the crop at regular interval. ➤ Keep vegetable crop weed free, apply irrigation regularly and send vegetable fruit to the market often grading.
Livestock	-	-	<ul style="list-style-type: none"> ➤ To provide adequate food & clean water for drinking and keep the animals clean shed. ➤ Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Give stored fodder with mineral mixture. Fodder should be stored at safe place. ➤ Feed silage & hay material along with concentrate feed. ➤ Use chaff cutter for fodder. ➤ For the control of tick and other parasites, spray deltemethrin or aemitzaz 2 ml in 1 litre of water. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.

	<p style="text-align: center;">Gramin Krushi Mausam Seva (GKMS) Agrometeorological Cell Department of Agronomy Junagadh Agricultural University Junagadh-362001</p> <p>E-mail: amfujnd@yahoo.com Phone: 0285-2670139</p>	
---	---	---

Agromet Advisory Service Bulletin for Gir Somnath District of South Saurashtra Agro-Climatic Zone

(Issued jointly by Junagadh Agricultural University and India Meteorological Department)

(#AZADI KA AMRIT MAHOTSAV#)

Day:- Friday

Date: 26-11-2021

Agromet Advisory Service bulletin. No:- 069 (2021-22)

Observed Weather during last Five days (Dated 21 st to 25 th November, 2021)					Weather Parameters	Weather Forecast upcoming Five days (Valid for 27 th Nov. to 01 st Dec., 2021) until 08:30 hrs of 01-12-2021				
21/11	22/11	23/11	24/11	25/11	Date	27/11	28/11	29/11	30/11	01/12
0	0	0	0	0	Rainfall(mm)	0	0	0	0	0
34.0	35.0	32.2	32.7	34.6	Tmax (°C)	35	34	34	34	32
24.6	25.2	24.7	22.6	21.1	Tmin (°C)	18	19	19	19	18
0	0	0	0	0	Cloud Cover	4	5	5	7	8
91	83	92	90	61	RH-I (%)	53	48	53	45	83
78	83	91	66	58	RH-II (%)	25	27	29	33	33
1.9	1.9	1.9	3.7	3.7	Wind speed (km/hr)	8.7	9.2	11.6	17.5	19.1
99	99	99	99	99	Wind direction	158	22	45	79	58

Agro meteorological Advisories

General Advice	Rain fall	As per IMD Forecast, There is forecasting of no rainfall occurs during next five days. Sky will be remain partly cloudy to cloudy.
	Temperature	Maximum and minimum temperature likely to be in range of 32 - 35°C and 18 - 19°C, respectively.
	RH (%)	The Relative Humidity is likely to be in the range of 25 – 83 %.
	Wind Direction and Wind Speed	Direction and speed of wind is to be S, N, NE & E and 8.7 – 19.1 km/h, respectively.

- + **Standardized Precipitation Index:** - For the period of **14th October to 10th November, 2021** for Gir Somnath district considered under the mildly wet condition.
- + **AS per ERFS products Above Normal rainfall may occur in South Saurashtra region during 01st to 07th December, 2021.**
- + **MEGHDOOT Mobile Application:**- **Download** the MEGHDOOT mobile application for Agromet Advisory Services and information on weather forecasting, which will provide

weather based agricultural advice to farmers for their location, crops and livestock in their local language.


Click on the following link to download the MEGHDOOT application:

Android users: <https://play.google.com/store/apps/details?id=com.aas.meghdoot>

iOS users: <https://apps.apple.com/in/app/meghdoot/id1474048155>

- ❖ Due to pandemic of CORONA (COVID-19), Farmers should maintain social distancing and safety precaution while handling farm machines and labour in the field.
- ❖ Farmers are advised to prepare their field after harvesting of kharif crops for the sowing of rabi season crops timely.
- ❖ Interculturing and hand weeding should be carried out in *rabi* sowing crops.
- ❖ To apply fertilizer and agrochemicals as per recommendation dose in *rabi* crops.

NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 46
(12.11.2021 to 18.11.2021) over Agricultural region



Agricultural vigour is good over isolated pockets of Gujarat.

Source: <https://www.star.nesdis.noaa.gov>

Crop	Variety/ Pest / Insect	Stage	Agro met Advisories
Wheat	Late Sowing:- GJW-173, Lok-1, GW-11 and MP - 3288	Sowing / Germination / vegetative	<ul style="list-style-type: none"> ➤ Late sowing of wheat crop done during 26th Nov. to 10th December duration. ➤ To control Termite, Removal and destruction of preceding crop plant parts. ➤ Apply well decomposed FYM to soil. Use castor, neem or karanj cake as organic manure.

			<ul style="list-style-type: none"> ➤ Seed treatment with bifenthrin 10 EC 200 ml or fipronil 5 SC 600 ml or chlorpyriphos 20 EC 400 ml in 5 litre of water for 100 kg of wheat seed. ➤ Apply irrigation to the crop at regular interval. ➤ Apply chlorpyriphos 20 EC 1.5 litre or fipronil 5 SC 1.6 litre in 100 kg of sand, mix it and spread it and give the light irrigation or through irrigation (chemigation). ➤ At 30-35 days after sowing spray 20 ml 2-4-D ethyle easter or Metasulfuron 0.8 gm/10 liters of water to control broad leafy weeds in wheat crop. ➤ Hand weeding operation should be carry out.
Chickpea	Aphid	Germination / vegetative	<ul style="list-style-type: none"> ➤ Apply irrigation to the crop at regular interval. ➤ Set up yellow stick trap @ 10/ha to monitor aphid population. ➤ Hand weeding and interculturing operation should be carry out.
Coriander	Aphid	Sowing	<ul style="list-style-type: none"> ➤ Over irrigation should be avoided. ➤ Seed treatment of thiamethoxam 70 WS 4.2 gram/kg against aphid. ➤ Apply irrigation to the crop at regular interval. ➤ Hand weeding and interculturing operation should be carry out.
Garlic	Thrips	Germination / vegetative	<ul style="list-style-type: none"> ➤ Apply irrigation to the crop at regular interval.

			<ul style="list-style-type: none"> ➤ For management of thrips, apply schedule spraying of Beauveria bassiana 1.15 WP (min 2 x 106 cfu/g), first spray at initiation of pest infestation 0.0035% (30 g/10 l of water). Subsequent second 0.007% (60 g/10 l of water) and third 0.009% (80 g/10 l of water) spray at 10 days interval. ➤ To chemical control of thrips, spray Dimethoate 30 EC (10 ml/10 l. of water) or Quinalphos 25 EC (20 ml/10 l. of water). ➤ Hand weeding operation should be carry out.
Kharif Groundnut	-	Threshing	<ul style="list-style-type: none"> ➤ To done the operation of threshing in groundnut as per upcoming weather condition.
Pigeon pea	Sterility mosaic virus	Branching / Flowering	<ul style="list-style-type: none"> ➤ To Spray Propergite 57 EC or Spiromesifen 22.9 SC 10 ml/10 l control of the sterility mosaic virus. ➤ Hand weeding and interculturing operation should be carry out. ➤ Apply irrigation to the crop at critical stage. ➤ For the control of sucking pest at 50 % flowering, spray 15 ml politrin OR 20 gm acephate OR 50 ml endoxacarb OR 25 ml profenophos in 10 liter of water.
Sugarcane	-	Vegetative	<ul style="list-style-type: none"> ➤ To the spray of agrochemicals during this period. ➤ Hand weeding and interculturing operation should be carry out. ➤ Apply irrigation to the crop at critical stage.

Vegetable crops	-	Sowing/ Flowering/ Fruiting	<ul style="list-style-type: none"> ➤ To the spray of agrochemicals during this period. ➤ Hand weeding and interculturing operation should be carry out. ➤ Apply irrigation to the crop at regular interval. ➤ Keep vegetable crop weed free, apply irrigation regularly and send vegetable fruit to the market often grading.
Livestock	-	-	<ul style="list-style-type: none"> ➤ To provide adequate food & clean water for drinking and keep the animals clean shed. ➤ Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Give stored fodder with mineral mixture. Fodder should be stored at safe place. ➤ Feed silage & hay material along with concentrate feed. ➤ Use chaff cutter for fodder. ➤ For the control of tick and other parasites, spray deltemethrin or aemitaz 2 ml in 1 litre of water. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.

Technical Officer
Gramin Krushi Mausam Seva (GKMS)
J.A.U., Junagadh