



ગ્રામીણ કૃષિ મૌસમ સેવા

હવામાન આગાહી આધારીત કૃષિસલાહ બુલેટીન

ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ

તા.૦૪-૦૫-૨૦૧૯ થી તા.૦૮-૦૫-૨૦૧૯



(ગ્રામીણ કૃષિ મૌસમ સેવા, સૂકી ખેતી સંશોધન કેન્દ્ર, જી.કૃ.યુ., તરઘડિયા અને ભારત મૌસમ વિભાગ દ્વારા સંપન્ન)

કૃષિહવામાન બુલેટીન નં. ૧૦/૨૦૧૯

પાછલા અઠવાડિયાનું રાજકોટનું હવામાન

હવામાન પરિબલો	
વરસાદ(મી.મી)	0.0
મહત્તમ તાપમાન (સે.ગ્રે)	40.0-43.4
લઘુત્તમ તાપમાન (સે.ગ્રે)	22.3-24.6
ભેજ(%) સવાર	62-78
ભેજ(%) બપોરબાદ	19-39
પવનની ગતિ (કિમી/કલાક)	8.0-11.4

હવામાન આગાહી: પત્રક-ક

ઉત્તરસૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ-હવામાન આગાહી

તા.૦૪-૦૫-૨૦૧૯ થી ૦૮-૦૫-૨૦૧૯

કૃષિ સલાહ:

સામાન્ય	<p>➤ ભારત સરકારશ્રીના હવામાન ખાતા તરફથી મળેલ હવામાન માહિતી અનુસાર અત્રેના ગ્રામીણકૃષિ મૌસમ સેવા વિભાગ, જુનાગઢ કૃષિ યુનિવર્સિટી, તરઘડિયા તરફથી જણાવવામાં આવે છે કે, ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના</p> <p>રાજકોટ: જીલ્લામાં આગામી તા.૦૪-૦૫-૨૦૧૯ થી તા.૦૮-૦૫-૨૦૧૯ દરમિયાન હવામાન સુકું, ગરમ અને મુખત્વે વાદળછાયું રહેશે. પવનની દિશા પશ્ચિમ અને નૈઋત્યની રહેવાની અને પવનની ઝડપ ૧૮ થી ૨૨ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૪૦-૪૧ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન ૨૪-૨૫ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.</p>					
	તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
	પેરામીટર	૦૪-૦૫-૧૯	૦૫-૦૫-૧૯	૦૬-૦૫-૧૯	૦૭-૦૫-૧૯	૦૮-૦૫-૧૯
	વરસાદ(મી.મી)	0	0	0	0	0
	મહત્તમતાપમાન (સે.ગ્રે)	40	40	40	41	41

લઘુતમતાપમાન (સે.ગ્રે)	24	24	24	25	24
વાદળનીસ્થિતિ(ઓકટા)	1	0	1	0	0
મહતમભેજ(%) સવાર	68	69	72	68	73
લઘુતમભેજ (%) બપોરબાદ	17	15	18	17	15
પવનનીગતિ(કિમી/કલાક)	22	20	19	18	18
પવનનીદિશા(ડીગ્રી)	257	249	244	258	255

મોરબી: જીલ્લામાં આગામી તા.૦૪-૦૫-૨૦૧૯ થી તા.૦૮-૦૫-૨૦૧૯ દરમ્યાન હવામાન સુકું, ગરમ અને મુખત્વે વાદળછાયું રહેશે. પવનની દિશા નૈઋત્ય અને પશ્ચિમની રહેવાની અને પવનની ઝડપ ૮ થી ૨૦ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહતમ તાપમાન દિવસ દરમ્યાન ૪૦-૪૧ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમ્યાન ૨૪-૨૫ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૪-૦૫-૧૯	૦૫-૦૫-૧૯	૦૬-૦૫-૧૯	૦૭-૦૫-૧૯	૦૮-૦૫-૧૯
વરસાદ(મી.મી)	0	0	0	0	0
મહતમતાપમાન (સે.ગ્રે)	40	40	40	41	41
લઘુતમતાપમાન (સે.ગ્રે)	24	24	24	25	24
વાદળનીસ્થિતિ(ઓકટા)	3	3	0	0	0
મહતમભેજ(%) સવાર	60	68	62	57	45
લઘુતમભેજ (%) બપોરબાદ	10	16	18	16	14
પવનનીગતિ(કિમી/કલાક)	19	20	16	8	10
પવનનીદિશા(ડીગ્રી)	227	222	231	235	261

જામનગર: જીલ્લામાં આગામી તા.૦૪-૦૫-૨૦૧૯ થી તા.૦૮-૦૫-૨૦૧૯ દરમ્યાન હવામાન સુકું, ગરમ અને યોખ્ખુ રહેશે પવનની દિશા પશ્ચિમ અને નૈઋત્યની રહેવાની અને પવનની ઝડપ ૧૬ થી ૨૦ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહતમ તાપમાન દિવસ દરમ્યાન ૩૪-૩૫ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમ્યાન ૨૫-૨૭ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૪-૦૫-૧૯	૦૫-૦૫-૧૯	૦૬-૦૫-૧૯	૦૭-૦૫-૧૯	૦૮-૦૫-૧૯
વરસાદ(મી.મી)	0	0	0	0	0
મહતમતાપમાન (સે.ગ્રે)	34	35	35	34	34
લઘુતમતાપમાન (સે.ગ્રે)	27	27	26	26	25
વાદળનીસ્થિતિ(ઓકટા)	0	0	0	0	0

મહતમભેજ(%) સવાર	81	81	84	84	84
લઘુતમભેજ (%) બપોરબાદ	47	44	47	45	44
પવનનીગતિ(કિમી/કલાક)	18	20	18	16	18
પવનનીદિશા(ડીગ્રી)	254	252	245	250	241

દેવભૂમિ દ્વારકા: જીલ્લામાં આગામી તા.૦૪-૦૫-૨૦૧૯ થી તા.૦૮-૦૫-૨૦૧૯ દરમ્યાન હવામાન સુકું, ગરમ અને યોખ્ખુ રહશે પવનની દિશા પશ્ચિમ અને નૈઋત્યની રહેવાની અને પવનની ઝડપ ૧૪ થી ૧૮ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહતમ તાપમાન દિવસ દરમ્યાન ૩૧-૩૨ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમ્યાન ૨૫-૨૭ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૪-૦૫-૧૯	૦૫-૦૫-૧૯	૦૬-૦૫-૧૯	૦૭-૦૫-૧૯	૦૮-૦૫-૧૯
વરસાદ(મી.મી)	0	0	0	0	0
મહતમતાપમાન (સે.ગ્રે)	31	32	32	31	31
લઘુતમતાપમાન (સે.ગ્રે)	27	27	26	26	25
વાદળનીસ્થિતિ(ઓકટા)	0	0	0	0	0
મહતમભેજ(%) સવાર	83	82	86	87	86
લઘુતમભેજ (%) બપોરબાદ	44	43	47	44	42
પવનનીગતિ(કિમી/કલાક)	15	18	16	14	16
પવનનીદિશા(ડીગ્રી)	254	255	246	247	237

અમરેલી: જીલ્લામાં આગામી તા.૦૪-૦૫-૨૦૧૯ થી તા.૦૮-૦૫-૨૦૧૯ દરમ્યાન હવામાન સુકું, ગરમ અને મુખત્વે વાદળછાયું રહશે. પવનની દિશા પશ્ચિમ અને નૈઋત્યની રહેવાની અને પવનની ઝડપ ૧૫ થી ૧૮ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહતમ તાપમાન દિવસ દરમ્યાન ૪૦-૪૦ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમ્યાન ૨૪-૨૫ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.



તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૪-૦૫-૧૯	૦૫-૦૫-૧૯	૦૬-૦૫-૧૯	૦૭-૦૫-૧૯	૦૮-૦૫-૧૯
વરસાદ(મી.મી)	0	0	0	0	0
મહતમતાપમાન (સે.ગ્રે)	40	40	40	40	40
લઘુતમતાપમાન (સે.ગ્રે)	24	24	24	25	24
વાદળનીસ્થિતિ(ઓકટા)	2	1	0	0	0
મહતમભેજ(%) સવાર	74	75	76	69	76
લઘુતમભેજ(%) બપોરબાદ	27	32	34	27	30



પવનનીગતિ(કિમી/કલાક)	18	17	16	16	15
પવનનીદિશા(ડીગ્રી)	249	239	247	277	269



સુરેન્દ્રનગર: જીલ્લામાં આગામી તા.૦૪-૦૫-૨૦૧૯ થી તા.૦૮-૦૫-૨૦૧૯ દરમ્યાન હવામાન સુકું, ગરમ અને મુખત્વે વાદળછાયું રહેશે. પવનની દિશા પશ્ચિમની રહેવાની અને પવનની ઝડપ ૧૮ થી ૨૨ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૪૦-૪૧ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૨૪-૨૫ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૪-૦૫-૧૯	૦૫-૦૫-૧૯	૦૬-૦૫-૧૯	૦૭-૦૫-૧૯	૦૮-૦૫-૧૯
વરસાદ(મી.મી)	0	0	0	0	0
મહત્તમતાપમાન (સે.ગ્રે)	40	40	40	41	41
લઘુત્તમતાપમાન (સે.ગ્રે)	24	24	24	25	24
વાદળનીસ્થિતિ(ઓકટા)	1	1	1	0	0
મહત્તમભેજ(%) સવાર	59	62	63	59	65
લઘુત્તમભેજ(%) બપોરબાદ	17	15	16	18	15
પવનનીગતિ(કિમી/કલાક)	22	20	19	18	20
પવનનીદિશા(ડીગ્રી)	263	251	252	267	263

- ૦૪ થી ૦૮-મે-૨૦૧૯ હવામાન સુકું, ગરમ અને મુખત્વે વાદળછાયું રહેશે પવનની દિશા પશ્ચિમ અને નૈઋત્યની મોટાભાગે રહેવાની અને પવનની ઝડપ ૮ થી ૨૨ કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૪૦-૪૧ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૨૪-૨૫ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.
- શિયાળું પાકોને સંગ્રહ કરતા પહેલા તેના દાણાને વ્યવસ્થિત સફાઈ કરી, દાણામાં ૮ ટકા કરતા ઓછો ભેજ રહે ત્યાં સુધી સૂર્ય તાપમાં સુકવી પછી સંગ્રહ કરવો. તેમજ જે ગોડાઉનમાં સંગ્રહ કરવાનો હોય તે ગોડાઉનમાંથી સંગ્રહેલ જુના દાણાઓ બહાર કાઢી વ્યવસ્થિત સફાઈ કરવી. તેમજ દીવાલોમાં પડેલ તિરાડોનું રીપેરીંગ કરી પૂરી દેવી અને ગોડાઉનમાં ૦.૫ ટકા મેલાથીયોન દવાના દ્રાવણનો છંટકાવ કરવો. દવાના છંટકાવ પછી ૭ થી ૮ દિવસ ગોડાઉન હવાયુસ્ત રીતે બંધ રાખવું. તેમજ દાણા ભરવા માટેના કોથળા સુર્યતાપમાં સૂકવવા જેથી જીવાતોના ઈંડા અને કોશેટાનો નાશ થાય.
- સૂંકા અને ગરમ હવામાનને કારણે ઉનાળુ શાકભાજી અને અન્ય પાકોને સવાર અથવા સાંજના સમયે પિયત આપવું.
- શિયાળુ પાકોની કાપણી પછી ખેતરમાં ઊંડી ખેડ કરવી જેથી કીટકોના ઈંડા અને કોશેટા, તેમજ રોગકારકો જેવા કે ફૂગ, જીવાણુનો સુર્યતાપમાં નાશ થાય.

મુખ્ય પાક	પાક અવસ્થા	રોગ/જીવાત/જાત	કૃષિ સલાહ (૧૦ લીટર પાણીમાં મિશ્ર કરી ભલામણ કરેલ દવા છાંટવી)
ઉનાળુ મગ અને અડદ	શીંગ બંધાવવી, દાણા ભરાવવા	સફેદ માખી, લીલી ઈયળ	સફેદ માખીના નિયંત્રણ માટે પ્રોફેનોફોસ ૨૦ મી.લી.અને લીંબોળીનું તેલ ૫૦ મીલી અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી.પાણીમાં મિશ્ર કરી છંટકાવ કરવો. લીલી ઈયળના નિયંત્રણ માટે સ્પીનોસાઇ ૩ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.
ઉનાળુ મગફળી	પોપટા બંધાવા, દાણા ભરાવવા 	-	• પોપટા બંધાવાની અવસ્થાએ મગફળીને હળવું પિયત આપવું.
ઉનાળુ તલ	ફૂલ/બૈઠા બેસવા 	પાનવાળનાર (માથા બાંધનાર) ઈયળ લીલી ઈયળ	ઉનાળુ તલમાં પાનવાળનાર(માથા બાંધનાર) ઈયળના નિયંત્રણ માટે લીંબોળીનું તેલ ૫૦ મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને પ્રોફેનોફોસ ૩૦ મિલી અથવા લેમ્બડા સાયહેલોથ્રીન ૧૫ મિલી દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. લીલી ઈયળના નિયંત્રણ માટે સ્પીનોસાઇ ૩ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. ➤ તેમજ તલના પાકને ફૂલ અવસ્થાએ હળવું પિયત આપવું. ત્યારબાદ જમીનની પ્રત મુજબ ૧૦-૧૨ દિવસના ગાળે પિયત આપવું.વધારે પ્રમાણમાં ટૂંકા અંતરે પિયત આપવા નહિ. જરૂર કરતા વધારે ભેજથી તલમાં સૂકારો અને કાળીયો રોગ આવવાની શક્યતા રહે છે.
ઉનાળુ બાજરી	દાણા ભરાવવા	-	જરૂરીયાત મુજબ પિયત આપવું.
ભીંડા	ફૂલ/ફળ	સફેદમાખી, પાન કથીરી	થીપ્સના નિયંત્રણ માટે લીંબોળીનું તેલ ૫૦ મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને ઈમીડાક્લોપ્રીડ ૩૦.૫ ટકા ૪ મિલી અથવા એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. જો પાન કથીરીનો ઉપદ્રવ હોય તો ઇથીઓન ૩૦ મિલી પ્રતિ પંપ

			છંટકાવ કરવો.
તુરીયા	ફૂલ/ફળ	થ્રીપ્સ/ સફેદ માખી	ઉનાળુ શાકભાજીમાં સફેદમાખીના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦ મિલિ અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને ઇમીડાક્લોપ્રીડ ૩૦.૫ ટકા ૪ મિલી અથવા એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.
દૂધી			
ગલકા			
કારેલા			
ગુવાર			
કાકડી			કથીરી તળછારો
રીંગણી અને ટમેટી	ફૂલ/ફળ	સફેદ માખી	<ul style="list-style-type: none"> લીબોળીનું તેલ ૫૦મિલિ અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી. પાણીમાં મિશ્ર કરી છંટકાવ કરવો. જો વધુ ઉપદ્રવ હોય તો ટ્રાયઝોફોસ ૪૦ ઇ.સી. ૨૦ મી.લી. અથવા એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. ફળ અને ડોકાની ઇયળોનો ઉપદ્રવ હોય તો નુકશાન પામેલ ફળ અને ડોકા તોડીને જમીનમાં દાંટી દેવા અને ક્લોરાનટ્રાનીલીપ્રોલ ૧૮.૫ એસ.સી (રીનાક્ષીપાયર) દવા ૩ મિલી અથવા ડીડીવીપી ૭૬ ઇસી ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. રીંગણી અને ટમેટીના પાકમાં પાન કથીરીના નિયંત્રણ માટે પ્રોપરગાઈટ દવા ૧૦ મિલી/૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.
મરચી	ફૂલ/ફળ	પાનની થ્રીપ્સ→  પાનની કુક્કડ-  કથીરી	<p>ટ્રાયઝોફોસ ૪૦ ઇ.સી. ૨૦ મી.લી. અથવા એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ અથવા પ્રોફેનોફોસ ૨૦ મી.લી.અને લીબોળીનું તેલ ૫૦ મીલી અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી.પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p> <p>વાયરસના કારણે પીળા થઈ કુક્કાય ગયેલ છોડ ઉપાડી તેનો જમીનમાં દાટી નાશ કરવો. ત્યાં સફેદ માખીથી ફેલાતો હોવાથી તેના નિયંત્રણ માટે હોસ્તાથીયોન દવા ૨૦ મી.લી./૧૦ લી. પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p> <p>જો પાન કથીરીનો ઉપદ્રવ હોય તો ઇથીઓન ૩૦ મિલી પ્રતિ પંપ</p>

			છાન્કાવ કરવો.
આંબો	ફળનો વિકાસ	ફળ માખી	કેરી લખોટી જેવડા કદની થાય ત્યારે ૨ ગ્રામ નેપ્થેલીન એસેટીક એસીડ અને ૨ કી.ગ્રા. યુરીયાને ૧૦૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો જેથી કેરીનું ખરણ ઓછું થાય. આંબામાં ફળમાખીના નિયંત્રણ માટે ડીડીવીપી ૪ મિલી અને મીથાયલયુજીનોલ ૧૦ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.
લીંબુ	ફળનો વિકાસ	પાન કથીરી 	ઊંચા તાપમાનને કારણે લીંબુના બગીચામાં પાન કથીરીનો ઉપદ્રવ વધુ હોય તો તેના નિયંત્રણ માટે પ્રોપરગાઈટ દવા ૧૦ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.
પશુ/ મરઘા 			તાપમાન વધારે રહેવાના કારણે ઢોરને દિવસના સમયે છાયડામાં રાખવા અને મિનરલ મિક્ચર યુક્ત પીવાના સ્વચ્છ પાણીની વ્યવસ્થા કરાવી. દુધાળા પશુઓને ઉનાળામાં ઠંડક આપવા પાણીનો છંટકાવ કરવો. ખોરાકમાં લીલા ચારનું પ્રમાણ વધારવું. દુધાળા પશુઓના આઉને ઝીંક ઓક્સાઈડ અથવા બોરિક પવડરથી બરાબર સાફ કરવું. દુધાળા પશુઓના આઉને ઝીંક ઓક્સાઈડ અથવા બોરિક પવડરથી બરાબર સાફ કરવું. પશુઓમાં ગળસુંઢો અને ગાંઠીયા તાવના રોગના નિયંત્રણ માટે રસીકરણ કરાવવું. ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટ્રાઝ ૨ મી.લી. ૧ લીટર પાણીમાં નાખીને છાંટવી. વિટામીન એ તથા ડી પુરકો આપવા. ફરમિયાની દવા આપવી.

જુકૃયુ/ટેકનિકલ ઓફિસર.(સુખે)/એગ્રોમેટ/બુલેટીન/ /૨૦૧૯
તરઘડિયા(રાજકોટ), તારીખ: ૦૩/૦૫/૨૦૧૯.

પ્રતિ,
તંત્રીશ્રી/ડાયરેક્ટરશ્રી,
રાજકોટ,

ઉપરોક્ત માહિતી આપના દૈનિક સમાચાર પત્રોમાં, રેડીયોમાં ગામના ચોરાના કાર્યક્રમમાં તથા ટીવીના ગ્રામ્ય કાર્યક્રમમાં વિના મૂલ્યે પ્રસિધ્ધ કરાવવા નમ્ર વિનતી છે. સહકારની અપેક્ષા સહ આભાર.

ટેકનીકલ ઓફિસર
સુકી ખેતીસંશોધન કેન્દ્ર
જુનાગઢ કૃષિ નિવર્સિટી
તરઘડીયા



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :942756433



Agro met Advisory Service bulletin No.10-for the Rajkot district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 04 to 08-May - 2019)

Significant past weather for the preceding week of Rajkot

Rainfall (mm)	0.0
Maximum temperature (°C)	40.0-43.4
Minimum temperature (°C)	22.3-24.6
Relative Humidity (%) I	62-78
Relative Humidity (%) II	19-39
Wind speed (Kmph)	8.0-11.4


Weather forecast until 08-30 hrs. of 08/05/2019







(1) RAJKOT DISTRICT:

	04/05/19	05/05/19	06/05/19	07/05/19	08/05/19
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	40	40	40	41	41
Min.Temp.(°C)	24	24	24	25	24
Total CC (octa)	1	0	1	0	0
RH-I (%)	68	69	72	68	73
RH-II (%)	17	15	18	17	15
Wind Speed (kmph)	22	20	19	18	18
Wind Direction(deg.)	257	249	244	258	255

Agro meteorological Advisories:

General Advice	<ul style="list-style-type: none">• Rain fall: Dry, hot and mainly clear weather forecasted for next five days.• Temperature: Maximum and minimum temperature likely to be in range of 40-41°C and 24-25°C, respectively.• Relative Humidity: The RH-I and RH-II is likely to be in the range of 68-73% & 15-18, respectively.• Wind Direction and Speed: Direction and speed of wind is likely to be W, SW and 18-22 km/h, respectively.• Before storage of rabi crops produce, grains should be properly cleaned and it should be dried properly to the recommended level of moisture content. Godowns should be properly cleaned by removing all the grains and other storage matter of previous crop. Cracks and crevices of the walls should be cleaned and repaired and whitewashing should be done. Those who can afford are advised to spray 0.5 % malathion solution and keep the chamber closed for 7-8 days. The gunny bags should be dried in the sun
-----------------------	---

	<p>so that eggs and pupating stages of insects as well as inoculum of diseases get destroyed.</p> <ul style="list-style-type: none"> As forecast of dry & hot weather conditions, farmers are advised for light irrigation in all vegetable crops. Irrigation should be done in the morning hours or in the evening hours. High temperature is suitable deep ploughing of fields after harvesting of rabi crops is advised so that eggs and pupating stages of insects, fungus and weed seeds are destroyed. 		
Crops	Stage	Insect/Pest/ Variety	Advisory
Green gram	Pod formation/Development	White flies & heliothis	Under present weather condition infestation of heliothis and whiteflies were observed in summer sown green gram & black gram. To control of heliothis spray spinosad 3ml/10lit of water and to control of whiteflies spray acetamiprid 20 % SP@ 5.0 gm and 500 ml of 5 % neem seed extract or neem seed oil 50 ml with Profenofos 20 ml or or Lemdacylhelothrin 10 ml / 10 lit of water.
Black Gram	Pod formation/Development	White flies & heliothis	
Summer Groundnut	Pod formation/development		Apply light irrigation at pod formation and development stages without stress.
Summer Sesame	Flowering/capsule formation/development	Leaf webber/heliothis	For the control Leaf Webber spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with Profenofos 20 ml or or Lemdacylhelothrin 10 ml / 10 lit of water and for control of heliothis spray spinosad 3ml/10lit of water. To prevent Phytophthora blight and wilt do not over irrigate sesame, apply light irrigation as per need only.
Summer Pearl millet	Grain development	-	Apply irrigation as per need.
Lady finger	Flowering/fruiting	White flies & mites	Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear. Due to prevailing high temperature, light irrigation at short interval is advised.
Ridge gourd	Flowering & fruiting	Thrips & white flies/	In all vegetable crops thrips infestation observed above ETL. Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of
Bottle gourd			

Sponge guard		Mites	water for whiteflies control. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites.
Bitter guard			
Cluster bean		Downy mildew	For the control of downy mildew spray mencozeb 40g/pump.
Brinjal& Tomato	Flowering/ fruiting	Sucking pest → & Fruit borer 	To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.
Chili	Flowering/ fruiting	Thrips →  Leaf curling Virus →  mites	In present weather conditions attack of mites has been reported. To control mite infestation spray propargite 57 % EC 10 ml/10 litre of water. In present weather conditions, rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of Imidacloprid @ 5 ml per 10litre of water is advised for control the vector, when sky remains clear. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water
Mango	Fruit development	Fruit fly	Spray mixture of DDVP 4 ml and Methyleugenol 10 ml in 10 litres of water for control of fruit fly. Apply light irrigation as mango fruit size become gram grain size. Apply irrigation as per need.
Lemon	Fruit development	Mite  	Due to high temperature mites infestation in lemon orchard observed. To control mite infestation spray propargite 57 % EC 10/10 liter of water.
Livestock & poultry 	Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day. Do Vaccination for Haemorrhagic septicemia (H.S.) disease and Blue Quarter (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give		

	deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.
--	--

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :942756433



Agro met Advisory Service bulletin No.-10 for the Morbi district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 04 to 08-May - 2019)


Weather forecast until 08-30 hrs. of 08/05/2019







(2) MORBI DISTRICT:

	04/05/19	05/05/19	06/05/19	07/05/19	08/05/19
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	40	40	40	41	41
Min.Temp.(°C)	24	24	24	25	24
Total CC (octa)	3	3	0	0	0
RH-I (%)	60	68	62	57	45
RH-II (%)	10	16	18	16	14
Wind Speed (kmph)	19	20	16	8	10
Wind Direction(deg.)	227	222	231	235	261

Agro meteorological Advisories:

General Advice	<ul style="list-style-type: none">• Rain fall: Dry, hot and partly cloudy weather forecasted for next five days.• Temperature: Maximum and minimum temperature likely to be in range of 40-41°C and 24-25°C, respectively.• Relative Humidity: The RH-I and RH-II is likely to be in the range of 45-68% & 10-18, respectively.• Wind Direction and Speed: Direction and speed of wind is likely to be SW, W and 8-20 km/h, respectively.• Before storage of rabi crops produce, grains should be properly cleaned and it should be dried properly to the recommended level of moisture content. Godowns should be properly cleaned by removing all the grains and other storage matter of previous crop. Cracks and crevices of the walls should be cleaned and repaired and whitewashing should be done. Those who can afford are advised to spray 0.5 % malathion solution and keep the chamber closed for 7-8 days. The gunny bags should be dried in the sun so that eggs and pupating stages of insects as well as inoculum of diseases get destroyed.• As forecast of dry & hot weather conditions, farmers are advised for light irrigation in all vegetable crops. Irrigation should be done in the morning hours or in the evening hours.• High temperature is suitable deep ploughing of fields after harvesting of rabi crops is advised so that eggs and pupating stages of insects, fungus and weed seeds are destroyed.
-----------------------	--

Crops	Stage	Insect/Pest/ Variety	Advisory
Green gram	Pod formation/Development	White flies & heliothis	Under present weather condition infestation of heliothis and whiteflies were observed in summer sown green gram & black gram. To control of heliothis spray spinosad 3ml/10lit of water and to control of whiteflies spray acetamiprid 20 % SP@ 5.0 gm and 500 ml of 5 % neem seed extract or neem seed oil 50 ml with Profenofos 20 ml or or Lemdacyhelothrin 10 ml / 10 lit of water.
Black Gram	Pod formation/Development	White flies & heliothis	Under present weather condition infestation of heliothis and whiteflies were observed in summer sown green gram & black gram. To control of heliothis spray spinosad 3ml/10lit of water and to control of whiteflies spray acetamiprid 20 % SP@ 5.0 gm and 500 ml of 5 % neem seed extract or neem seed oil 50 ml with Profenofos 20 ml or or Lemdacyhelothrin 10 ml / 10 lit of water.
Summer Groundnut	Pod formation/development		Apply light irrigation at pod formation and development stages without stress.
Summer Sesame	Flowering/capsule formation/development	Leaf webber/heliothis	For the control Leaf Webber spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with Profenofos 20 ml or or Lemdacyhelothrin 10 ml / 10 lit of water and for control of heliothis spray spinosad 3ml/10lit of water. To prevent Phytophthora blight and wilt do not over irrigate sesame, apply light irrigation as per need only.
Summer Pearl millet	Grain development	-	Apply irrigation as per need.
Lady finger	Flowering/fruiting	White flies & mites	Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear. Due to prevailing high temperature, light irrigation at short interval is advised.
Ridge gourd	Flowering & fruiting	Thrips & white flies/	In all vegetable crops thrips infestation observed above ETL. Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites. For the control of downy mildew spray mencozeb 40g/pump.
Bottle gourd			
Sponge guard		Mites	
Bitter guard			
Cluster bean		Downy mildew	

Brinjal& Tomato	Flowering/ fruiting	Sucking pest → & Fruit borer 	<p>To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.</p>
Chili	Flowering/ fruiting	Thrips →  Leaf curling Virus →  mites	<p>In present weather conditions attack of mites has been reported. To control mite infestation spray propargite 57 % EC 10 ml/10 litre of water. In present weather conditions, rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of Imidacloprid @ 5 ml per 10litre of water is advised for control the vector, when sky remains clear.</p> <p>If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water</p>
Mango	Fruit development	Fruit fly	<p>Spray mixture of DDVP 4 ml and Methyleugenol 10 ml in 10 litres of water for control of fruit fly.</p> <p>Apply light irrigation as mango fruit size become gram grain size.</p> <p>Apply irrigation as per need.</p>
Lemon	Fruit development	Mite 	<p>Due to high temperature mites infestation in lemon orchard observed. To control mite infestation spray propargite 57 % EC 10/10 liter of water.</p> 
Livestock & poultry 	<p>Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day. Do Vaccination for Haemorrhagic septicemia (H.S.) disease and Blue Quarter (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-10 for the Jamnagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 04 to 08-May - 2019)


Weather forecast until 08-30 hrs of 08/05/2019







(3) JAMNAGAR DISTRICT:

	04/05/19	05/05/19	06/05/19	07/05/19	08/05/19
Rainfall (mm)	0	0	0	0	0
Max. Temp.(°C)	34	35	35	34	34
Min. Temp.(°C)	27	27	26	26	25
Total CC (octa)	0	0	0	0	0
RH-I (%)	81	81	84	84	84
RH-II (%)	47	44	47	45	44
Wind Speed (kmph)	18	20	18	16	18
Wind Direction(deg.)	254	252	245	250	241

Agro meteorological Advisories:-

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry, hot and clear sky weather forecasted for next five days. • Temperature: Maximum and minimum temperature likely to be in range of 34-35°C and 25-27°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 81-84% & 44-47, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be W, SW and 16-20 km/h, respectively. • Before storage of rabi crops produce, grains should be properly cleaned and it should be dried properly to the recommended level of moisture content. Godowns should be properly cleaned by removing all the grains and other storage matter of previous crop. Cracks and crevices of the walls should be cleaned and repaired and whitewashing should be done. Those who can afford are advised to spray 0.5 % malathion solution and keep the chamber closed for 7-8 days. The gunny bags should be dried in the sun so that eggs and pupating stages of insects as well as inoculum of diseases get destroyed. • As forecast of dry & hot weather conditions, farmers are advised for light irrigation in all vegetable crops. Irrigation should be done in the morning hours or in the evening hours. • High temperature is suitable deep ploughing of fields after harvesting of rabi crops is advised so that eggs and pupating stages of insects, fungus and weed seeds are destroyed. 		
Crops	Stage	Insect/Pest/ Variety	Advisory

Green gram	Pod formation/Development	White flies & heliothis	Under present weather condition infestation of heliothis and whiteflies were observed in summer sown green gram & black gram. To control of heliothis spray spinosad 3ml/10lit of water and to control of whiteflies spray acetamiprid 20 % SP@ 5.0 gm and 500 ml of 5 % neem seed extract or neem seed oil 50 ml with Profenofos 20 ml or or Lemdacyhelothrin 10 ml / 10 lit of water.
Black Gram	Pod formation/Development	White flies & heliothis	
Summer Groundnut	Pod formation/development		Apply light irrigation at pod formation and development stages without stress.
Summer Sesame	Flowering/capsule formation/development	Leaf webber/heliothis	For the control Leaf Webber spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with Profenofos 20 ml or or Lemdacyhelothrin 10 ml / 10 lit of water and for control of heliothis spray spinosad 3ml/10lit of water. To prevent Phytophthora blight and wilt do not over irrigate sesame, apply light irrigation as per need only.
Summer Pearl millet	Grain development	-	Apply irrigation as per need.
Lady finger	Flowering/fruiting	White flies & mites	Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear. Due to prevailing high temperature, light irrigation at short interval is advised.
Ridge gourd	Flowering & fruiting	Thrips & white flies/	In all vegetable crops thrips infestation observed above ETL. Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites. For the control of downy mildew spray mencozeb 40g/pump.
Bottle gourd			
Sponge guard		Mites	
Bitter guard			
Cluster bean		Downy mildew	

Brinjal& Tomato	Flowering/ fruiting	Sucking pest → & Fruit borer 	<p>To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.</p>
Chili	Flowering/ fruiting	Thrips →  Leaf curling Virus →  mites	<p>In present weather conditions attack of mites has been reported. To control mite infestation spray propargite 57 % EC 10 ml/10 litre of water. In present weather conditions, rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of Imidacloprid @ 5 ml per 10litre of water is advised for control the vector, when sky remains clear.</p> <p>If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water</p>
Mango	Fruit development	Fruit fly	<p>Spray mixture of DDVP 4 ml and Methyleugenol 10 ml in 10 litres of water for control of fruit fly.</p> <p>Apply light irrigation as mango fruit size become gram grain size.</p> <p>Apply irrigation as per need.</p>
Lemon	Fruit development	Mite 	<p>Due to high temperature mites infestation in lemon orchard observed. To control mite infestation spray propargite 57 % EC 10/10 liter of water.</p> 
Livestock & poultry 	<p>Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day. Do Vaccination for Haemorrhagic septicemia (H.S.) disease and Blue Quarter (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-10 for the Devbhumi Dwarka district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 04 to 08-May - 2019)


Weather forecast until 08-30 hrs of 08/05/2019







(4) DEVBHUMI DWARKA DISTRICT:

	04/05/19	05/05/19	06/05/19	07/05/19	08/05/19
Rainfall (mm)	0	0	0	0	0
Max. Temp.(°C)	31	32	32	31	31
Min. Temp.(°C)	27	27	26	26	25
Total CC (octa)	0	0	0	0	0
RH-I (%)	83	82	86	87	86
RH-II (%)	44	43	47	44	42
Wind Speed (kmph)	15	18	16	14	16
Wind Direction(deg.)	254	255	246	247	237

Agro meteorological Advisories:-

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry, hot and clear sky weather forecasted for next five days. • Temperature: Maximum and minimum temperature likely to be in range of 31-32°C and 25-27°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 82-87% & 42-47, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be W, SW and 14-18 km/h, respectively. • Before storage of rabi crops produce, grains should be properly cleaned and it should be dried properly to the recommended level of moisture content. Godowns should be properly cleaned by removing all the grains and other storage matter of previous crop. Cracks and crevices of the walls should be cleaned and repaired and whitewashing should be done. Those who can afford are advised to spray 0.5 % malathion solution and keep the chamber closed for 7-8 days. The gunny bags should be dried in the sun so that eggs and pupating stages of insects as well as inoculum of diseases get destroyed. • As forecast of dry & hot weather conditions, farmers are advised for light irrigation in all vegetable crops. Irrigation should be done in the morning hours or in the evening hours. • High temperature is suitable deep ploughing of fields after harvesting of rabi crops is advised so that eggs and pupating stages of insects, fungus and weed seeds are destroyed. 			
Crops	Stage	Insect/Pest/ Variety	Advisory	

Green gram	Pod formation/Development	White flies & heliothis	Under present weather condition infestation of heliothis and whiteflies were observed in summer sown green gram & black gram. To control of heliothis spray spinosad 3ml/10lit of water and to control of whiteflies spray acetamiprid 20 % SP@ 5.0 gm and 500 ml of 5 % neem seed extract or neem seed oil 50 ml with Profenofos 20 ml or or Lemdacyhelothrin 10 ml / 10 lit of water.
Black Gram	Pod formation/Development	White flies & heliothis	
Summer Groundnut	Pod formation/development		Apply light irrigation at pod formation and development stages without stress.
Summer Sesame	Flowering/capsule formation/development	Leaf webber/heliothis	For the control Leaf Webber spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with Profenofos 20 ml or or Lemdacyhelothrin 10 ml / 10 lit of water and for control of heliothis spray spinosad 3ml/10lit of water. To prevent Phytophthora blight and wilt do not over irrigate sesame, apply light irrigation as per need only.
Summer Pearl millet	Grain development	-	Apply irrigation as per need.
Lady finger	Flowering/fruiting	White flies & mites	Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear. Due to prevailing high temperature, light irrigation at short interval is advised.
Ridge gourd	Flowering & fruiting	Thrips & white flies/	In all vegetable crops thrips infestation observed above ETL. Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control.
Bottle gourd			
Sponge guard		Mites	If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites.
Bitter guard		Downy mildew	For the control of downy mildew spray mencozeb 40g/pump.
Cluster bean			

Brinjal& Tomato	Flowering/ fruiting	Sucking pest → & Fruit borer 	<p>To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.</p>
Chili	Flowering/ fruiting	Thrips →  Leaf curling Virus →  mites	<p>In present weather conditions attack of mites has been reported. To control mite infestation spray propargite 57 % EC 10 ml/10 litre of water. In present weather conditions, rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of Imidacloprid @ 5 ml per 10litre of water is advised for control the vector, when sky remains clear.</p> <p>If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water</p>
Mango	Fruit development	Fruit fly	<p>Spray mixture of DDVP 4 ml and Methyleugenol 10 ml in 10 litres of water for control of fruit fly.</p> <p>Apply light irrigation as mango fruit size become gram grain size.</p> <p>Apply irrigation as per need.</p>
Lemon	Fruit development	Mite 	<p>Due to high temperature mites infestation in lemon orchard observed. To control mite infestation spray propargite 57 % EC 10/10 liter of water.</p> 
Livestock & poultry 	<p>Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day. Do Vaccination for Haemorrhagic septicemia (H.S.) disease and Blue Quarter (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-10 for the Amreli district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 04 to 08-May - 2019)


Weather forecast until 08-30 hrs of 08/05/2019







(5) AMRELI DISTRICT:

	04/05/19	05/05/19	06/05/19	07/05/19	08/05/19
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	40	40	40	40	40
Min.Temp.(°C)	24	24	24	25	24
Total CC (octa)	2	1	0	0	0
RH-I (%)	74	75	76	69	76
RH-II (%)	27	32	34	27	30
Wind Speed (kmph)	18	17	16	16	15
Wind Direction(deg.)	249	239	247	277	269

Agro meteorological Advisories:

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry, hot and mainly clear weather forecasted for next five days. • Temperature: Maximum and minimum temperature likely to be in range of 40-40°C and 24-25°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 69-76% & 27-34, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be W, SW and 15-18 km/h, respectively. • Before storage of rabi crops produce, grains should be properly cleaned and it should be dried properly to the recommended level of moisture content. Godowns should be properly cleaned by removing all the grains and other storage matter of previous crop. Cracks and crevices of the walls should be cleaned and repaired and whitewashing should be done. Those who can afford are advised to spray 0.5 % malathion solution and keep the chamber closed for 7-8 days. The gunny bags should be dried in the sun so that eggs and pupating stages of insects as well as inoculum of diseases get destroyed. • As forecast of dry & hot weather conditions, farmers are advised for light irrigation in all vegetable crops. Irrigation should be done in the morning hours or in the evening hours. • High temperature is suitable deep ploughing of fields after harvesting of rabi crops is advised so that eggs and pupating stages of insects, fungus and weed seeds are destroyed. 		
Crops	Stage	Insect/Pest/ Variety	Advisory

Green gram	Pod formation/Development	White flies & heliothis	Under present weather condition infestation of heliothis and whiteflies were observed in summer sown green gram & black gram. To control of heliothis spray spinosad 3ml/10lit of water and to control of whiteflies spray acetamiprid 20 % SP@ 5.0 gm and 500 ml of 5 % neem seed extract or neem seed oil 50 ml with Profenofos 20 ml or or Lemdacyhelothrin 10 ml / 10 lit of water.
Black Gram	Pod formation/Development	White flies & heliothis	
Summer Groundnut	Pod formation/development		Apply light irrigation at pod formation and development stages without stress.
Summer Sesame	Flowering/capsule formation/development	Leaf webber/heliothis	For the control Leaf Webber spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with Profenofos 20 ml or or Lemdacyhelothrin 10 ml / 10 lit of water and for control of heliothis spray spinosad 3ml/10lit of water. To prevent Phytophthora blight and wilt do not over irrigate sesame, apply light irrigation as per need only.
Summer Pearl millet	Grain development	-	Apply irrigation as per need.
Lady finger	Flowering/fruiting	White flies & mites	Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear. Due to prevailing high temperature, light irrigation at short interval is advised.
Ridge gourd	Flowering & fruiting	Thrips & white flies/	In all vegetable crops thrips infestation observed above ETL. Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites. For the control of downy mildew spray mencozeb 40g/pump.
Bottle gourd			
Sponge guard		Mites	
Bitter guard			
Cluster bean		Downy mildew	

Brinjal& Tomato	Flowering/ fruiting	Sucking pest → & Fruit borer 	<p>To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.</p>
Chili	Flowering/ fruiting	Thrips →  Leaf curling Virus →  mites	<p>In present weather conditions attack of mites has been reported. To control mite infestation spray propargite 57 % EC 10 ml/10 litre of water. In present weather conditions, rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of Imidacloprid @ 5 ml per 10litre of water is advised for control the vector, when sky remains clear.</p> <p>If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water</p>
Mango	Fruit development	Fruit fly	<p>Spray mixture of DDVP 4 ml and Methyleugenol 10 ml in 10 litres of water for control of fruit fly.</p> <p>Apply light irrigation as mango fruit size become gram grain size.</p> <p>Apply irrigation as per need.</p>
Lemon	Fruit development	Mite  	<p>Due to high temperature mites infestation in lemon orchard observed. To control mite infestation spray propargite 57 % EC 10/10 liter of water.</p>
Livestock & poultry 	<p>Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day. Do Vaccination for Haemorrhagic septicemia (H.S.) disease and Blue Quarter (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell:94277 28523



Agro met Advisory Service bulletin No.-10 for the Surendranagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 04 to 08-May - 2019)


Weather forecast until 08-30 hrs of 08/05/2019







(6) SURENDRANAGAR DISTRICT:

	04/05/19	05/05/19	06/05/19	07/05/19	08/05/19
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	40	40	40	41	41
Min.Temp.(°C)	24	24	24	25	24
Total CC (octa)	1	1	1	0	0
RH-I (%)	59	62	63	59	65
RH-II (%)	17	15	16	18	15
Wind Speed (kmph)	22	20	19	18	20
Wind Direction(deg.)	263	251	252	267	263

Agro meteorological advisories:

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry, hot and mainly clear weather forecasted for next five days. • Temperature: Maximum and minimum temperature likely to be in range of 40-41°C and 24-25°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 59-65% & 15-18, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be W and 18-22 km/h, respectively. • Before storage of rabi crops produce, grains should be properly cleaned and it should be dried properly to the recommended level of moisture content. Godowns should be properly cleaned by removing all the grains and other storage matter of previous crop. Cracks and crevices of the walls should be cleaned and repaired and whitewashing should be done. Those who can afford are advised to spray 0.5 % malathion solution and keep the chamber closed for 7-8 days. The gunny bags should be dried in the sun so that eggs and pupating stages of insects as well as inoculum of diseases get destroyed. • As forecast of dry & hot weather conditions, farmers are advised for light irrigation in all vegetable crops. Irrigation should be done in the morning hours or in the evening hours. • High temperature is suitable deep ploughing of fields after harvesting of rabi crops is advised so that eggs and pupating stages of insects, fungus and weed seeds are destroyed. 			
Crops	Stage	Insect/Pest/ Variety	Advisory	

Green gram	Pod formation/Development	White flies & heliothis	Under present weather condition infestation of heliothis and whiteflies were observed in summer sown green gram & black gram. To control of heliothis spray spinosad 3ml/10lit of water and to control of whiteflies spray acetamiprid 20 % SP@ 5.0 gm and 500 ml of 5 % neem seed extract or neem seed oil 50 ml with Profenofos 20 ml or or Lemdacyhelothrin 10 ml / 10 lit of water.
Black Gram	Pod formation/Development	White flies & heliothis	
Summer Groundnut	Pod formation/development		Apply light irrigation at pod formation and development stages without stress.
Summer Sesame	Flowering/capsule formation/development	Leaf webber/heliothis	For the control Leaf Webber spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with Profenofos 20 ml or or Lemdacyhelothrin 10 ml / 10 lit of water and for control of heliothis spray spinosad 3ml/10lit of water. To prevent Phytophthora blight and wilt do not over irrigate sesame, apply light irrigation as per need only.
Summer Pearl millet	Grain development	-	Apply irrigation as per need.
Lady finger	Flowering/fruiting	White flies & mites	Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear. Due to prevailing high temperature, light irrigation at short interval is advised.
Ridge gourd	Flowering & fruiting	Thrips & white flies/	In all vegetable crops thrips infestation observed above ETL. Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control.
Bottle gourd			
Sponge guard		Mites	If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites.
Bitter guard		Downy mildew	For the control of downy mildew spray mencozeb 40g/pump.
Cluster bean			

Brinjal& Tomato	Flowering/ fruiting	Sucking pest → & Fruit borer 	<p>To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.</p>
Chili	Flowering/ fruiting	Thrips →  Leaf curling Virus →  mites	<p>In present weather conditions attack of mites has been reported. To control mite infestation spray propargite 57 % EC 10 ml/10 litre of water. In present weather conditions, rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of Imidacloprid @ 5 ml per 10litre of water is advised for control the vector, when sky remains clear.</p> <p>If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water</p>
Mango	Fruit development	Fruit fly	<p>Spray mixture of DDVP 4 ml and Methyleugenol 10 ml in 10 litres of water for control of fruit fly.</p> <p>Apply light irrigation as mango fruit size become gram grain size.</p> <p>Apply irrigation as per need.</p>
Lemon	Fruit development	Mite 	<p>Due to high temperature mites infestation in lemon orchard observed. To control mite infestation spray propargite 57 % EC 10/10 liter of water.</p> 
Livestock & poultry 	<p>Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day. Do Vaccination for Haemorrhagic septicemia (H.S.) disease and Blue Quarter (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.