



ગ્રામીણ કૃષિ મૌસમ સેવા
હવામાન આગાહી આધારીત કૃષિસલાહ બુલેટીન
ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ
તા.૦૫-૦૯-૨૦૧૮ થી તા.૦૯-૦૯-૨૦૧૮



(ગ્રામીણ કૃષિ મૌસમ સેવા, સૂકી ખેતી સંશોધન કેન્દ્ર, જી.કૃ.યુ., તરઘડિયા અને ભારત મૌસમ વિભાગ દ્વારા સંપન્ન)

કૃષિહવામાન બુલેટીન નં. ૪૫/૨૦૧૮

પાછલા અઠવાડિયાનું રાજકોટનું હવામાન

હવામાન પરિબલો	
વરસાદ(મી.મી)	20.1
મહત્તમ તાપમાન (સે.ગ્રે)	27.4-29.9
લઘુતમ તાપમાન (સે.ગ્રે)	22.2-22.8
ભેજ(%) સવાર	88-91
ભેજ(%) બપોરબાદ	79-90
પવનની ગતિ (કિમી/કલાક)	4.6-6.4

હવામાન આગાહી: પત્રક-ક

ઉત્તરસૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ-હવામાન આગાહી

તા.૦૫-૦૯-૨૦૧૮ થી ૦૯-૦૯-૨૦૧૮

કૃષિ સલાહ:

સામાન્ય	<p>➤ ભારત સરકારશ્રીના હવામાન ખાતા તરફથી મળેલ હવામાન માહિતી અનુસાર અત્રેના ગ્રામીણકૃષિ મૌસમ સેવા વિભાગ, જુનાગઢ કૃષિ યુનિવર્સિટી, તરઘડિયા તરફથી જાણાવવામાં આવે છે કે, ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના</p> <p>રાજકોટ: જીલ્લામાં આગામી તા.૦૫-૦૯-૨૦૧૮ થી તા ૦૯-૦૯-૨૦૧૮ દરમિયાન હવામાન ભેજવાળું અને વાદળછાયું રહેશે. પવનની દિશા પશ્ચિમ અને નૈઋત્યની રહેવાની અને પવનની ઝડપ ૯ થી ૧૮ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૨૯-૩૦ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમિયાન ૨૨-૨૩ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.</p>																																				
	<table border="1"><thead><tr><th>તારીખ</th><th>દિવસ-૧</th><th>દિવસ-૨</th><th>દિવસ-૩</th><th>દિવસ-૪</th><th>દિવસ-૫</th></tr></thead><tbody><tr><td>પેશામીટર</td><td>૦૫-૦૯-૧૮</td><td>૦૬-૦૯-૧૮</td><td>૦૭-૦૯-૧૮</td><td>૦૮-૦૯-૧૮</td><td>૦૯-૦૯-૧૮</td></tr><tr><td>વરસાદ(મી.મી)</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></tr><tr><td>મહત્તમતાપમાન (સે.ગ્રે)</td><td>29</td><td>30</td><td>30</td><td>30</td><td>29</td></tr><tr><td>લઘુતમતાપમાન (સે.ગ્રે)</td><td>23</td><td>23</td><td>23</td><td>23</td><td>22</td></tr><tr><td>વાદળનીસ્થિતિ(ઓકટા)</td><td>5</td><td>7</td><td>3</td><td>2</td><td>3</td></tr></tbody></table>	તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫	પેશામીટર	૦૫-૦૯-૧૮	૦૬-૦૯-૧૮	૦૭-૦૯-૧૮	૦૮-૦૯-૧૮	૦૯-૦૯-૧૮	વરસાદ(મી.મી)	0	0	0	0	0	મહત્તમતાપમાન (સે.ગ્રે)	29	30	30	30	29	લઘુતમતાપમાન (સે.ગ્રે)	23	23	23	23	22	વાદળનીસ્થિતિ(ઓકટા)	5	7	3	2	3
તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫																																
પેશામીટર	૦૫-૦૯-૧૮	૦૬-૦૯-૧૮	૦૭-૦૯-૧૮	૦૮-૦૯-૧૮	૦૯-૦૯-૧૮																																
વરસાદ(મી.મી)	0	0	0	0	0																																
મહત્તમતાપમાન (સે.ગ્રે)	29	30	30	30	29																																
લઘુતમતાપમાન (સે.ગ્રે)	23	23	23	23	22																																
વાદળનીસ્થિતિ(ઓકટા)	5	7	3	2	3																																

મહત્તમભેજ(%) સવાર	80	84	80	82	85
લઘુત્તમભેજ (%) બપોરબાદ	60	62	50	49	54
પવનનીગતિ(મી/કલાક)	18	13	13	10	9
પવનનીદિશા(ડીગ્રી)	248	241	236	224	227

જામનગર: જીલ્લામાં આગામી તા.૦૫-૦૯-૨૦૧૮ થી તા.૦૯-૦૯-૨૦૧૮ દરમ્યાન હવામાન ભેજવાળું અને વાદળછાયું રહેશે. પવનની દિશા પશ્ચિમ અને નૈઋત્યની રહેવાની અને પવનની ઝડપ ૧૧ થી ૨૫ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૨૮-૨૯ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૨૫-૨૬ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૫-૦૯-૧૮	૦૬-૦૯-૧૮	૦૭-૦૯-૧૮	૦૮-૦૯-૧૮	૦૯-૦૯-૧૮
વરસાદ(મી.મી)	0	0	0	0	0
મહત્તમતાપમાન (સે.ગ્રે)	29	29	28	29	28
લઘુત્તમતાપમાન (સે.ગ્રે)	25	25	25	26	26
વાદળનીસ્થિતિ(ઓકટા)	6	7	3	3	4
મહત્તમભેજ(%) સવાર	77	79	76	77	79
લઘુત્તમભેજ(%) બપોરબાદ	67	68	57	59	62
પવનનીગતિ(કીમી/કલાક)	25	20	19	13	11
પવનનીદિશા(ડીગ્રી)	249	242	243	224	214

અમરેલી: જીલ્લામાં આગામી તા.૦૫-૦૯-૨૦૧૮ થી તા.૦૯-૦૯-૨૦૧૮ દરમ્યાન હવામાન ભેજવાળું અને વાદળછાયું રહેશે. પવનની દિશા પશ્ચિમ અને નૈઋત્યની રહેવાની અને પવનની ઝડપ ૭ થી ૧૪ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૩૦-૩૧ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૨૨-૨૨ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૫-૦૯-૧૮	૦૬-૦૯-૧૮	૦૭-૦૯-૧૮	૦૮-૦૯-૧૮	૦૯-૦૯-૧૮
વરસાદ(મી.મી)	0	0	0	0	0
મહત્તમતાપમાન (સે.ગ્રે)	31	31	31	30	31
લઘુત્તમતાપમાન (સે.ગ્રે)	22	22	22	22	22
વાદળનીસ્થિતિ(ઓકટા)	5	6	4	2	3
મહત્તમભેજ(%) સવાર	84	87	83	82	85
લઘુત્તમભેજ(%) બપોરબાદ	65	65	61	55	55




પવનનીગતિકિમી/કલાક)	14	12	12	9	7
પવનનીદિશા(ડીગ્રી)	253	240	250	247	249


સુરેન્દ્રનગર: જીલ્લામાં આગામી તા.૦૫-૦૯-૨૦૧૮ થી તા.૦૯-૦૯-૨૦૧૮ દરમ્યાન હવામાન ભેજવાળું અને વાદળછાયું રહેશે. પવનની દિશા નૈઋત્યની રહેવાની અને પવનની ઝડપ ૧૨ થી ૧૯ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૨૯-૩૦ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૨૨-૨૩ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૫-૦૯-૧૮	૦૬-૦૯-૧૮	૦૭-૦૯-૧૮	૦૮-૦૯-૧૮	૦૯-૦૯-૧૮
વરસાદ(મી.મી)	0	0	0	0	0
મહત્તમતાપમાન (સે.ગ્રે)	29	30	30	30	29
લઘુત્તમતાપમાન (સે.ગ્રે)	23	23	23	23	22
વાદળનીસ્થિતિ(ઓકટા)	6	7	2	1	2
મહત્તમભેજ(%) સવાર	79	79	77	76	79
લઘુત્તમભેજ(%) બપોરબાદ	56	55	44	44	45
પવનનીગતિકિમી/કલાક)	19	16	14	12	12
પવનનીદિશા(ડીગ્રી)	246	241	236	239	247



- ૦૫ થી ૦૯-સપ્ટેમ્બર દરમ્યાન હવામાન ભેજવાળું અને વાદળછાયું રહેશે પવનની દિશા પશ્ચિમ અને નૈઋત્યની રહેવાની અને પવનની ઝડપ ૭ થી ૧૯ કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૨૯-૩૧ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૨૨-૨૩ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.
- સુરેન્દ્રનગર અને મોરબી જીલ્લાનાં જે વિસ્તારમાં પાછોતરી વાવણી થયેલ માટે તલ પૂર્વા ૧, દિવેલા GC-3 , GCH-4, 6 & 7, તુવેર બી.ડી.એન.-૨, વૈશાલી, જી.જી.પી-૧, ચારની જુવાર એસ-૧૦૪૯, ગુન્ધરી, GFS-3, GAFS-11, CSV-21F, સોયાબીન ગુજ.-૧, અને ૩ ની વાવણી કરવી.

મુખ્ય પાક	પાક અવસ્થા	રોગ/જીવાત/જાત	કૃષિ સલાહ
			(૧૦ લીટર પાણીમાં મિશ્ર કરી ભલામણ કરેલ દવા છાંટવી)

મગફળી	ફૂલ/સુયા	<p>સફેદ ફૂગ</p>  <p>ચીતરી અને ગેરુ</p>  <p>સફેદ મુંડા</p> 	<ul style="list-style-type: none"> ➤ મગફળીમાં સફેદ ફૂગનો ઉપદ્રવ અટકાવવા ટ્રાયકોડર્મા ફૂગ ૨.૫ કી.ગ્રા./હે. મુજબ ગળતીયા દેશી ખાતર અથવા દિવેલાના ખોળ ૫૦૦ કી.ગ્રા./હે. મિશ્ર કરીને જમીનમાં આપવું. ➤ પાનના ટપકાના રોગના આગોતરા નિયંત્રણ માટે હેક્ઝાકોનાઝોલ ૧૫ મિલી અથવા ટેબ્યુંકોનાઝોલ ૧૦ મિલી ૧૦ લીટર પાણીમા મિશ્ર કરી છંટકાવ કરવો. ➤ ફોરેટ ૧૦ ટકા દાણાદાર દવા ૨૫ કી.ગ્રા. પ્રતિ હેક્ટરે જમીનમાં પુરતો ભેજ હોય ત્યારે નાખવી અથવા ક્લોરોપાયરીફોસ દવા ૪ લીટર પ્રતિ હેક્ટરે પિયત પાણી સાથે અથવા ક્લોરોપાયરીફોસ દવા ૨૫ મિલી ૧૦ લીટર પાણીમા મિશ્ર કરી જમીનમાં થડની ફરતે રેડવું. ➤ આગોતરી મગફળી પીળી પડેલ હોય તો ૮-૧૦ કી.ગ્રા. એમોનિયમ સલ્ફેટ ખાતર પ્રતિ વીધા આપવું. તેમજ વરાપે હીરાકસી ૧૫૦ ગ્રામ અને લીંબુના ફૂલ ૧૫ ગ્રામ ૧ પમ્પમાં નાખીને છંટકાવ કરવો.
કપાસ	ચાપવા	<p>શ્રીખસ</p> <p>ગુલાબી ઇયળ</p>	<ul style="list-style-type: none"> ➤ આ હવામાન પરિસ્થિતિમાં કપાસમાં શ્રીખસનો વધુ ઉપદ્રવ જોવા મળે છે જેના નિયંત્રણ માટે પ્રોફેનોફોસ ૫૦ % અથવા ઈથીઓન ૫૦ % ઇસી ૨૦ મીલી સાથે લીંબોળીનું તેલ ૫૦ મીલી અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. ➤ આગોતરા કપાસમાં ગુલાબી ઇયળના ઉપદ્રવની મોજણી માટે નર કુદાને એકત્રીત કરવા માટે હેક્ટરે ૬-૭ ગુલાબી ઇયળના ફેરોમેન ટ્રેપ મુકવા અને જો વધુ ઉપદ્રવ જણાય તો ૪૦ ની સંખ્યામાં ફેરોમેન ટ્રેપ ગોઠવવાથી ઇયળોનું શરુઆતની અવસ્થાથી જ

			<p>નિયંત્રણ કરી શકાય. કપાસમાં ગુલાબી ઈયળથી નુકશાન પામેલ ફૂલ અને ચાપવા તોડીને જમીનમાં દાટી દેવા. વધુ નુકશાન હોય તો પ્રોફેનોફોસ ૫૦% ઇ.સી. ૨૦ મી.લી. અથવા સાયપરમેથ્રીન ૧૦ મિલી અને બીવેરીયા બાસીઆના ફૂગનો પાઉડર ૪૦ ગ્રામ ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p> <p>➤ સુરેન્દ્રનગર જીલ્લામાં બાકી રહી ગયેલ વિસ્તારમાં વાવણી લાયક વરસાદ થયે તુવેર, દિવેલા, મગ, ચોળી, અડદ, તલ, અજમો, સુવા, વરયાળી અને ઘાસચારાની જુવારનું વાવેતર કરવું.</p>
તલ	ફૂલ/કેપ્સુલ	પાનવાળનાર(માથા બાંધનાર) ઈયળ	<p>કવીનાલફોસ ૨૦ મિલી અથવા કાર્ટેપ હાયડ્રોકલોરાઈડ ૧૫ ગ્રામ ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p>  <p>આગોતરી વાવણીના તલમાં ભુકીછારો અને ફાયટોપ્થોરા બ્લાઈટ રોગનો ઉપદ્રવ ના થાય તે માટે રીડોમિલ દવાનો ૨૦ મિલી/૧૦ લિટર પાણીમાં મિશ્ર કરી ૧૦ દિવસે બે છંટકાવ કરવા.</p>
બાજરી	ગાભ	-	<p>➤ વરસાદ ખેંચાયેલ હોય જમીનમાં ભેજ જળવાય રહે તે માટે આંતરખેડ અને નિંદામણ કરવું.</p>
જુવાર/ મકાઈ	વૃદ્ધિ	કુપરું કોરી ખાનાર ઈયળ	<p>૨૫-૩૦ દિવસની જુવારમાં યુરીયા ૮૬ કી.ગ્રા./હે આપવું. તેમજ કુપરાની ઈયળના ઉપદ્રવનું સતત નિરીક્ષણ કરતા રહેવું. જરૂર જણાયે વધુ ઉપદ્રવ હોય તો સાયપરમેથ્રીન ૨૫ ટકા ૧૦ મીલી અને લીંબોળીનું તેલ ૫૦ મીલી અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p>

કઠોળ પાકો	ફૂલ	થડ માખી	થડમાખીના નિયંત્રણ માટે ડીડીવીપી ૭ મિલી અને ફેન્વાલીરેટ ૧૫ મિલી/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. આગતર મગ અને ચોળીની પાકેલ શીંગોની સમયસર વીણી કરવી.
દિવેલા	વરસાદ ખેયાતા હવે વાવણી લાયક વરસાદ થયે દિવેલાની GC-3, GCH-4,6, અથવા 7 જાત પસંદ કરવી.		
તુવેર	વરસાદ ખેયાતા હવે વાવણી લાયક વરસાદ થયે બીડીએન-૨ અને વૈશાલી અથવા GJP-1 જાતનું વાવેતર કરવું અને પાયામાં ૨૫ કી.ગ્રા. નાઈટ્રોજન અને ૫૦ કી.ગ્રા. ફોસ્ફરસ આપવો.		
શાકભાજી	ધરું ઉછેર/રોપણી વાવેતર	મરચી,ટમેટી, રીંગણ, અને કોબીઝના ધરું તૈયાર કરવા (ગ્રીન શેડ નેટમાં અને નેટની ઊંચાઈ ૬.૫ ફૂટ રાખવી) ગાદી ક્યારા બનાવી તેમાં બીજનું વાવેતર કરવું. જો ધરું તૈયાર હોય તો મરચી,ટમેટી, રીંગણ, અને કોબીઝની વરસાદ થયે રોપણી કરવી. તુરિયા, કરેલા, દુધી, ગલકા,પરવળ, કાકડી, ગુવાર, ભીંડો, ચોળી, તુવેર, મૂળા,તાંદળજો, પાલકનું વાવેતર કરવું.	
ભીંડા	વૃદ્ધિ/કુલ/ શીંગ	શ્રીપ્સ,સફેદ માખી શીંગ કોરી ખાનાર ઈયળ પાન કથીરી	<ul style="list-style-type: none"> ➤ તેના નીયંત્રણ માટે ટ્રાયએફેસ ૪૦ ટકા ૨૫ મિલી અને લીબોળીનું તેલ ૫૦ મિલી અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મિલી/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. ➤ તેના ફળ અને ડોકા કોરી ખાનાર ઈયળનો ઉપદ્રવ વધુ હોય તો સ્પીનોસાડ ૩ મિલી /૧૦ લીટર પાણીમા ઓગાળી છંટકાવ કરવો. ➤ પાન કથીરીનો ઉપદ્રવ વધુ હોય તો તેના નિયંત્રણ માટે પ્રોપરગાઈટ દવા ૧૦ મિલી ૧૦ લીટર પાણીમા મિશ્ર કરી છંટકાવ કરવો.
તુરીયા, દુધી, ગલકા કારેલા અને કાકડી	વૃદ્ધિ/કુલ/ફળ	પાન કોરીયું , શ્રીપ્સ, સફેદ માખી, પાન કથીરી	<ul style="list-style-type: none"> ➤ લીબોળીનો અર્ક ૫૦૦ મિલી અથવા લીબોળીનું તેલ ૫૦ મીલીનો છંટકાવ ૧૫ દિવસના અંતરે ચાર વખત કરવો. વધુ ઉપદ્રવ હોય તો એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ / ૧૦ લીટર પાણીમા મિશ્ર કરી છંટકાવ કરવો. ➤ પાન કથીરીના નિયંત્રણ માટે પ્રોપરગાઈટ દવા ૧૦ મિલી ૧૦ લીટર પાણીમા મિશ્ર કરી છંટકાવ કરવો.

		તલછારો	<ul style="list-style-type: none"> મેન્કોઝેબ ૨૭ ગ્રામ અથવા ક્લોરોથેલોનીલ ૨૭ ગ્રામ અથવા ફોઝેટાઈલ ૨૦ ગ્રામના ૧૫ દિવસના અંતરે ત્રણ છંટકાવ કરવા.
ગુવાર	વૃદ્ધિ/કુલ/શીંગ	થ્રીપ્સ, સફેદ માખી	<ul style="list-style-type: none"> તેના નિયંત્રણ માટે એસીટામાપ્રીડ ૨૦ ટકા ૫ ગ્રામ અને લીબોળીનું તેલ ૫૦ મીલી અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.
રીંગણી અને ટમેટી	ફુલ/ફળ	થ્રીપ્સ, સફેદ માખી, ફળ અને ડોકા કોરી ખાનાર ઈયળ	<ul style="list-style-type: none"> લીબોળીનું તેલ ૫૦ મિલી અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. જો વધુ ઉપદ્રવ હોય તો ઈથીઓન ૫૦ ટકા ૨૦ મિલી /૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. રીંગણી અને ટમેટીમાં ફળ અને ડોકા કોરી ખાનાર ઈયળનો ઉપદ્રવ વધુ હોય તો રીનાક્ષીપાયર ૧૮.૫ ટકા ૪ મિલી /૧૦ લીટર પાણીમાં ઓગાળી છંટકાવ કરવો.
મરચી	વૃદ્ધિ/કુલ/ફળ	થ્રીપ્સ સફેદ માખી 	<ul style="list-style-type: none"> પ્રોફેનોફોસ ૨૦ મીલી અને લીબોળીનું તેલ ૫૦ મિલી અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મિલી ૧૦ લી. પાણીમાં મિશ્ર કરી છંટકાવ કરવો.
લીંબુ	ફળનો વિકાસ	સાયલા અને થ્રીપ્સ  બદામી ટપકાં પાનકોરીયુ	<p>તેના નિયંત્રણ માટે ઈમીડાક્લોપ્રીડ ૩૦.૫ ટકા ૫ મીલી અને લીબોળીનું તેલ ૫૦ મીલી અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. વધુ ઉપદ્રવ હોય તો ટ્રાયઝોફોસ ૪૦ ટકા ૨૦/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p> <p>બદામી ટપકાના નિયંત્રણ માટે કોપર ઓક્સીક્લોરાઈડ ૪૦ ગ્રામ અને ૧.૫ ગ્રામ સ્ટ્રેપ્ટોસાયક્લીનના બે છંટકાવ ૧૫ દિવસના ગળે કરવા.</p> <p>પાનકોરીયુના નિયંત્રણ માટે ડીડીવીપી ૧૦ મી.લી. ૧૦ લિટર પાણીમાં મિશ્ર કરીને છાંટવી.</p>

પશુ/મરઘા



દુધાળા પશુના આઉને ઝીંક ઓક્સાઈડ અથવા બોરીક પાવડર થી બરાબર સાફ કરવું. તેમજ ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટ્રાલ ૨ મી.લી. એક લીટર પાણીમાં નાખીને છાંટવી અને કરમીયાની ગોળી આપવી તેમજ ટીબી અને જે.ડી.ના રોગની પરખ કરાવવી તેમજ બુસેલ્લાનું રસીકરણ કરાવવું. વરસાદના સમયમાં પશુઓને સુરક્ષિત રાખવા. ચોમાસા દરમ્યાન પશુઓના રહેઠાણની જગ્યાની વ્યવસ્થિત સફાઈ રાખવી અને માખીઓના ઉપદ્રવથી બચવા ફિનાઈલ છાંટવું.

ખુલ્લામાં રાખેલ ઘાસસારો વરસાદથી બગડે નહિ તે માટે જરૂરી પગલા ભરવા.

જુક્યુ/ટેકનિકલ ઓફિસર.(સુખે)/એગ્રોમેટ/બુલેટીન/ /૨૦૧૮
તરઘડિયા(રાજકોટ), તારીખ:૦૪/૦૮/૨૦૧૮.

પ્રતિ,
તંત્રીશ્રી/ડાયરેક્ટરશ્રી,
રાજકોટ,

ઉપરોક્ત માહિતી આપના દૈનિક સમાચાર પત્રોમાં, રેડીયોમાં ગામના ચોરાના કાર્યક્રમમાં તથા ટીવીના ગ્રામ્ય કાર્યક્રમમાં વિના મૂલ્યે પ્રસિધ્ધ કરાવવા નમ્ર વિનતી છે. સહકારની અપેક્ષા સહ આભાર.

ટેકનીકલ ઓફિસર
સુકી ખેતીસંશોધન કેન્દ્ર
જુનાગઢ કૃષિ યુનિવર્સિટી
તરઘડીયા



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-45 for the Rajkot district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 05 to 09-September- 2018)

Significant past weather for the preceding week of Rajkot

Rainfall (mm)	20.1
Maximum temperature (°C)	27.4-29.9
Minimum temperature (°C)	22.2-22.8
Relative Humidity (%) I	88-91
Relative Humidity (%) II	79-90
Wind speed (Kmph)	4.6-6.4




Weather forecast until 08-30 hrs of 09/09/2018


(1) RAJKOT DISTRICT:



	05/09/18	06/09/18	07/09/18	08/09/18	09/09/18
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	29	30	30	30	29
Min.Temp.(°C)	23	23	23	23	22
Total CC (octa)	5	7	3	2	3
RH-I (%)	80	84	80	82	85
RH-II (%)	60	62	50	49	54
Wind Speed (kmph)	18	13	13	10	9
Wind Direction(deg.)	248	241	236	224	227

Agro meteorological Advisories:

General Advice	<ul style="list-style-type: none">• Rain fall: Humid and Partly cloudy to generally cloudy weather forecasted for next five days.• Temperature: Maximum and minimum temperature likely to be in range of 29-30°C and 22-23°C, respectively.• Relative Humidity: The RH-I and RH-II is likely to be in the range of 80-85% & 49-62, respectively.• Wind Direction and Speed: Direction and speed of wind is likely to be W, SW and 9-18 km/h, respectively.• Apply supplementary dose of urea fertilizer as per recommendation of respective
-----------------------	--

	<p>crop.</p> <ul style="list-style-type: none"> • Carry out interculturing and weeding as per need. 		
Crops	Stage	Insect/Pest/ Variety	Advisory
Kharif roundnut	Flowering/pegging/pod formation	<p>Stem rot</p> <p>Leaf spot-Tikka</p>  <p>White grub</p> 	<p>Apply 2.5 kg/ha Trichoderma fungus with well decomposed FYM OR castor cake @ 500 kg/ha as preventive measures against Sclerotium rolfsii stem rot.</p> <p>Due to continuous humid & cloudy weather spray Tebuconazole 10 ml/10 lit of to prevent leaf spot infestation.</p> <p>Do soil application of Phorate 10 G 25 kg/ha or Drenching of Chlorpyrifos or Quinalphos 4 liters/ha.</p>
Cotton	Square	<p>Jassid & Thrips</p> <p>Pink ball worm</p>	<p>Under present weather condition, severe attack of thrips persist in cotton since last two weeks, spray profenfos 20 ml or Ethion 20 ml + imidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 ml / 10 lit of water ml/10 lit of water.</p> <p>For the monitoring of pink bollworm infestation in cotton crop farmers are advised to put pheromone traps 6-8/ha in early sown crop. If infestation found pick and burry the infested flowers/bolls in soil for control of pink bollworm infestation in cotton. Also make spray of profenfos 50 EC 20 ml or cypermethrin 10 ml in 10 liters of water. Due to continuous humid weather spray Mancozeb 27 gram/10 lit of water at 15 days interval to prevent leaf spot infestation.</p>
Sesame	Flowering/capsule formation	Leaf webber	<p>Apply supplementary dose of nitrogen(Urea 54 kg/ha) at 30 DAS.</p> <p>Spray Cartep Hydrochloride 25 gram ml in 10 litres of water.</p> 

Fodder Sorghum & Maize	Heading	Sorghum-S-1049, Gundari, GFS-4 & 5 Maize-African tall	Monitor the crop for shoot fly infestation and if require spray cypermethrin 25 EC 5 ml in 10 liters of water.
Castor	Establishment/branching	GC-3,GCH-4,6,7	Procure good quality of seeds for kharif sowing. Carry out sowing if sufficient rain occurs.
Pigeon pea	Establishment	BDN-2,Vaishali,GJP-1	Procure good quality of seeds for kharif sowing. Carry out sowing if sufficient rain occurs.
Pulses	Flowering& pod formation/ ripening	-	Carry out interculturing and weeding. For the control stem fly infestation in Black gram spray DDVP 7 ml and fenvalerae 15 mlin 10 litres of water. Peak ripen pods of cow pea & green gram in time.
Vegetables	Nursery /Transplanting	Farmers are advised for preparation of nursery for tomato, chilli, brinjal and early cauliflower (maturing in the month of September). It is also advised to use shed nets, which helps in reducing vector borne diseases and produce healthy seedlings. Nursery should be covered by net at the height of 6.5 feet for protection against sunlight. Transplanting of brinjal, chilli and early cauliflower are advised for those farmers whose seedlings are ready.	
Lady finger	Flowering/fruiting	White fly, Shoot and fruit borer Mites	To control them spray neem seed extract 500 ml or neem oil 40 ml and Trizophos 40 % EC @ 25 ml / 10 lit of water at 15 days interval. Sparyneem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval(4 spray).If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear. Spraying of Ethion @ 1.5-2 ml/litre of water is advised.
Ridge gourd/ Bottle gourd/ Sponge gourd/ Bitter gourd/ Cucumbers	Flowering/Fruiting	White fly,thripsMite Downy mildew &Alternaria blight 	Sparyneem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 sprays) or acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval. To control mite infestation spray propargite 57 % EC 10/10 liter of water.
Cluster bean	Growing/fruiting	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval

Brinjal and tomato	Growing/fruiting	Sucking pest, Fruit and shoot borer	To control sucking pest spray neem seed extract 500 ml or neem oil 40 ml and Ethion 50 % EC 20 ml / 10 lit of water at 15 days interval. To control shoot and fruit borer, infested fruits and shoots should be collected and buried inside the soil. Installation of pheromone trap @ 2-3 traps per acre is advised for monitoring of borers. If pest population is high, spraying of Renaxypyr 18.5 EC @ 4 ml/10 litres of water is advised when sky remains clear.
Chili	Flowering & fruiting	Thrips → 	Under present weather conditions severe attack of thrips has been reported. Spray 20 ml profenofos in 10 litres of water.
Lemon	Fruit development	Psylla and thrips Citrus canker Leaf minor	To prevent damage of psylla & thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water. Spray copper oxychloride 40 gm or streptocyclie 1.5 gm in 10 litre of water Spray Dichlorvosh 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.
Livestock & poultry 	Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. Testing should be done. Do Brucellous calfwood vaccination. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Due to chance of rainfall keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid flies and mosquitoes. Do Arrangement for artificial light to birds to maintain egg production in poultry farm.		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-45 for the Jamnagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 05 to 09-September - 2018)




Weather forecast until 08-30 hrs of 09/09/2018


(2) JAMNAGAR DISTRICT:



	05/09/18	06/09/18	07/09/18	08/09/18	09/09/18
Rainfall (mm)	0	0	0	0	0
Max. Temp.(°C)	29	29	28	29	28
Min. Temp.(°C)	25	25	25	26	26
Total CC (octa)	6	7	3	3	4
RH-I (%)	77	79	76	77	79
RH-II (%)	67	68	57	59	62
Wind Speed (kmph)	25	20	19	13	11
Wind Direction(deg.)	249	242	243	224	214

Agro meteorological Advisories:-

General Advice	<ul style="list-style-type: none"> • Rain fall: Humid and Partly cloudy to generally cloudy weather forecasted for next five days. • Temperature: Maximum and minimum temperature likely to be in range of 28-29°C and 25-26°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 76-79% & 57-68, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be W, SW and 11-25 km/h, respectively. • Apply supplementary dose of urea fertilizer as per recommendation of respective crop. • Carry out interculturing and weeding as per need. 		
Crops	Stage	Insect/Pest/ Variety	Advisory
<i>Kharif roundnut</i>	Flowering/pegging/pod formation	Stem rot Leaf spot-Tikka	Apply 2.5 kg/ha Trichoderma fungus with well decomposed FYM OR castor cake @ 500 kg/ha as preventive measures against Sclerotium rolfsii stem rot. Due to continuous humid & cloudy weather spray

		 White grub 	<p>Tebuconazole 10 ml/10 lit of to prevent leaf spot infestation.</p> <p>Do soil application of Phorate 10 G 25 kg/ha or Drenching of Chlorpyrifos or Quinalphos 4 liters/ha.</p>
Cotton	Square	Jassid & Thrips Pink ball worm	<p>Under present weather condition, severe attack of thrips persist in cotton since last two weeks, spray profenfos 20 ml or Ethion 20 ml + imidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 mla / 10 lit of water ml/10 lit of water.</p> <p>For the monitoring of pink bollworm infestation in cotton crop farmers are advised to put pheromone traps 6-8/ha in early sown crop. If infestation found pick and burry the infested flowers/bolls in soil for control of pink bollworm infestation in cotton. Also make spray of profenfos 50 EC 20 ml or cypermethrin 10 ml in 10 liters of water. Due to continuous humid weather spray Mancozeb 27 gram/10 lit of water at 15 days interval to prevent leaf spot infestation.</p>
Sesame	Flowering/capsule formation	Leaf webber	<p>Apply supplementary dose of nitrogen(Urea 54 kg/ha) at 30 DAS.</p> <p>Spray Cartep Hydrochloride 25 gram ml in 10 litres of water.</p> 
Fodder Sorghum & Maize	Heading	Sorghum-S-1049, Gundari, GFS-4 & 5 Maize-African tall	Monitor the crop for shoot fly infestation and if require spray cypermethrin 25 EC 5 ml in 10 liters of water.
Castor	Establishment/branching	GC-3,GCH-4,6,7	Procure good quality of seeds for kharif sowing. Carry out sowing if sufficient rain occurs.
Pigeon pea	Establishment	BDN-2,Vaishali,GJP-1	Procure good quality of seeds for kharif sowing. Carry out sowing if sufficient rain occurs.
Pulses	Flowering&	-	Carry out interculturing and weeding.

	pod formation/ ripening		For the control stem fly infestation in Black gram spray DDVP 7 ml and fenvalerae 15 ml in 10 litres of water. Peak ripen pods of cow pea & green gram in time.
Vegetables	Nursery /Transplanting	Farmers are advised for preparation of nursery for tomato, chilli, brinjal and early cauliflower (maturing in the month of September). It is also advised to use shed nets, which helps in reducing vector borne diseases and produce healthy seedlings. Nursery should be covered by net at the height of 6.5 feet for protection against sunlight. Transplanting of brinjal, chilli and early cauliflower are advised for those farmers whose seedlings are ready.	
Lady finger	Flowering/fruit ing	White fly, Shoot and fruit borer Mites	To control them spray neem seed extract 500 ml or neem oil 40 ml and Trizophos 40 % EC @ 25 ml / 10 lit of water at 15 days interval. Spray neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 spray). If pest population is high, spraying of Spinosad 48 EC @ 1 ml / 4 litres of water is advised when sky remains clear. Spraying of Ethion @ 1.5-2 ml/litre of water is advised.
Ridge gourd/ Bottle gourd/ Sponge gourd/ Bitter gourd/ Cucumbers	Flowering/Fru iting	White fly, thrips, Mite Downy mildew & Alternaria blight 	Spray neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 sprays) or acetamiprid 20 % SP @ 5.0 gm / 10 lit of water at 15 days interval. Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosetyl 20 g/10 lit of water at 15 days interval. To control mite infestation spray propargite 57 % EC 10/10 liter of water.
Cluster bean	Growing/fru iting	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 ml or neem oil 40 ml and acetamiprid 20 % SP @ 5.0 gm / 10 lit of water at 15 days interval
Brinjal and tomato	Growing/fru iting	Sucking pest, Fruit and shoot borer	To control sucking pest spray neem seed extract 500 ml or neem oil 40 ml and Ethion 50 % EC 20 ml / 10 lit of water at 15 days interval. To control shoot and fruit borer, infested fruits and shoots should be collected and buried inside the soil. Installation of pheromone trap @ 2-3 traps per acre is advised for monitoring of borers. If pest population is high, spraying of Renaxypyr 18.5 EC @ 4 ml/10 litres of water is advised when sky remains clear.

Chili	Flowering & fruiting	Thrips → 	Under present weather conditions severe attack of thrips has been reported. Spray 20 ml profenofos in 10 litres of water.
Lemon	Fruit development	Psylla and thrips Citrus canker Leaf minor	To prevent damage of psylla&thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water. Spray copper oxychloride 40 gm or streptocyclie 1.5 gm in 10 litre of water Spray Dichlorvosh 10 ml or Quinalphos 20 ml in 10 litres of water for control of leafminor.
Livestock & poultry 	Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. Testing should be done. Do Brucellousiscalfwood vaccination. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Due to chance of rainfall keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid flies and mosquitoes. Do Arrangement for artificial light to birds to maintain egg production in poultry farm.		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-45 for the Amreli district

(Issued jointly by DFERS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 05 to 09-September - 2018)




Weather forecast until 08-30 hrs of 09/09/2018


(3) AMRELI DISTRICT:



	05/09/18	06/09/18	07/09/18	08/09/18	09/09/18
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	31	31	31	30	31
Min.Temp.(°C)	22	22	22	22	22
Total CC (octa)	5	6	4	2	3
RH-I (%)	84	87	83	82	85
RH-II (%)	65	65	61	55	55
Wind Speed (kmph)	14	12	12	9	7
Wind Direction(deg.)	253	240	250	247	249

Agro meteorological Advisories :

General Advice	<ul style="list-style-type: none"> • Rain fall: Humid and Partly cloudy to generally cloudy weather forecasted for next five days. • Temperature: Maximum and minimum temperature likely to be in range of 30-31°C and 22-22°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 82-87% & 55-65, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be W, SW and 7-14 km/h, respectively. • Apply supplementary dose of urea fertilizer as per recommendation of respective crop. • Carry out interculturing and weeding as per need. 		
Crops	Stage	Insect/Pest/ Variety	Advisory
<i>Kharif roundnut</i>	Flowering/pegging/pod formation	Stem rot	Apply 2.5 kg/ha Trichoderma fungus with well decomposed FYM OR castor cake @ 500 kg/ha as preventive measures against Sclerotium rolfsii stem rot.

		<p>Leaf spot-Tikka</p>  <p>White grub</p> 	<p>Due to continuous humid & cloudy weather spray Tebuconazole 10 ml/10 lit of to prevent leaf spot infestation.</p> <p>Do soil application of Phorate 10 G 25 kg/ha or Drenching of Chlorpyrifos or Quinalphos 4 liters/ha.</p>
Cotton	Square	<p>Jassid & Thrips</p> <p>Pink ball worm</p>	<p>Under present weather condition, severe attack of thrips persist in cotton since last two weeks, spray profenfos 20 ml or Ethion 20 ml + imidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 mla / 10 lit of water ml/10 lit of water.</p> <p>For the monitoring of pink bollworm infestation in cotton crop farmers are advised to put pheromone traps 6-8/ha in early sown crop. If infestation found pick and burry the infested flowers/bolls in soil for control of pink bollworm infestation in cotton. Also make spray of profenfos 50 EC 20 ml or cypermethrin 10 ml in 10 liters of water. Due to continuous humid weather spray Mancozeb 27 gram/10 lit of water at 15 days interval to prevent leaf spot infestation.</p>
Sesame	Flowering/capsule formation	Leaf webber	<p>Apply supplementary dose of nitrogen(Urea 54 kg/ha) at 30 DAS.</p> <p>Spray Cartep Hydrochloride 25 gram ml in 10 litres of water.</p> 
Fodder Sorghum & Maize	Heading	Sorghum-S-1049, Gundari, GFS-4 & 5 Maize-African tall	Monitor the crop for shoot fly infestation and if require spray cypermethrin 25 EC 5 ml in 10 liters of water.
Castor	Establishment/branching	GC-3,GCH-4,6,7	Procure good quality of seeds for kharif sowing. Carry out sowing if sufficient rain occurs.

Pigeon pea	Establishment	BDN-2, Vaishali, GJP-1	Procure good quality of seeds for kharif sowing. Carry out sowing if sufficient rain occurs.
Pulses	Flowering & pod formation/ ripening	-	Carry out interculturing and weeding. For the control stem fly infestation in Black gram spray DDVP 7 ml and fenvalerae 15 ml in 10 litres of water. Peak ripen pods of cow pea & green gram in time.
Vegetables	Nursery /Transplanting	Farmers are advised for preparation of nursery for tomato, chilli, brinjal and early cauliflower (maturing in the month of September). It is also advised to use shed nets, which helps in reducing vector borne diseases and produce healthy seedlings. Nursery should be covered by net at the height of 6.5 feet for protection against sunlight. Transplanting of brinjal, chilli and early cauliflower are advised for those farmers whose seedlings are ready.	
Lady finger	Flowering/fruiting	White fly, Shoot and fruit borer Mites	To control them spray neem seed extract 500 ml or neem oil 40 ml and Trizophos 40 % EC @ 25 ml / 10 lit of water at 15 days interval. Spray neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 spray). If pest population is high, spraying of Spinosad 48 EC @ 1 ml / 4 litres of water is advised when sky remains clear. Spraying of Ethion @ 1.5-2 ml/litre of water is advised.
Ridge gourd/ Bottle gourd/ Sponge gourd/ Bitter gourd/ Cucumbers	Flowering/Fruiting	White fly, thrips, Mite Downy mildew & Alternaria blight 	Spray neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 sprays) or acetamiprid 20 % SP @ 5.0 gm / 10 lit of water at 15 days interval. Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosetyl 20 g/10 lit of water at 15 days interval. To control mite infestation spray propargite 57 % EC 10/10 liter of water.
Cluster bean	Growing/fruiting	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 ml or neem oil 40 ml and acetamiprid 20 % SP @ 5.0 gm / 10 lit of water at 15 days interval
Brinjal and tomato	Growing/fruiting	Sucking pest, Fruit and shoot borer	To control sucking pest spray neem seed extract 500 ml or neem oil 40 ml and Ethion 50 % EC 20 ml / 10 lit of water at 15 days interval. To control shoot and fruit borer, infested fruits and shoots should be collected and buried inside the soil. Installation of pheromone trap @ 2-3 traps per acre is advised for monitoring of borers. If pest population is high, spraying of Renaxypyr 18.5 EC @ 4 ml/10 litres of

			water is advised when sky remains clear.
Chili	Flowering & fruiting	Thrips → 	Under present weather conditions severe attack of thrips has been reported. Spray 20 ml profenofos in 10 litres of water.
Lemon	Fruit development	Psylla and thrips Citrus canker Leaf minor	To prevent damage of psylla&thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water. Spray copper oxychloride 40 gm or streptocyclie 1.5 gm in 10 litre of water Spray Dichlorvosh 10 ml or Quinalphos 20 ml in 10 litres of water for control of leafminor.
Livestock & poultry 	Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. Testing should be done. Do Brucellousiscalfwood vaccination. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Due to chance of rainfall keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid flies and mosquitoes.Do Arrangement for artificial light to birds to maintain egg production in poultry farm.		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell:94277 28523



Agro met Advisory Service bulletin No.-45 for the Surendranagar district

(Issued jointly by DFRRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 05 to 09-September - 2018)




Weather forecast until 08-30 hrs of 09/09/2018


(4) SURENDRANAGAR DISTRICT:



	05/09/18	06/09/18	07/09/18	08/09/18	09/09/18
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	29	30	30	30	29
Min.Temp.(°C)	23	23	23	23	22
Total CC (octa)	6	7	2	1	2
RH-I (%)	79	79	77	76	79
RH-II (%)	56	55	44	44	45
Wind Speed (kmph)	19	16	14	12	12
Wind Direction(deg.)	246	241	236	239	247

Agro meteorological advisories:

General Advice	<ul style="list-style-type: none">• Rain fall: Humid and Partly cloudy to generally cloudy weather forecasted for next five days.• Temperature: Maximum and minimum temperature likely to be in range of 29-30°C and 22-23°C, respectively.• Relative Humidity: The RH-I and RH-II is likely to be in the range of 76-79% & 44-56, respectively.• Wind Direction and Speed: Direction and speed of wind is likely to be SW and 12-19 km/h, respectively.• Apply supplementary dose of urea fertilizer as per recommendation of respective crop.• In Surendranagar district Talukas under late onset of monsoon farmer are advised to sow Sesame (Purva-1), Castor(GC-3,GCH-4,6, 7), Sorghum (Gundhari, GFS-3, GAFS-11,CSV-21F, S-1049), Pigeon pea (BDN-2,Vaishali,GJP-1), Soybean (GS-1,3).		
Crops	Stage	Insect/Pest/ Variety	Advisory

<p>Kharif roundnut</p>	<p>Flowering/pegging/pod formation</p>	<p>Stem rot</p> <p>Leaf spot-Tikka</p>  <p>White grub</p> 	<p>Apply 2.5 kg/ha Trichoderma fungus with well decomposed FYM OR castor cake @ 500 kg/ha as preventive measures against Sclerotium rolfsii stem rot.</p> <p>Due to continuous humid & cloudy weather spray Tebuconazole 10 ml/10 lit of to prevent leaf spot infestation.</p> <p>Do soil application of Phorate 10 G 25 kg/ha or Drenching of Chlorpyrifos or Quinalphos 4 liters/ha.</p>
<p>Cotton</p>	<p>Square</p>	<p>Jassid & Thrips</p> <p>Pink ball worm</p>	<p>Under present weather condition, severe attack of thrips persist in cotton since last two weeks, spray profenfos 20 ml or Ethion 20 ml + imidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 ml / 10 lit of water ml/10 lit of water.</p> <p>For the monitoring of pink bollworm infestation in cotton crop farmers are advised to put pheromone traps 6-8/ha in early sown crop. If infestation found pick and bury the infested flowers/bolls in soil for control of pink bollworm infestation in cotton. Also make spray of profenfos 50 EC 20 ml or cypermethrin 10 ml in 10 liters of water. Due to continuous humid weather spray Mancozeb 27 gram/10 lit of water at 15 days interval to prevent leaf spot infestation.</p>
<p>Sesame</p>	<p>Flowering/capsule formation</p>	<p>Leaf webber</p>	<p>Apply supplementary dose of nitrogen(Urea 54 kg/ha) at 30 DAS.</p> <p>Spray Cartep Hydrochloride 25 gram ml in 10 litres of water.</p> 
<p>Fodder Sorghum & Maize</p>	<p>Heading</p>	<p>Sorghum-S-1049, Gundari, GFS-4 & 5 Maize-African tall</p>	<p>Monitor the crop for shoot fly infestation and if require spray cypermethrin 25 EC 5 ml in 10 liters of water.</p>

Castor	Establishment/ branching	GC-3,GCH-4,6,7	Procure good quality of seeds for kharif sowing. Carry out sowing if sufficient rain occurs.
Pigeon pea	Establishment	BDN- 2,Vaishali,GJP-1	Procure good quality of seeds for kharif sowing. Carry out sowing if sufficient rain occurs.
Pulses	Flowering& pod formation/ ripening	-	Carry out interculturing and weeding. For the control stem fly infestation in Black gram spray DDVP 7 ml and fenvalerae 15 mlin 10 litres of water. Peak ripen pods of cow pea & green gram in time.
Vegetables	Nursery /Transplanting	Farmers are advised for preparation of nursery for tomato, chilli, brinjal and early cauliflower (maturing in the month of September). It is also advised to use shed nets, which helps in reducing vector borne diseases and produce healthy seedlings. Nursery should be covered by net at the height of 6.5 feet for protection against sunlight. Transplanting of brinjal, chilli and early cauliflower are advised for those farmers whose seedlings are ready.	
Lady finger	Flowering/fruit ing	White fly, Shoot and fruit borer Mites	To control them spray neem seed extract 500 ml or neem oil 40 ml and Trizophos 40 % EC @ 25 ml / 10 lit of water at 15 days interval. Sparyneem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval(4 spray).If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear. Spraying of Ethion @ 1.5-2 ml/litre of water is advised.
Ridge gourd/ Bottle gourd/ Sponge gourd/ Bitter gourd/ Cucumbers	Flowering/Fruiti ng	White fly,thripsMite Downy mildew &Alternaria  blight	Sparyneem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 sprays) or acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval. To control mite infestation spray propargite 57 % EC 10/10 liter of water.
Cluster bean	Growing/fruiti ng	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval
Brinjal and tomato	Growing/fruiti ng	Sucking pest, Fruit and shoot borer	To control sucking pest spray neem seed extract 500 ml or neem oil 40 ml and Ethion 50 % EC 20 ml / 10 lit of water at 15 days interval. To control shoot and fruit borer, infested fruits and shoots should be collected and buried inside the soil.

			Installation of pheromone trap @ 2-3 traps per acre is advised for monitoring of borers. If pest population is high, spraying of Renaxypyr 18.5 EC @ 4 ml/10 litres of water is advised when sky remains clear.
Chili	Flowering & fruiting	Thrips → 	Under present weather conditions severe attack of thrips has been reported. Spray 20 ml profenofos in 10 litres of water.
Lemon	Fruit development	Psylla and thrips Citrus canker Leaf minor	To prevent damage of psylla & thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water. Spray copper oxychloride 40 gm or streptocyclie 1.5 gm in 10 litre of water Spray Dichlorvosh 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.
Livestock & poultry 	Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. Testing should be done. Do Brucellous calfwood vaccination. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Due to chance of rainfall keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid flies and mosquitoes. Do Arrangement for artificial light to birds to maintain egg production in poultry farm.		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.