



ગ્રામીણ કૃષિ મૌસમ સેવા  
હવામાન આગાહી આધારીત કૃષિસલાહ બુલેટીન  
ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ  
તા.૦૯-૦૪-૨૦૨૦ થી તા.૧૩-૦૫-૨૦૨૦



(ગ્રામીણ કૃષિ મૌસમ સેવા, સૂકી ખેતી સંશોધન કેન્દ્ર, જી.કે.યુ., તરઘડિયા અને ભારત મૌસમ વિભાગ દ્વારા સંપન્ન)

કૃષિ હવામાન બુલેટીન નં. ૧૨/૨૦૨૦

પાછલા અઠવાડિયાનું રાજકોટનું હવામાન

હવામાન પરિબલો	
વરસાદ(મી.મી)	0.0
મહત્તમ તાપમાન( સે.ગ્રે)	40.8-43.1
લઘુતમ તાપમાન( સે.ગ્રે)	24.1-25.5
ભેજ (%)સવાર	70-81
ભેજ (%)બપોરબાદ	18-27
પવનની ગતિ( કિમી/કલાક)	8.5-11.6

હવામાન આગાહી :પત્રક-ક

ઉત્તરસૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ-હવામાન આગાહી

તા.૦૯-૦૫-૨૦૨૦ થી ૧૩-૦૫-૨૦૨૦

કૃષિ સલાહ :

સામાન્ય	<p>➤ ભારત સરકારશ્રીના હવામાન ખાતા તરફથી મળેલ હવામાન માહિતી અનુસાર અત્રેના ગ્રામીણકૃષિ મૌસમ સેવા વિભાગ, જુનાગઢ કૃષિ યુનિવર્સિટી, તરઘડિયા તરફથી જણાવવામાં આવે છે કે, ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના</p> <p>રાજકોટ :જીલ્લામાં આગામી તા.૦૯-૦૫-૨૦૨૦ થી તા.૧૩-૦૫-૨૦૨૦ દરમિયાન હવામાન સૂકું, ગરમ અને યોખ્ખું રહેશે. પવનની દિશા પશ્ચિમની રહેવાની અને પવનની ઝડપ ૧૦ થી ૧૩ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૪૩-૪૪ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમિયાન ૨૪-૨૫ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.</p>					
તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫	
પેરામીટર	૦૯-૦૫-૨૦	૧૦-૦૫-૨૦	૧૧-૦૫-૨૦	૧૨-૦૫-૨૦	૧૩-૦૫-૨૦	
વરસાદ(મી.મી.)	0	0	0	0	0	
મહત્તમતાપમાન( સે.ગ્રે)	44	43	43	43	43	
લઘુતમતાપમાન( સે.ગ્રે)	25	25	24	25	25	
વાદળનીસ્થિતિ( ઓકટા)	0	0	0	0	0	
મહત્તમભેજ (%)સવાર	82	80	83	84	84	
લઘુતમભેજ (%) બપોરબાદ	30	28	26	25	26	

પવનનીગતિ(કિમી/કલાક)	13	10	12	12	12
પવનનીદિશા(ડીગ્રી)	260	268	275	270	269

**મોરબી :** જીલ્લામાં આગામી તા.૦૯-૦૫-૨૦૨૦ થી તા.૧૩-૦૫-૨૦૨૦ દરમ્યાન હવામાન સુકું, ગરમ અને યોખ્ખું રહેશે. પવનની દિશા પશ્ચિમ, વાયવ્ય અને નૈઋત્યની રહેવાની અને પવનની ઝડપ ૬ થી ૧૨ કીમી/કલાક રહેવાની શક્યતા છે . આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૪૩-૪૪ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૨૪-૨૫ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૯-૦૫-૨૦	૧૦-૦૫-૨૦	૧૧-૦૫-૨૦	૧૨-૦૫-૨૦	૧૩-૦૫-૨૦
વરસાદ(મી.મી.)	0	0	0	0	0
મહત્તમતાપમાન( સે.ગ્રે)	44	43	43	43	43
લઘુત્તમતાપમાન( સે.ગ્રે)	25	25	24	25	25
વાદળનીસ્થિતિ( ઓકટા)	0	0	0	0	0
મહત્તમભેજ (%)સવાર	46	39	38	39	43
લઘુત્તમભેજ (%) બપોરબાદ	10	11	12	11	11
પવનનીગતિ(કિમી/કલાક)	8	10	6	8	12
પવનનીદિશા(ડીગ્રી)	272	335	237	257	290

**જામનગર:** જીલ્લામાં આગામી તા.૦૯-૦૫-૨૦૨૦ થી તા.૧૩-૦૫-૨૦૨૦ દરમ્યાન હવામાન સુકું, ગરમ અને યોખ્ખું રહેશે. પવનની દિશા પશ્ચિમની રહેવાની અને પવનની ઝડપ ૧૧ થી ૧૩ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૩૮-૩૯ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૨૫-૨૬ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૯-૦૫-૨૦	૧૦-૦૫-૨૦	૧૧-૦૫-૨૦	૧૨-૦૫-૨૦	૧૩-૦૫-૨૦
વરસાદ(મી.મી.)	0	0	0	0	0
મહત્તમતાપમાન( સે.ગ્રે)	38	38	39	38	38
લઘુત્તમતાપમાન( સે.ગ્રે)	26	26	26	25	25
વાદળનીસ્થિતિ( ઓકટા)	0	0	0	0	0
મહત્તમભેજ (%)સવાર	88	88	88	88	89
લઘુત્તમભેજ (%) બપોરબાદ	53	51	49	49	51
પવનનીગતિ(કિમી/કલાક)	13	11	11	12	13
પવનનીદિશા(ડીગ્રી)	255	256	273	266	270

**દેવભૂમિ દ્વારકા :** જીલ્લામાં આગામી તા.૦૯-૦૫-૨૦૧૯ થી તા.૧૩-૦૫-૨૦૨૦ દરમ્યાન હવામાન સુકું, ગરમ રહેશે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૩૧-૩૩ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૨૬-૨૭ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ પેરામીટર	દિવસ-૧ ૦૯-૦૫-૨૦	દિવસ-૨ ૧૦-૦૫-૨૦	દિવસ-૩ ૧૧-૦૫-૨૦	દિવસ-૪ ૧૨-૦૫-૨૦	દિવસ-૫ ૧૩-૦૫-૨૦
વરસાદ(મી.મી.)	0	0	0	0	0
મહત્તમતાપમાન( સે.ગ્રે)	32	33	32	32	31
લઘુતમતાપમાન( સે.ગ્રે)	27	27	26	26	27
વાદળનીસ્થિતિ( ઓકટા)					
મહત્તમભેજ (%)સવાર					
લઘુતમભેજ (%) બપોરબાદ					
પવનનીગતિ(કિમી/કલાક)					
પવનનીદિશા(ડીગ્રી)					

**અમરેલી :** જીલ્લામાં આગામી તા.૦૯-૦૫-૨૦૨૦ થી તા.૧૩-૦૫-૨૦૨૦ દરમ્યાન હવામાન સુકું, ગરમ અને યોખ્ખું રહેશે. પવનની દિશા પશ્ચિમ અને વાયવ્યની રહેવાની અને પવનની ઝડપ ૭ થી ૧૦ કીમી/કલાક રહેની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૪૩-૪૪ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમ્યાન ૨૪-૨૫ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ પેરામીટર	દિવસ-૧ ૦૯-૦૫-૨૦	દિવસ-૨ ૧૦-૦૫-૨૦	દિવસ-૩ ૧૧-૦૫-૨૦	દિવસ-૪ ૧૨-૦૫-૨૦	દિવસ-૫ ૧૩-૦૫-૨૦
વરસાદ(મી.મી.)	0	0	0	0	0
મહત્તમતાપમાન( સે.ગ્રે)	44	43	43	43	43
લઘુતમતાપમાન( સે.ગ્રે)	25	25	24	25	25
વાદળનીસ્થિતિ( ઓકટા)	0	0	0	0	0
મહત્તમભેજ (%)સવાર	78	74	74	75	76
લઘુતમભેજ (%) બપોરબાદ	34	33	31	32	30
પવનનીગતિ(કિમી/કલાક)	10	9	9	7	8
પવનનીદિશા(ડીગ્રી)	288	300	303	272	300


**સુરેન્દ્રનગર :** જીલ્લામાં આગામી તા.૦૯-૦૫-૨૦૨૦ થી તા.૧૩-૦૫-૨૦૨૦ દરમ્યાન હવામાન સુકું, ગરમ અને યોખ્ખું રહેશે. પવનની દિશા પશ્ચિમની રહેવાની અને પવનની ઝડપ ૧૩ થી ૧૫ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૪૩-૪૪ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમ્યાન ૨૬-૨૭ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.


તારીખ પેરામીટર	દિવસ-૧ ૦૯-૦૫-૨૦	દિવસ-૨ ૧૦-૦૫-૨૦	દિવસ-૩ ૧૧-૦૫-૨૦	દિવસ-૪ ૧૨-૦૫-૨૦	દિવસ-૫ ૧૩-૦૫-૨૦
વરસાદ(મી.મી.)	0	0	0	0	0
મહત્તમતાપમાન( સે.ગ્રે)	44	43	43	43	43
લઘુતમતાપમાન( સે.ગ્રે)	27	27	26	27	27
વાદળનીસ્થિતિ( ઓકટા)	0	0	0	1	0


મહત્તમભેજ (%)સવાર	74	72	74	77	77
લઘુત્તમભેજ (%) બપોરબાદ	27	25	25	24	24
પવનનીગતિ(કિમી/કલાક)	15	13	14	14	14
પવનનીદિશા(ડીગ્રી)	263	273	278	279	268

- ૦૯ થી ૧૩-મે -૨૦૨૦ દરમ્યાન હવામાન સુકું, ગરમ અને ચોખ્ખું રહેશે. પવનની દિશા પશ્ચિમની મોટાભાગે રહેવાની અને પવનની ઝડપ ૬ થી ૧૫ કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૩૧-૪૪ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૨૪-૨૭ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.
- કોરોના વાયરસને ફેલાતો અટકાવવા બીનજરૂરી અવર જવર ટાળવી, કોઈપણ સંજોગોમાં સમુહમાં ભેગા થવું નહિ, ટોળે વળીને બેસવું નહિ.
- વારંવાર સાબુથી હાથ ધોવા અને પછી જ આંખ, કાન અને મોને અડકવું.
- મોં પર કપડું (માસ્ક) બાંધી રાખવું, છોંક અને ઉઘરસ વખતે મો અને નાક આડે રુમાલ અવશ્ય રાખવો.
- કાપણી, શ્રેશિંગ અને અન્ય ખેતી વિષયક કામગીરી દરમ્યાન સામાજીક અંતર (બે વ્યક્તિ વચ્ચે ૬ થી ૭ ફૂટ અંતર) જાળવવું તેમજ ભોજન અને આરામના સમયે અલગ અલગ દુર બેસવું.
- ખેત ઓજારોની આપ-લે દરમ્યાન ઓજારોને સાબુના પાણીથી સાફ કર્યા બાદ જ એક બીજાને આપવા અને ઉપયોગમાં લેવા.
- શરદી, ઉઘરસ, કફ અને તાવ જેવું લાગે તો નજીકના આરોગ્ય કેન્દ્રનો સંપર્ક સાધવો.
- ખેત પેદાશના વેચાણ અને બીજ, ખાતર અને દવાઓની ખરીદી દરમ્યાન મોં પર કપડું (માસ્ક) બાંધી રાખવું અને બે વ્યક્તિ વચ્ચે ૬ થી ૭ ફૂટ અંતર જાળવવું.
- આગામી તા. ૧૪ થી ૧૭ મે દરમ્યાન હવામાન ચોખ્ખું, સુકું અને ગરમ લુ પડવાની શક્યતા રહેશે.
- આ હવામાન પરિસ્થિતિમાં ઉનાળુ શાકભાજી, તલ, અને ઘાસચારાના પાકોને સવારે અથવા સાંજના સમયે જરૂરીયાત મુજબ હળવું પિયત આપવું.
- આકરા તાપમાં બપોરના સમયે બહાર નીકળવું નહિ, પુષ્કળ પાણી પીવું, હલકા સુતરાઉ કપડા પહેરવા અને માથા પર ટોપી પહેરવી. તેમજ પશુઓને બપોરના સમયે વૃક્ષોના છાયામાં અથવા શેડમાં રાખવા.

મુખ્ય પાક	પાક અવસ્થા	રોગ/જીવાત/જાત	કૃષિ સલાહ (૧૦ લીટરપાણીમાં મિશ્ર કરી ભલામણ કરેલ દવા છાંટવી )
ખાતરની ખાડ	પાકું ખાતર પલટાવવું	-	પશુઓના મળમૂત્ર અને ખેત કચરામાંથી પૂર્ણ સદી ગયેલ ખાતરને ખાકાંથી બહાર કાઢવું. 

<p><b>ઉનાળુ મગફળી</b></p>	<p>ડોડવા બંધાવવા અને દાણા ભરાવા</p>	<p><b>કથીરી</b></p>	<ul style="list-style-type: none"> <li>• ડોડવા બન્યાવાની અવસ્થાએ મગફળીને હળવું પિયત આપવું.</li> <li>• ગરમ હવામાનને કારને મગફળીમાં લાલ કથીરીનો ઉપદ્રવ થવાની શક્યતા હોય સતત નિરિક્ષણ કરતા રહેવું. જો ઉપદ્રવ જણાય તો ઇથીઓન દવા ૨૦ મી.લી. અથવા પ્રોપરગાઈટ ૫૭ ઇ.સી. ૧૦ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.</li> </ul> 
<p><b>ઉનાળુ તલ</b></p>	<p>બેઢાંમાં દાણા ભરાવવા અને દેહધાર્મિક પરિપક્વ અવસ્થા</p>	<p>-</p>	<p>➤ પરિપક્વ અવસ્થાએ સમયસર કાપણી કરવી.</p>
<p><b>ઉનાળુ મગ અને અડદ</b></p>	<p>ફૂલ અને શીંગ બંધાવવી</p>	<p>લીલી ઇયળ</p>	<p>ઉનાળુ મગ અને અડદમાં જો ઇયળોનો વધુ ઉપદ્રવ હોય તો ઇન્ડોકસાકાર્બ ૧૦ મીલી/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p>
<p><b>સરગવો</b></p>	<p>ફૂલ/શીંગ</p>	<p>ફૂલ અને પાન ખાનાર ઇયળ</p>	<p>સરગવામાં ફૂલ અને પાન ખાનાર ઇયળના નિયંત્રણ માટે લીબોળીનું તેલ એઝાદીરેક્ટીન ૧૫૦૦ પીપીએમ ૫૦ મીલી અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી/ ૧૦ લી.પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p> <p>તેમજ પુખ્ત કીટકના નિયંત્રણ માટે પ્રતિ હેક્ટર પ્રકાશ પિંજર મુકવા.</p> <p>વધુ ઉપદ્રવ હોય તો મેલાથીયોન ૫૦ ઇસી ૨૦ મિલી/ ૧૦ લી.પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p>
<p><b>ભીંડા</b></p>	<p>ફૂલ/ફળ</p>	<p>પાન કથીરી</p>	<p>ગરમ હવામાનના કારણે ભીંડામાં જો પાન કથીરીનો ઉપદ્રવ હોય તો ઇથીઓન ૩૦ મિલી પ્રતિ પંપ છંટકાવ કરવો.</p> <p>ગરમ હવામાનને કારણે ભીંડામાં સવાર અથવા સાંજના સમયે ટૂંકા અંતરે ફાળવું પિયત આપવું.</p> <p>ભીંડામાં શીન્ગોની વીણી કર્યા બાદ ૧૫-૨૦ કી.ગ્રા. યુરિયા ખાતર પ્રતિ હેક્ટર આપવું.</p>

<p>તુરીયા, દૂધી, ગલકા, કારેલા, ગુવાર, કાકડી, તરબુચ</p>	<p>ફૂલ/ફળ</p>	<p>શ્રીપ્સ/ સફેદ માખી  કથીરી  તળછારો</p>	<p>ઉનાળુ શાકભાજીમાં સફેદમાખીના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦ મિલિ અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને ઈમીડાક્લોપ્રીડ ૩૦.૫ ટકા ૪ મિલી અથવા એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. જો પાન કથીરીનો ઉપદ્રવ હોય તો ઇથીઓન ૩૦ મિલી પ્રતિ પંપ છાન્કાવ કરવો. વેલાવાળા શાકભાજીમાં તળછારાના નિયંત્રણ માટે મેન્કોઝેબ ૪૦ ગ્રામ પ્રતિ પંપમાં મિશ્ર કરી છંટકાવ કરવો. વેલાવાળા શાકભાજીમાં પરાગનયનની પ્રક્રિયા સારી થાય તે માટે હળવા અને ટુંકા ગાળાના પિયતથી જમીનમાં ભેજ જાળવવો જેથી વધુ ઉત્પાદન મેળવી શકાય.</p>
<p>રીંગણી</p>	<p>ફૂલ/ફળ</p>	<p>સફેદ માખી  ડુંબ અને → ફળ કોરી ખાનાર ઈયળ  પાન કથીરી</p>	<p>➤ સફેદ માખીના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦ મિલિ અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને ડાયફેન્થાયુરોન ૫૦ ટકા વે.પા. ૧૬ ગ્રામ અથવા ટ્રાયઝોફોસ ૪૦ ઈસી ૨૫ મિ.લી. દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. ➤ ફળ અને ડોકાની ઈયળોનો ઉપદ્રવ હોય તો નુકશાન પામેલ ફળ અને ડોકા તોડીને જમીનમાં દાંટી દેવા અને ક્લોરાનટ્રાનીલીપ્રોલ ૧૮.૫ એસ.સી (રીનાક્ષીપાયર) દવા ૩ મિલી અથવા ડીડીવીપી ૭૬ ઈસી ૫ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. ➤ રીંગણામાં જો પાનકથીરીનો ઉપદ્રવ હોય તો પ્રોપરગાઈટ ૫૭ ઈ.સી. ૧૫ મિલી પ્રતિ પંપ છંટકાવ કરવો.</p>
<p>મરચી</p>	<p>વૃદ્ધિ/ફૂલ/ફળ</p>	<p>શ્રીપ્સ</p>	<p>➤ માર્ચમાં શ્રીપ્સના નિયંત્રણ માટે પ્રોફેનોફોસ ૨૦ મીલી અને લીબોળીનું તેલ ૫૦ મિલી અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મિલી ૧૦ લી. પાણીમાં મિશ્ર કરી છંટકાવ કરવો. શ્રીપ્સનો વધુ ઉપદ્રવ હોય તો સ્પીનોસાદ અથવા સ્પીનેટોરમ દવા ૫ મિલી. પ્રતિ પંપ છાંટવી.</p>
<p>લીબુ</p>	<p>ફળનો વિકાસ</p>	<p>પાન કથીરી </p>	<p>ઊંચા તાપમાનને કારણે લીબુના બગીયામાં પાન કથીરીનો ઉપદ્રવ વધુ હોય તો તેના નિયંત્રણ માટે પ્રોપરગાઈટ દવા ૧૦ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. લીબુના બગીયાને નિયત અંતરે સમયસર પિયત આપવું જેથી ફળ ફાટે નહિ.</p>
<p>આંબો</p>	<p>ફળનો વિકાસ</p>	<p>ફળમાખી</p>	<p>કેરી લખોટી જેવડા કદની થાય ત્યારે ૨ ગ્રામ નેપ્થેલીન એસેટીક એસીડ અને ૨ કી.ગ્રા. યુરીયાને ૧૦૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો જેથી કેરીનું ખરણ ઓછું થાય. આંબામાં ફળમાખીના નિયંત્રણ માટે ફેન્થીઓન ૧૦ મિલી અથવા ડીડીવીપી ૪ મિલી અને મીથાયલયુજીનોલ ૧૦ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p>

<p>પશુ/ મરઘા</p> 	<p>તાપમાન વધારે રહેવાના કારણે ઢોરને દિવસના સમયે છાયડામાં રાખવા અને મિનરલ મિક્શર યુક્ત પીવાના સ્વચ્છ પાણીની વ્યવસ્થા કરાવી. દુધાળા પશુઓને ઉનાળામાં ઠંડક આપવા પાણીનો છંટકાવ કરવો. ખોરાકમાં લીલા ચારનું પ્રમાણ વધારવું. દુધાળા પશુઓના આઉને ઝીંક ઓક્સાઇડ અથવા બોરિક પાવડરથી બરાબર સાફ કરવું. પશુઓમાં ગળસુંઢો અને ગાંઠીયા તાવના રોગના નિયંત્રણ માટે રસીકરણ કરાવવું. ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટ્રાઝ ૨ મી.લી. ૧ લીટર પાણીમાં નાખીને છાંટવી. વિટામીન એ તથા ડી પુરકો આપવા. કરમિયાની દવા આપવી.</p>
--	---

જુફ્યુ/ટેકનિકલ ઓફિસર).સુખે/(એગ્રોમેટ/બુલેટીન / / ૨૦૨૦  
તરઘડિયા(રાજકોટ), તારીખ:૦૮/૦૫/૨૦૨૦.

પ્રતિ,  
તંત્રીશ્રી/ડાયરેક્ટરશ્રી,  
રાજકોટ,

ઉપરોક્ત માહિતી આપના દૈનિક સમાચાર પત્રોમાં, રેડીયોમાં ગામના ચોરાના કાર્યક્રમમાં તથા ટીવીના ગ્રામ્ય કાર્યક્રમમાં વિના મૂલ્યે પ્રસિધ્ધ કરાવવા નમ્ર વિનતી છે. સહકારની અપેક્ષા સહ આભાર.

ટેક્નીકલ ઓફિસર  
સુકી ખેતીસંશોધન કેન્દ્ર  
જુનાગઢ કૃષિ નિવર્સિટી  
તરઘડીયા



**Gramin Krushi Mausam Seva**  
**Main Dry Farming Research Station**  
**Junagadh Agricultural University**  
**Targhadia (Rajkot) - 360 003**  
**Phone: (0281)2784260(O), 2331686(R)**  
**Fax : (0281)2784722, Cell :94269 38235**



**Agromet Advisory Service bulletin No.12 -for the Rajkot district**

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 09 to 13-May - 2020)

**Significant past weather for the preceding week of Rajkot**

Rainfall (mm)	0.0
Maximum temperature (°C)	40.8-43.1
Minimum temperature (°C)	24.1-25.5
Relative Humidity (%) I	70-81
Relative Humidity (%) II	18-27
Wind speed (Kmph)	8.5-11.6

**Weather forecast until 08-30 hrs. of 13/05/2020**




**(1) RAJKOT DISTRICT:**


	09/05/20	10/05/20	11/05/20	12/05/20	13/05/20
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	44	43	43	43	43
Min.Temp.(°C)	25	25	24	25	25
Total CC (octa)	0	0	0	0	0
RH-I (%)	82	80	83	84	84
RH-II (%)	30	28	26	25	26
Wind Speed (kmph)	13	10	12	12	12
Wind Direction(deg.)	260	268	275	270	269




**Agro meteorological Advisories:**

<b>General Advice</b>	<ul style="list-style-type: none"><li>• <b>Rain fall: Dry, hot and clear weather forecasted next five days</b></li><li>• <b>Temperature:</b> Maximum and minimum temperature likely to be in range of 43-44°C and 24-25°C, respectively.</li><li>• <b>Relative Humidity:</b> The RH-I and RH-II is likely to be in the range of 80-84% &amp; 25-30, respectively.</li><li>• <b>Wind Direction and Speed:</b> Direction and speed of wind is likely to be W and 10-13 km/h, respectively.</li><li>• <b>Keeping in view of threat of COVID-19 spread farmers are advised to follow the government guidelines, personal hygiene, use mask, ensure hand washing with soap at reasonable interval, social distancing of 6 feet during harvesting and other</b></li></ul>
-----------------------	---



	<p><b>agricultural operation in all crops.</b></p> <ul style="list-style-type: none"> <li>• <b>Keep social distancing of 5-6 feet in selling of agricultural produces in Marketing yard and also during buying of agricultural inputs.</b></li> <li>• <b>Weather outlook for subsequent week: Dry, hot and clear weather very likely to observe during 14-17 May-2020.</b></li> <li>• <b>Due to prevailing dry and hot weather conditions, farmers are advised for light irrigation in all vegetable crops. Irrigation should be done in the morning hours or in the evening hours.</b></li> <li>• <b>Under hot weather a. Avoid heat exposure. b. Wear light weight, light coloured, loose cotton clothes. c. Cover your head. d. Keep the animal under tree shade or shed during noon hours.</b></li> <li>• <b>Deep ploughing of fields after harvesting of rabi crops is advised so that eggs and pupating stages of insects, fungus, bacteria and weed seeds are destroyed.</b></li> </ul>		
<b>Crops</b>	<b>Stage</b>	<b>Insect/Pest/ Variety</b>	<b>Advisory</b>
<b>Farm compost/ FYM</b>	Turning of well rotted compost & FYM	-	<p>Dig out completely composted material made from animal dung, urine, litter and farm waste.</p> 
<b>Summer Groundnut</b>	Pod formation/development  	<b>Red Mite</b>  	<p><b>Due to hot weather monitor the crop against the attack of mite in groundnut is advised. If population is found more then spraying of Ethion @ 1.5-2 ml or Propargite 57 % EC @ 10 ml /litre of water is advised.</b></p> <p>Apply light irrigation at pod formation and development stages.</p>
<b>Summer Sesame</b>	Physiological Maturity	-	Harvest in time at ripening stage before seed shattering.
<b>Green gram &amp; Black gram</b>	Flowering & pod formation	Heliiothis	<p>Under present weather condition infestation of heliiothis and whiteflies were observed in summer sown green gram &amp; black gram. To control heliiothis spray spinosad 3 ml/10 lit of water with 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml / 10 lit of water.</p>

<b>Drum stick/ Moringa</b>	Flowering & fruiting	Bud worm & leaf caterpillar	<p><b>Monitor the crop for infestation of Bud worm &amp; leaf caterpillar, if require spray 500 ml of 5 % neem seed extract (Azadirachtin) or neem seed oil (Azadirachtin 1500 ppm) 50 ml with / 10 lit of water.</b></p> <p><b>Use light trap 1-2/ha to attract &amp; kill the adult.</b></p> <p><b>Under heavy infestation make spray of Malathion 50 EC 20 ml/10 litre of water.</b></p>
<b>Lady finger</b>	Flowering/fruiting	Mites	<p>Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised.</p> <p>Due to prevailing high temperature, light irrigation at short interval is advised.</p> <p>After harvesting of mature okra, urea should be applied @ 15-20 kg/ha.</p>
<b>Ridge gourd, Bottle gourd, Sponge gourd, Bitter gourd, Cluster bean, Water melon</b>	Flowering & fruiting	<p>White flies/</p> <p>Mites</p> <p>Downy mildew</p>	<p>In all vegetable crops whitefly infestation observed above ETL. Spray 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml with acetamiprid 20 % SP@ 5.0 gm or <b>imidacloprid 30.5 SC</b> 3 ml/10 lit of water for whiteflies control.</p> <p>If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites in cucurbitaceous .</p> <p>For the control of downy mildew spray mencozeb 40g/pump.</p> <p>Optimum moisture level should be maintained in cucurbitaceous crops by light and frequent irrigation to increase pollination and thus increase yield of the crop.</p>
<b>Brinjal</b>	Flowering/fruiting	<p><b>Sucking pest</b> →</p> <p><b>Shoot &amp; Fruit borer</b></p>  <p>Mites</p>	<p><b>Monitor the crop for infestation of white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised. To control shoot and fruit borer in brinjal and tomato crops, infested fruits and shoots should be collected and buried inside the soil. If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</b></p> <p><b>If infestation of mites found above ETL in brinjal, then make spraying of Propargite 57 % EC @ 10 ml/ 10 litre of water for the control of Mites under clear sky.</b></p>

<b>Chili</b>	Flowering & fruiting	<b>Thrips</b> → 	<p>Under present weather conditions severe attack of thrips has been reported. Spray profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of <b>spinosad or spinetoram</b> 3 ml per 10 liter of water.</p>
<b>Mango</b>	Fruit development	Fruit fly	<p>Spray mixture of DDVP 4 ml and Methyleugenol 10 ml in 10 litres of water for control of fruit fly. Apply irrigation as per need.</p>
<b>Lemon</b>	Fruit development	<b>Mite</b> 	<p>Due to high temperature mites infestation in lemon orchard observed. To control mite infestation spray propargite 57 % EC 10/10 liter of water.</p> <p>To prevent cracking of fruits apply irrigation in time at regular interval.</p>
<b>Livestock &amp; poultry</b> 	<p>Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day. Do Vaccination for <b>Haemorrhagic septicemia</b> (H.S.) disease and <b>Blue Quarter</b> (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.</p>		

**Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.**



**Gramin Krushi Mausam Seva**  
**Main Dry Farming Research Station**  
**Junagadh Agricultural University**  
**Targhadia (Rajkot) - 360 003**  
**Phone: (0281)2784260(O), 2331686(R)**  
**Fax : ( 0281)2784722, Cell :9426938235**



**Agro met Advisory Service bulletin No.-12 for the Morbi district**

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 09 to 13-May - 2020)




**Weather forecast until 08-30 hrs. of 13/05/2020**



**(2) MORBI DISTRICT:**




	09/05/20	10/05/20	11/05/20	12/05/20	13/05/20
Rainfall (mm)	0	0	0	0	0
Max. Temp.(°C)	44	43	43	43	43
Min. Temp.(°C)	25	25	24	25	25
Total CC (octa)	0	0	0	0	0
RH-I (%)	46	39	38	39	43
RH-II (%)	10	11	12	11	11
Wind Speed (kmph)	8	10	6	8	12
Wind Direction(deg.)	272	335	237	257	290

**Agro meteorological Advisories:-**

<b>General Advice</b>	<ul style="list-style-type: none"><li>• <b>Rain fall: Dry, hot and clear weather forecasted next five days</b></li><li>• <b>Temperature:</b> Maximum and minimum temperature likely to be in range of 43-44°C and 24-25°C, respectively.</li><li>• <b>Relative Humidity:</b> The RH-I and RH-II is likely to be in the range of 38-46% &amp; 10-12, respectively.</li><li>• <b>Wind Direction and Speed:</b> Direction and speed of wind is likely to be W, NW, SW and 6-12 km/h, respectively.</li><li>• <b>Keeping in view of threat of COVID-19 spread farmers are advised to follow the government guidelines, personal hygiene, use mask, ensure hand washing with soap at reasonable interval, social distancing of 6 feet during harvesting and other agricultural operation in all crops.</b></li><li>• <b>Keep social distancing of 5-6 feet in selling of agricultural produces in Marketing yard and also during buying of agricultural inputs.</b></li><li>• <b>Weather outlook for subsequent week: Dry, hot and clear weather very likely to observe during 14-17 May-2020.</b></li><li>• <b>Due to prevailing dry and hot weather conditions, farmers are advised for light irrigation in all vegetable crops. Irrigation should be done in the morning hours</b></li></ul>
-----------------------	--

	<p><b>or in the evening hours.</b></p> <ul style="list-style-type: none"> <li>• Under hot weather a. Avoid heat exposure. b. Wear light weight, light coloured, loose cotton clothes. c. Cover your head. d. Keep the animal under tree shade or shed during noon hours.</li> <li>• Deep ploughing of fields after harvesting of rabi crops is advised so that eggs and pupating stages of insects, fungus, bacteria and weed seeds are destroyed.</li> </ul>		
Crops	Stage	Insect/Pest/ Variety	Advisory
<b>Farm compost/ FYM</b>	Turning of well rotted compost & FYM	-	<p>Dig out completely composted material made from animal dung, urine, litter and farm waste.</p> 
<b>Summer Groundnut</b>	Pod formation/development  	<b>Red Mite</b>  	<p>Due to hot weather monitor the crop against the attack of mite in groundnut is advised. If population is found more then spraying of Ethion @ 1.5-2 ml <b>or</b> Propargite 57 % EC @ 10 ml /litre of water is advised.</p> <p>Apply light irrigation at pod formation and development stages.</p>
<b>Summer Sesame</b>	Physiological Maturity	-	Harvest in time at ripening stage before seed shattering.
<b>Green gram &amp; Black gram</b>	Flowering & pod formation	Heliothis	Under present weather condition infestation of heliothis and whiteflies were observed in summer sown green gram & black gram. To control heliothis spray spinosad 3 ml/10 lit of water with 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml / 10 lit of water.
<b>Drum stick/ Moringa</b>	Flowering & fruiting	Bud worm & leaf caterpillar	<p><b>Monitor the crop for infestation of Bud worm &amp; leaf caterpillar, if require spray 500 ml of 5 % neem seed extract (Azadirachtin) or neem seed oil (Azadirachtin 1500 ppm) 50 ml with / 10 lit of water.</b></p> <p><b>Use light trap 1-2/ha to attract &amp; kill the adult.</b></p> <p><b>Under heavy infestation make spray of Malathion 50 EC 20 ml/10 litre of water.</b></p>

<b>Lady finger</b>	Flowering/fruiting	Mites	<p>Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised.</p> <p>Due to prevailing high temperature, light irrigation at short interval is advised.</p> <p>After harvesting of mature okra, urea should be applied @ 15-20 kg/ha.</p>
<b>Ridge gourd, Bottle gourd, Sponge guard, Bitter gourd, Cluster bean, Water melon</b>	Flowering & fruiting	<p>White flies/</p> <p>Mites</p> <p>Downy mildew</p>	<p>In all vegetable crops whitefly infestation observed above ETL. Spray 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml with acetamiprid 20 % SP@ 5.0 gm or <b>imidacloprid 30.5 SC</b> 3 ml/10 lit of water for whiteflies control.</p> <p>If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites in cucurbitaceous .</p> <p>For the control of downy mildew spray mencozeb 40g/pump.</p> <p>Optimum moisture level should be maintained in cucurbitaceous crops by light and frequent irrigation to increase pollination and thus increase yield of the crop.</p>
<b>Brinjal</b>	Flowering/fruiting	<p><b>Sucking pest</b> →</p> <p><b>Shoot &amp; Fruit borer</b></p>  <p>Mites</p>	<p><b>Monitor the crop for infestation of white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised. To control shoot and fruit borer in brinjal and tomato crops, infested fruits and shoots should be collected and buried inside the soil. If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</b></p> <p><b>If infestation of miteis found aboe ETL in brinjal, then make spraying of Propargite 57 % EC @ 10 ml/ 10 litre of water for the control of Mites under clear sky.</b></p>
<b>Chili</b>	Flowering & fruiting	<p><b>Thrips</b> →</p> 	<p><b>Under present weather conditions severe attack of thrips has been reported. Spray profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</b></p>
<b>Mango</b>	Fruit development	Fruit fly	<p>Spray mixture of DDVP 4 ml and Methyleugenol 10 ml in 10 litres of water for control of fruit fly.</p> <p>Apply irrigation as per need.</p>

<p><b>Lemon</b></p>	<p>Fruit development</p>	<p><b>Mite</b></p> 	<p>Due to high temperature mites infestation in lemon orchard observed. To control mite infestation spray propargite 57 % EC 10/10 liter of water.</p>  <p>To prevent cracking of fruits apply irrigation in time at regular interval.</p>
<p><b>Livestock &amp; poultry</b></p> 	<p>Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day. Do Vaccination for <b>Haemorrhagic septicemia</b> (H.S.) disease and <b>Blue Quarter</b> (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.</p>		

**Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.**



**Gramin Krushi Mausam Seva**  
**Main Dry Farming Research Station**  
**Junagadh Agricultural University**  
**Targhadia (Rajkot) - 360 003**  
**Phone: (0281)2784260(O), 2331686(R)**  
**Fax : ( 0281)2784722, Cell :9426938235**



**Agro met Advisory Service bulletin No.-12 for the Jamnagar district**

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 09 to 13-May - 2020)

**Weather forecast until 08-30 hrs. of 13/05/2020**




**(3) JAMNAGAR DISTRICT:**



	09/05/20	10/05/20	11/05/20	12/05/20	13/05/20
Rainfall (mm)	0	0	0	0	0
Max. Temp.(°C)	38	38	39	38	38
Min. Temp.(°C)	26	26	26	25	25
Total CC (octa)	0	0	0	0	0
RH-I (%)	88	88	88	88	89
RH-II (%)	53	51	49	49	51
Wind Speed (kmph)	13	11	11	12	13
Wind Direction(deg.)	255	256	273	266	270




**Agro meteorological Advisories:-**

<b>General Advice</b>	<ul style="list-style-type: none"><li>• <b>Rain fall: Dry, hot and clear weather forecasted next five days</b></li><li>• <b>Temperature:</b> Maximum and minimum temperature likely to be in range of 38-39°C and 25-26°C, respectively.</li><li>• <b>Relative Humidity:</b> The RH-I and RH-II is likely to be in the range of 88-89% &amp; 49-53, respectively.</li><li>• <b>Wind Direction and Speed:</b> Direction and speed of wind is likely to be W and 11-13 km/h, respectively.</li><li>• <b>Keeping in view of threat of COVID-19 spread farmers are advised to follow the government guidelines, personal hygiene, use mask, ensure hand washing with soap at reasonable interval, social distancing of 6 feet during harvesting and other agricultural operation in all crops.</b></li><li>• <b>Keep social distancing of 5-6 feet in selling of agricultural produces in Marketing yard and also during buying of agricultural inputs.</b></li><li>• <b>Weather outlook for subsequent week: Dry, hot and clear weather very likely to observe during 14-17 May-2020.</b></li><li>• <b>Due to prevailing dry and hot weather conditions, farmers are advised for light irrigation in all vegetable crops. Irrigation should be done in the morning hours</b></li></ul>
-----------------------	---



	<p><b>or in the evening hours.</b></p> <ul style="list-style-type: none"> <li>• Under hot weather a. Avoid heat exposure. b. Wear light weight, light coloured, loose cotton clothes. c. Cover your head. d. Keep the animal under tree shade or shed during noon hours.</li> <li>• Deep ploughing of fields after harvesting of rabi crops is advised so that eggs and pupating stages of insects, fungus, bacteria and weed seeds are destroyed.</li> </ul>		
Crops	Stage	Insect/Pest/ Variety	Advisory
<b>Farm compost/ FYM</b>	Turning of well rotted compost & FYM	-	<p>Dig out completely composted material made from animal dung, urine, litter and farm waste.</p> 
<b>Summer Groundnut</b>	Pod formation/development 	<b>Red Mite</b> 	<p>Due to hot weather monitor the crop against the attack of mite in groundnut is advised. If population is found more then spraying of Ethion @ 1.5-2 ml <b>or</b> Propargite 57 % EC @ 10 ml /litre of water is advised.</p> <p>Apply light irrigation at pod formation and development stages.</p>
<b>Summer Sesame</b>	Physiological Maturity	-	Harvest in time at ripening stage before seed shattering.
<b>Green gram &amp; Black gram</b>	Flowering & pod formation	Heliothis	Under present weather condition infestation of heliothis and whiteflies were observed in summer sown green gram & black gram. To control heliothis spray spinosad 3 ml/10 lit of water with 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml / 10 lit of water.
<b>Drum stick/ Moringa</b>	Flowering & fruiting	Bud worm & leaf caterpillar	<p><b>Monitor the crop for infestation of Bud worm &amp; leaf caterpillar, if require spray 500 ml of 5 % neem seed extract (Azadirachtin) or neem seed oil (Azadirachtin 1500 ppm) 50 ml with / 10 lit of water.</b></p> <p><b>Use light trap 1-2/ha to attract &amp; kill the adult.</b></p> <p><b>Under heavy infestation make spray of Malathion 50 EC 20 ml/10 litre of water.</b></p>

<b>Lady finger</b>	Flowering/fruiting	Mites	<p>Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised.</p> <p>Due to prevailing high temperature, light irrigation at short interval is advised.</p> <p>After harvesting of mature okra, urea should be applied @ 15-20 kg/ha.</p>
<b>Ridge gourd, Bottle gourd, Sponge guard, Bitter gourd, Cluster bean, Water melon</b>	Flowering & fruiting	<p>White flies/</p> <p>Mites</p> <p>Downy mildew</p>	<p>In all vegetable crops whitefly infestation observed above ETL. Spray 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml with acetamiprid 20 % SP@ 5.0 gm or <b>imidacloprid 30.5 SC</b> 3 ml/10 lit of water for whiteflies control.</p> <p>If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites in cucurbitaceous .</p> <p>For the control of downy mildew spray mencozeb 40g/pump.</p> <p>Optimum moisture level should be maintained in cucurbitaceous crops by light and frequent irrigation to increase pollination and thus increase yield of the crop.</p>
<b>Brinjal</b>	Flowering/fruiting	<p><b>Sucking pest</b> →</p> <p><b>Shoot &amp; Fruit borer</b></p>  <p>Mites</p>	<p><b>Monitor the crop for infestation of white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised. To control shoot and fruit borer in brinjal and tomato crops, infested fruits and shoots should be collected and buried inside the soil. If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</b></p> <p><b>If infestation of miteis found aboe ETL in brinjal, then make spraying of Propargite 57 % EC @ 10 ml/ 10 litre of water for the control of Mites under clear sky.</b></p>
<b>Chili</b>	Flowering & fruiting	<p><b>Thrips</b> →</p> 	<p><b>Under present weather conditions severe attack of thrips has been reported. Spray profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</b></p>
<b>Mango</b>	Fruit development	Fruit fly	<p>Spray mixture of DDVP 4 ml and Methyleugenol 10 ml in 10 litres of water for control of fruit fly.</p> <p>Apply irrigation as per need.</p>

<p><b>Lemon</b></p>	<p>Fruit development</p>	<p><b>Mite</b></p> 	<p>Due to high temperature mites infestation in lemon orchard observed. To control mite infestation spray propargite 57 % EC 10/10 liter of water.</p>  <p>To prevent cracking of fruits apply irrigation in time at regular interval.</p>
<p><b>Livestock &amp; poultry</b></p> 	<p>Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day. Do Vaccination for <b>Haemorrhagic septicemia</b> (H.S.) disease and <b>Blue Quarter</b> (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.</p>		

**Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.**



**Gramin Krushi Mausam Seva**  
**Main Dry Farming Research Station**  
**Junagadh Agricultural University**  
**Targhadia (Rajkot) - 360 003**  
**Phone: (0281)2784260(O), 2331686(R)**  
**Fax : ( 0281)2784722, Cell :9426938235**



**Agro met Advisory Service bulletin No.-12 for the Devbhumi Dwarka district**

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 09 to 13-May - 2020)

**Weather forecast until 08-30 hrs. of 13/05/2020**




**(4) DEVBHUMI DWARKA DISTRICT:**



	09/05/20	10/05/20	11/05/20	12/05/20	13/05/20
Rainfall (mm)	0	0	0	0	0
Max. Temp.(°C)	32	33	32	32	31
Min. Temp.(°C)	27	27	26	26	27
Total CC (octa)					
RH-I (%)					
RH-II (%)					
Wind Speed (kmph)					
Wind Direction(deg.)					




**Agro meteorological Advisories:-**

**General Advice**

- **Rain fall: Dry, hot weather forecasted next five days**
- **Temperature:** Maximum and minimum temperature likely to be in range of 31-33°C and 26-27°C, respectively.
- **Keeping in view of threat of COVID-19 spread farmers are advised to follow the government guidelines, personal hygiene, use mask, ensure hand washing with soap at reasonable interval, social distancing of 6 feet during harvesting and other agricultural operation in all crops.**
- **Keep social distancing of 5-6 feet in selling of agricultural produces in Marketing yard and also during buying of agricultural inputs.**
- **Weather outlook for subsequent week: Dry, hot and clear weather very likely to observe during 14-17 May-2020.**
- **Due to prevailing dry and hot weather conditions, farmers are advised for light irrigation in all vegetable crops. Irrigation should be done in the morning hours or in the evening hours.**
- **Under hot weather a. Avoid heat exposure. b. Wear light weight, light coloured, loose cotton clothes. c. Cover your head. d. Keep the animal under tree shade or shed during noon hours.**
- **Deep ploughing of fields after harvesting of rabi crops is advised so that eggs and**

pupating stages of insects, fungus, bacteria and weed seeds are destroyed.			
Crops	Stage	Insect/Pest/ Variety	Advisory
<b>Farm compost/ FYM</b>	Turning of well rotted compost & FYM	-	Dig out completely composted material made from animal dung, urine, litter and farm waste. 
<b>Summer Groundnut</b>	Pod formation/development 	<b>Red Mite</b> 	<b>Due to hot weather monitor the crop against the attack of mite in groundnut is advised. If population is found more then spraying of Ethion @ 1.5-2 ml or Propargite 57 % EC @ 10 ml /litre of water is advised.</b> Apply light irrigation at pod formation and development stages.
<b>Summer Sesame</b>	Physiological Maturity	-	Harvest in time at ripening stage before seed shattering.
<b>Green gram &amp; Black gram</b>	Flowering & pod formation	Heliothis	Under present weather condition infestation of heliothis and whiteflies were observed in summer sown green gram & black gram. To control heliothis spray spinosad 3 ml/10 lit of water with 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml / 10 lit of water.
<b>Drum stick/ Moringa</b>	Flowering & fruiting	Bud worm & leaf caterpillar	<b>Monitor the crop for infestation of Bud worm &amp; leaf caterpillar, if require spray 500 ml of 5 % neem seed extract (Azadirachtin) or neem seed oil (Azadirachtin 1500 ppm) 50 ml with / 10 lit of water.</b> <b>Use light trap 1-2/ha to attract &amp; kill the adult.</b> <b>Under heavy infestation make spray of Malathion 50 EC 20 ml/10 litre of water.</b>
<b>Lady finger</b>	Flowering/fruiting	Mites	Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised. Due to prevailing high temperature, light irrigation

			at short interval is advised. After harvesting of mature okra, urea should be applied @ 15-20 kg/ha.
<b>Ridge gourd, Bottle gourd, Sponge guard, Bitter gourd, Cluster bean, Water melon</b>	Flowering & fruiting	White flies/  Mites  Downy mildew	In all vegetable crops whitefly infestation observed above ETL. Spray 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml with acetamiprid 20 % SP@ 5.0 gm or <b>imidacloprid 30.5 SC</b> 3 ml/10 lit of water for whiteflies control. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites in cucurbiteous . For the control of downy mildew spray mencozeb 40g/pump. Optimum moisture level should be maintained in cucurbitaceous crops by light and frequent irrigation to increase pollination and thus increase yield of the crop.
<b>Brinjal</b>	Flowering/ fruiting	<b>Sucking pest</b> →  <b>Shoot &amp; Fruit borer</b>    Mites	<b>Monitor the crop for infestation of white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised. To control shoot and fruit borer in brinjal and tomato crops, infested fruits and shoots should be collected and buried inside the soil. If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</b>  <b>If infestation of mites found aboe ETL in brinjal, then make spraying of Propargite 57 % EC @ 10 ml/ 10 litre of water for the control of Mites under clear sky.</b>
<b>Chili</b>	Flowering & fruiting	<b>Thrips</b> → 	<b>Under present weather conditions severe attack of thrips has been reported. Spray profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</b>
<b>Mango</b>	Fruit development	Fruit fly	Spray mixture of DDVP 4 ml and Methyleugenol 10 ml in 10 litres of water for control of fruit fly. Apply irrigation as per need.

<p><b>Lemon</b></p>	<p>Fruit development</p>	<p><b>Mite</b></p> 	<p>Due to high temperature mites infestation in lemon orchard observed. To control mite infestation spray propargite 57 % EC 10/10 liter of water.</p>  <p>To prevent cracking of fruits apply irrigation in time at regular interval.</p>
<p><b>Livestock &amp; poultry</b></p> 	<p>Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day. Do Vaccination for <b>Haemorrhagic septicemia</b> (H.S.) disease and <b>Blue Quarter</b> (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.</p>		

**Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.**



**Gramin Krushi Mausam Seva**  
**Main Dry Farming Research Station**  
**Junagadh Agricultural University**  
**Targhadia (Rajkot) - 360 003**  
**Phone: (0281)2784260(O), 2331686(R)**  
**Fax : ( 0281)2784722, Cell :9426938235**



**Agro met Advisory Service bulletin No.-12 for the Amreli district**

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 09 to 13-May - 2020)

**Weather forecast until 08-30 hrs. of 13/05/2020**




**(5) AMRELI DISTRICT:**



	09/05/20	10/05/20	11/05/20	12/05/20	13/05/20
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	44	43	43	43	43
Min.Temp.(°C)	25	25	24	25	25
Total CC (octa)	0	0	0	0	0
RH-I (%)	78	74	74	75	76
RH-II (%)	34	33	31	32	30
Wind Speed (kmph)	10	9	9	7	8
Wind Direction(deg.)	288	300	303	272	300




**Agro meteorological Advisories:**

<b>General Advice</b>	<ul style="list-style-type: none"><li>• <b>Rain fall: Dry, hot and clear weather forecasted next five days</b></li><li>• <b>Temperature:</b> Maximum and minimum temperature likely to be in range of 43-44°C and 24-25°C, respectively.</li><li>• <b>Relative Humidity:</b> The RH-I and RH-II is likely to be in the range of 74-78% &amp; 30-34, respectively.</li><li>• <b>Wind Direction and Speed:</b> Direction and speed of wind is likely to be W, NW and 7-10 km/h, respectively.</li><li>• <b>Keeping in view of threat of COVID-19 spread farmers are advised to follow the government guidelines, personal hygiene, use mask, ensure hand washing with soap at reasonable interval, social distancing of 6 feet during harvesting and other agricultural operation in all crops.</b></li><li>• <b>Keep social distancing of 5-6 feet in selling of agricultural produces in Marketing yard and also during buying of agricultural inputs.</b></li><li>• <b>Weather outlook for subsequent week: Dry, hot and clear weather very likely to observe during 14-17 May-2020.</b></li><li>• <b>Due to prevailing dry and hot weather conditions, farmers are advised for light irrigation in all vegetable crops. Irrigation should be done in the morning hours</b></li></ul>
-----------------------	--



	<p><b>or in the evening hours.</b></p> <ul style="list-style-type: none"> <li><b>Under hot weather a. Avoid heat exposure. b. Wear light weight, light coloured, loose cotton clothes. c. Cover your head. d. Keep the animal under tree shade or shed during noon hours.</b></li> <li><b>Deep ploughing of fields after harvesting of rabi crops is advised so that eggs and pupating stages of insects, fungus, bacteria and weed seeds are destroyed.</b></li> </ul>		
<b>Crops</b>	<b>Stage</b>	<b>Insect/Pest/ Variety</b>	<b>Advisory</b>
<b>Farm compost/ FYM</b>	Turning of well rotted compost & FYM	-	<p>Dig out completely composted material made from animal dung, urine, litter and farm waste.</p> 
<b>Summer Groundnut</b>	Pod formation/development  	<b>Red Mite</b>  	<p><b>Due to hot weather monitor the crop against the attack of mite in groundnut is advised. If population is found more then spraying of Ethion @ 1.5-2 ml or Propargite 57 % EC @ 10 ml /litre of water is advised.</b></p> <p>Apply light irrigation at pod formation and development stages.</p>
<b>Summer Sesame</b>	Physiological Maturity	-	Harvest in time at ripening stage before seed shattering.
<b>Green gram &amp; Black gram</b>	Flowering & pod formation	Heliothis	Under present weather condition infestation of heliothis and whiteflies were observed in summer sown green gram & black gram. To control heliothis spray spinosad 3 ml/10 lit of water with 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml / 10 lit of water.
<b>Drum stick/ Moringa</b>	Flowering & fruiting	Bud worm & leaf caterpillar	<p><b>Monitor the crop for infestation of Bud worm &amp; leaf caterpillar, if require spray 500 ml of 5 % neem seed extract (Azadirachtin) or neem seed oil (Azadirachtin 1500 ppm) 50 ml with / 10 lit of water.</b></p> <p><b>Use light trap 1-2/ha to attract &amp; kill the adult.</b></p> <p><b>Under heavy infestation make spray of Malathion 50 EC 20 ml/10 litre of water.</b></p>

<b>Lady finger</b>	Flowering/fruiting	Mites	<p>Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised.</p> <p>Due to prevailing high temperature, light irrigation at short interval is advised.</p> <p>After harvesting of mature okra, urea should be applied @ 15-20 kg/ha.</p>
<b>Ridge gourd, Bottle gourd, Sponge guard, Bitter gourd, Cluster bean, Water melon</b>	Flowering & fruiting	<p>White flies/</p> <p>Mites</p> <p>Downy mildew</p>	<p>In all vegetable crops whitefly infestation observed above ETL. Spray 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml with acetamiprid 20 % SP@ 5.0 gm or <b>imidacloprid 30.5 SC</b> 3 ml/10 lit of water for whiteflies control.</p> <p>If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites in cucurbitaceous .</p> <p>For the control of downy mildew spray mencozeb 40g/pump.</p> <p>Optimum moisture level should be maintained in cucurbitaceous crops by light and frequent irrigation to increase pollination and thus increase yield of the crop.</p>
<b>Brinjal</b>	Flowering/fruiting	<p><b>Sucking pest</b> →</p> <p><b>Shoot &amp; Fruit borer</b></p>  <p>Mites</p>	<p><b>Monitor the crop for infestation of white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised. To control shoot and fruit borer in brinjal and tomato crops, infested fruits and shoots should be collected and buried inside the soil. If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</b></p> <p><b>If infestation of miteis found aboe ETL in brinjal, then make spraying of Propargite 57 % EC @ 10 ml/ 10 litre of water for the control of Mites under clear sky.</b></p>
<b>Chili</b>	Flowering & fruiting	<p><b>Thrips</b> →</p> 	<p><b>Under present weather conditions severe attack of thrips has been reported. Spray profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</b></p>
<b>Mango</b>	Fruit development	Fruit fly	<p>Spray mixture of DDVP 4 ml and Methyleugenol 10 ml in 10 litres of water for control of fruit fly.</p> <p>Apply irrigation as per need.</p>

<p><b>Lemon</b></p>	<p>Fruit development</p>	<p><b>Mite</b></p> 	<p>Due to high temperature mites infestation in lemon orchard observed. To control mite infestation spray propargite 57 % EC 10/10 liter of water.</p>  <p>To prevent cracking of fruits apply irrigation in time at regular interval.</p>
<p><b>Livestock &amp; poultry</b></p> 	<p>Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day. Do Vaccination for <b>Haemorrhagic septicemia</b> (H.S.) disease and <b>Blue Quarter</b> (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.</p>		

**Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.**



**Gramin Krushi Mausam Seva**  
**Main Dry Farming Research Station**  
**Junagadh Agricultural University**  
**Targhadia (Rajkot) - 360 003**  
**Phone: (0281)2784260(O), 2331686(R)**  
**Fax : (0281)2784722, Cell:9426938235**



**Agro met Advisory Service bulletin No.-12 for the Surendranagar district**

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 09 to 13-May - 2020)

**Weather forecast until 08-30 hrs of 13/05/2020**




**(6) SURENDRANAGAR DISTRICT:**



	09/05/20	10/05/20	11/05/20	12/05/20	13/05/20
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	44	43	43	43	43
Min.Temp.(°C)	27	27	26	27	27
Total CC (octa)	0	0	0	1	0
RH-I (%)	74	72	74	77	77
RH-II (%)	27	25	25	24	24
Wind Speed (kmph)	15	13	14	14	14
Wind Direction(deg.)	263	273	278	279	268




**Agro meteorological advisories:**

**General Advice**

- **Rain fall: Dry, hot and clear weather forecasted next five days**
- **Temperature:** Maximum and minimum temperature likely to be in range of 43-44°C and 26-27°C, respectively.
- **Relative Humidity:** The RH-I and RH-II is likely to be in the range of 72-77% & 24-27, respectively.
- **Wind Direction and Speed:** Direction and speed of wind is likely to be W and 13-15 km/h, respectively.
- **Keeping in view of threat of COVID-19 spread farmers are advised to follow the government guidelines, personal hygiene, use mask, ensure hand washing with soap at reasonable interval, social distancing of 6 feet during harvesting and other agricultural operation in all crops.**
- **Keep social distancing of 5-6 feet in selling of agricultural produces in Marketing yard and also during buying of agricultural inputs.**
- **Weather outlook for subsequent week: Dry, hot and clear weather very likely to observe during 14-17 May-2020.**
- **Due to prevailing dry and hot weather conditions, farmers are advised for light irrigation in all vegetable crops. Irrigation should be done in the morning hours**

	<p><b>or in the evening hours.</b></p> <ul style="list-style-type: none"> <li>• Under hot weather a. Avoid heat exposure. b. Wear light weight, light coloured, loose cotton clothes. c. Cover your head. d. Keep the animal under tree shade or shed during noon hours.</li> <li>• Deep ploughing of fields after harvesting of rabi crops is advised so that eggs and pupating stages of insects, fungus, bacteria and weed seeds are destroyed.</li> </ul>		
Crops	Stage	Insect/Pest/ Variety	Advisory
<b>Farm compost/ FYM</b>	Turning of well rotted compost & FYM	-	<p>Dig out completely composted material made from animal dung, urine, litter and farm waste.</p> 
<b>Summer Groundnut</b>	Pod formation/development  	<b>Red Mite</b>  	<p>Due to hot weather monitor the crop against the attack of mite in groundnut is advised. If population is found more then spraying of Ethion @ 1.5-2 ml <b>or</b> Propargite 57 % EC @ 10 ml /litre of water is advised.</p> <p>Apply light irrigation at pod formation and development stages.</p>
<b>Summer Sesame</b>	Physiological Maturity	-	Harvest in time at ripening stage before seed shattering.
<b>Green gram &amp; Black gram</b>	Flowering & pod formation	Heliothis	Under present weather condition infestation of heliothis and whiteflies were observed in summer sown green gram & black gram. To control heliothis spray spinosad 3 ml/10 lit of water with 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml / 10 lit of water.
<b>Drum stick/ Moringa</b>	Flowering & fruiting	Bud worm & leaf caterpillar	<p><b>Monitor the crop for infestation of Bud worm &amp; leaf caterpillar, if require spray 500 ml of 5 % neem seed extract (Azadirachtin) or neem seed oil (Azadirachtin 1500 ppm) 50 ml with / 10 lit of water.</b></p> <p><b>Use light trap 1-2/ha to attract &amp; kill the adult.</b></p> <p><b>Under heavy infestation make spray of Malathion 50 EC 20 ml/10 litre of water.</b></p>

<b>Lady finger</b>	Flowering/fruiting	Mites	<p>Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised.</p> <p>Due to prevailing high temperature, light irrigation at short interval is advised.</p> <p>After harvesting of mature okra, urea should be applied @ 15-20 kg/ha.</p>
<b>Ridge gourd, Bottle gourd, Sponge guard, Bitter gourd, Cluster bean, Water melon</b>	Flowering & fruiting	<p>White flies/</p> <p>Mites</p> <p>Downy mildew</p>	<p>In all vegetable crops whitefly infestation observed above ETL. Spray 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml with acetamiprid 20 % SP@ 5.0 gm or <b>imidacloprid 30.5 SC</b> 3 ml/10 lit of water for whiteflies control.</p> <p>If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites in cucurbitaceous .</p> <p>For the control of downy mildew spray mencozeb 40g/pump.</p> <p>Optimum moisture level should be maintained in cucurbitaceous crops by light and frequent irrigation to increase pollination and thus increase yield of the crop.</p>
<b>Brinjal</b>	Flowering/fruiting	<p><b>Sucking pest</b> →</p> <p><b>Shoot &amp; Fruit borer</b></p>  <p>Mites</p>	<p><b>Monitor the crop for infestation of white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised. To control shoot and fruit borer in brinjal and tomato crops, infested fruits and shoots should be collected and buried inside the soil. If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</b></p> <p><b>If infestation of miteis found aboe ETL in brinjal, then make spraying of Propargite 57 % EC @ 10 ml/ 10 litre of water for the control of Mites under clear sky.</b></p>
<b>Chili</b>	Flowering & fruiting	<p><b>Thrips</b> →</p> 	<p><b>Under present weather conditions severe attack of thrips has been reported. Spray profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</b></p>
<b>Mango</b>	Fruit development	Fruit fly	<p>Spray mixture of DDVP 4 ml and Methyleugenol 10 ml in 10 litres of water for control of fruit fly.</p> <p>Apply irrigation as per need.</p>

<p><b>Lemon</b></p>	<p>Fruit development</p>	<p><b>Mite</b></p> 	<p>Due to high temperature mites infestation in lemon orchard observed. To control mite infestation spray propargite 57 % EC 10/10 liter of water.</p>  <p>To prevent cracking of fruits apply irrigation in time at regular interval.</p>
<p><b>Livestock &amp; poultry</b></p> 	<p>Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day. Do Vaccination for <b>Haemorrhagic septicemia</b> (H.S.) disease and <b>Blue Quarter</b> (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.</p>		

**Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.**